

Topics in algorithmic randomness and computable analysis

Jason Rute

August 2013

Carnegie Mellon University

Thesis committee:

Jeremy Avigad (Advisor)

James Cummings

Richard Statman

Bjørn Kjos-Hanssen

Submitted in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

Abstract

This dissertation develops connections between algorithmic randomness and computable analysis.

In the first part, it is shown that computable randomness can be defined robustly on all computable probability spaces, and that computable randomness is preserved by a.e. computable isomorphisms between spaces. Further applications are also given.

In the second part, a number of almost-everywhere convergence theorems are looked at using computable analysis and algorithmic randomness. These include various martingale convergence theorems and almost-everywhere differentiability theorems. General conditions are given for when the rate of convergence is computable and for when convergence takes place on the Schnorr random points. Examples are provided to show that these almost-everywhere convergence theorems characterize Schnorr randomness.

Acknowledgements

I am grateful to my family. Their unconditional support over the years gave me the courage to follow my dreams and to work hard at attaining them.

Jeremy Avigad has been all I could ask for in an advisor, and even more. I am incredibly lucky for having ending up under his direction.

The computability theory community has been incredibly supportive, even when I was just starting out. This especially includes Laurent Bienvenu, Johanna Franklin, Bjørn Kjos-Hanssen, Mathieu Hoyrup, Joseph Miller, André Nies, Jan Reimann, and Stephan Simpson. Their encouragement, proof reading, dinner conversations, advice, and invitations to visit have been invaluable to me.

Lastly, I am indebted to Marla Slusky for her patience and encouragement. She was literally cheering me on in the last few weeks of writing my dissertation.

Contents

Introduction.....	1
Computable randomness and betting for computable probability spaces.....	3
Algorithmic randomness, martingales and differentiability	34

Introduction

Algorithmic randomness is a branch of computability theory which is concerned with the properties of objects which behave randomly with respect to computable statistical tests. While there are many notions of “computable statistical test”, and therefore many notions of randomness, the majority of the papers written on algorithmic randomness have focused on infinite binary sequences $2^{\mathbb{N}}$ under the coin-flipping (or Lebesgue) probability measure λ . For example the recent monographs of Downey and Hirschfeldt [3] and Nies [9] almost entirely focus on $(2^{\mathbb{N}}, \lambda)$.

However, a number of researchers have started to look at randomness on other spaces and other measures. This includes work by Levin [8] on Martin-Löf randomness for other probability measures, as well as the work by Fouché [4] on Martin-Löf random Brownian motion. Hoyrup and Rojas [7] have done much to give a robust theory of Martin-Löf randomness on other spaces.

Parallel to this, a number of researchers have taken an interest in the computability of measure theory and probability. This goes back to the constructive works of Šanin [11] and Bishop [1]. More recently, it was discovered that many almost-everywhere theorems in analysis characterize the most common randomness notions [12, 6, 10, 5, 2]. This is an exciting new direction connecting computable analysis and algorithmic randomness.

This dissertation contributes to this direction. It is made up of two chapters. Each can be read independently of the other.

The first chapter, “Computable randomness and betting for computable probability spaces”, gives a definition of computable randomness on other computable probability spaces. Unlike other common randomness notions, it is not completely obvious how to define computable randomness on other spaces. I give a number of examples showing that my definition is the correct one.

The second chapter, “Algorithmic randomness, martingales and differentiability”, gives new results in the computability of a.e. convergence results. In particular, I show that under certain conditions, most a.e. convergence theorems characterize Schnorr randomness. I also develop a general theory of measurable functions and Schnorr randomness which I hope will be of use to other researchers.

References

- [1] Errett Bishop. *Foundations of constructive analysis*. McGraw-Hill Book Co., New York, 1967.
- [2] Vasco Brattka, Joseph S. Miller, and André Nies. Randomness and differentiability. *Submitted*.
- [3] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010.
- [4] Willem L. Fouché. The descriptive complexity of Brownian motion. *Adv. Math.*, 155(2):317–343, 2000.
- [5] Peter Gács, Mathieu Hoyrup, and Cristóbal Rojas. Randomness on computable probability spaces—a dynamical point of view. *Theory Comput. Syst.*, 48(3):465–485, 2011.
- [6] Mathieu Hoyrup and Cristóbal Rojas. Applications of effective probability theory to Martin-Löf randomness. In *Automata, languages and programming. Part I*, volume 5555 of *Lecture Notes in Comput. Sci.*, pages 549–561. Springer, Berlin, 2009.
- [7] Mathieu Hoyrup and Cristóbal Rojas. Computability of probability measures and Martin-Löf randomness over metric spaces. *Inform. and Comput.*, 207(7):830–847, 2009.

- [8] L. A. Levin. Uniform tests for randomness. *Dokl. Akad. Nauk SSSR*, 227(1):33–35, 1976.
- [9] André Nies. *Computability and randomness*, volume 51 of *Oxford Logic Guides*. Oxford University Press, Oxford, 2009.
- [10] Noopur Pathak. A computational aspect of the Lebesgue differentiation theorem. *J. Log. Anal.*, 1(9):15, 2009.
- [11] N. A. Šanin. *Constructive real numbers and constructive function spaces*. Translated from the Russian by E. Mendelson. Translations of Mathematical Monographs, Vol. 21. American Mathematical Society, Providence, R.I., 1968.
- [12] V. V. V'yugin. Ergodic theorems for individual random sequences. *Theoret. Comput. Sci.*, 207(2):343–361, 1998.

COMPUTABLE RANDOMNESS AND BETTING FOR COMPUTABLE PROBABILITY SPACES

ABSTRACT. Unlike Martin-Löf randomness and Schnorr randomness, computable randomness has not been defined, except for a few ad hoc cases, outside of Cantor space. This paper offers such a definition (actually, many equivalent definitions), and further, provides a general method for abstracting “bit-wise” definitions of randomness from Cantor space to arbitrary computable probability spaces. This same method is also applied to give machine characterizations of computable and Schnorr randomness for computable probability spaces, extending the previous known results. This paper also addresses “Schnorr’s Critique” that gambling characterizations of Martin-Löf randomness are not computable enough. The paper contains a new type of randomness—endomorphism randomness—which the author hopes will shed light on the open question of whether Kolmogorov-Loveland randomness is equivalent to Martin-Löf randomness. It ends with ideas on how to extend this work to layerwise-computable structures, non-computable probability spaces, computable topological spaces, and measures defined by π -systems. It also ends with a possible definition of K-triviality for computable probability spaces.

CONTENTS

1. Introduction	3
2. Computable randomness on 2^ω	5
3. Computable probability spaces and algorithmic randomness	6
4. Almost-everywhere decidable cell decompositions	8
5. Computable randomness on computable probability spaces	10
6. Machine characterizations of computable and Schnorr randomness	14
7. Computable randomness and isomorphisms	15
8. Generalizing randomness to computable probability spaces	19
9. Betting strategies and Kolmogorov-Loveland randomness	20
10. Endomorphism randomness	24
11. Further directions	26
References	27

1. INTRODUCTION

The subjects of measure theory and probability are filled with a number of theorems stating that some property holds “almost everywhere” or “almost surely.” Informally, these theorems state that if one starts with a random point, then the desired result is true. The field of algorithmic randomness has been very successful in making this notion formal: by restricting oneself to computable tests for non-randomness, one can achieve a measure-one set of points that behave as desired. The most prominent such notion of randomness is Martin-Löf randomness. However, Schnorr [35] gave an argument—which is now known as Schnorr’s Critique—that Martin-Löf randomness does not have a sufficiently computable characterization. He offered two weaker-but-more-computable alternatives: Schnorr randomness and computable randomness. All three randomness notions are interesting and robust, and further each has been closely linked to computable analysis (for example [12, 18, 33, 38]).

Computable randomness, however, is the only one of the three that has not been defined for arbitrary computable probability spaces. The usual definition is specifically for Cantor space (i.e. the space 2^ω of infinite binary strings), or by analogy, spaces such as 3^ω . Namely, a string $x \in 2^\omega$ is said to be computably random (in the fair-coin measure) if, roughly speaking, one cannot win arbitrarily large amounts of money using a computable betting strategy to gamble on the bits of x . (See Section 2 for a formal definition.)

2010 *Mathematics Subject Classification.* 03D32, 68Q30.

This work has been partially supported by NSF grant DMS1068829.

While it is customary to say a real $x \in [0, 1]$ is computably random if its binary expansion is computably random in 2^ω , it was only recently shown [12] that this is the same as saying that, for example, the ternary expansion of x is computably random in 3^ω . In other words, computable randomness is base invariant.

In this paper, I use a method for extending the “bit-wise” definitions of randomness on Cantor space to arbitrary computable probability spaces. The method is based on previous methods given by Gács [17] and later Hoyrup and Rojas [22] of dividing a space into cells. However, to successfully extend a randomness notion (such that the new definition agrees with the former on 2^ω), one must show a property similar to base invariance. I do this for computable randomness.

An outline of the paper is as follows. Section 2 defines computable randomness on 2^ω , both for the fair-coin measure and for other computable probability measures on Cantor space. Unlike previous treatments (for example Bienvenu and Merkle [6]) I address the important pathological case where the measure may have null open sets.

Section 3 gives background on computable analysis, computable probability spaces, and algorithmic randomness.

Section 4 presents the concepts of an almost-everywhere decidable set (due to Hoyrup and Rojas [22]) and an a.e. decidable cell decomposition (which is similar to work of Hoyrup and Rojas [22] and Gács [17]). Recall that the topology of 2^ω is generated by the collection of basic open sets of the form $[\sigma]^\prec = \{x \in 2^\omega \mid x \succ \sigma\}$ where $x \succ \sigma$ means σ is an initial segment of x . Further, any Borel measure μ of 2^ω is determined by the values $\mu([\sigma]^\prec)$. The main idea of this paper is that for a computable probability space (\mathcal{X}, μ) one can replace the basic open sets of 2^ω (which are decidable) with an indexed family of “almost-everywhere decidable” sets $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$ which behave in much the same way. I call each such indexed family a cell decomposition of the space. This allows one to effortlessly transfer a definition from Cantor space to any computable probability space.

Section 5 applies this method to computable randomness, giving a variety of equivalent definitions based on martingales and other tests. More importantly, I show this definition is invariant under the choice of cell decomposition. Similar to the base-invariance proof of Brattka, Miller and Nies [12, 37], my proof uses computable analysis. However, their method does not apply here. (Their proof uses differentiability and the fact that every atomless measure on $[0, 1]$ is naturally equivalent to a measure on 2^ω . The situation is more complicated in the general case. One does not have differentiability, and one must consider absolutely-continuous measures instead of mere atomless ones.)

Section 6 gives a machine characterization of computable and Schnorr randomness for computable probability spaces. This combines the machine characterizations of computable randomness and Schnorr randomness (respectively, Mihailović [14, Theorem 7.1.25] and Downey, Griffiths, and LaForte [13]) with the machine characterization of Martin-Löf randomness on arbitrary computable probability spaces (Gács [16] and Hoyrup and Rojas [22]).

Section 7 shows a correspondence between cell decompositions of a computable probability space (\mathcal{X}, μ) and isomorphisms from (\mathcal{X}, μ) to Cantor space. I also show computable randomness is preserved by isomorphisms between computable probability spaces, giving yet another characterization of computable randomness. However, unlike other notions of randomness, computable randomness is not preserved by mere morphisms (almost-everywhere computable, measure-preserving maps).

Section 8 gives three equivalent methods to extend a randomness notion to all computable probability measures. It also gives the conditions for when this new randomness notion agrees with the original one.

Section 9 asks how the method of this paper applies to Kolmogorov-Loveland randomness, another notion of randomness defined by gambling. The result is that the natural extension of Kolmogorov-Loveland randomness to arbitrary computable probability measures is Martin-Löf randomness. However, I do not answer the important open question as to whether Kolmogorov-Loveland randomness and Martin-Löf randomness are equivalent. Nonetheless, I do believe this answers Schnorr’s Critique, namely that Martin-Löf randomness does have a natural definition in terms of computable betting strategies.

Section 10 explores a new notion of randomness in between Martin-Löf randomness and Kolmogorov-Loveland randomness, possibly equal to both. It is called endomorphism randomness.

Last, in Section 11, I suggest ways to generalize the method of this paper to a larger class of isomorphisms and cell decompositions. I also suggest methods for extending computable randomness to a larger class of probability spaces, including non-computable probability spaces, computable topological spaces, and measures defined by π -systems. Drawing on Section 6, I suggest a possible definition of K-triviality for

computable probability spaces. Finally, I ask what can be known about the interplay between randomness, morphisms, and isomorphisms.

2. COMPUTABLE RANDOMNESS ON 2^ω

Before exploring computable randomness on arbitrary computable probability spaces, a useful intermediate step will be to consider computable probability measures on Cantor space.

We fix notation: $2^{<\omega}$ is the space of finite binary strings; 2^ω is the space of infinite binary strings; ε is the empty string; $\sigma \prec \tau$ and $\sigma \prec x$ mean σ is a proper initial segment of $\tau \in 2^{<\omega}$ or $x \in 2^\omega$; and $[\sigma]^\prec = \{x \in 2^\omega \mid \sigma \prec x\}$ is a BASIC OPEN SET or CYLINDER SET. Also for $\sigma \in 2^{<\omega}$ (or $x \in 2^\omega$), $\sigma(n)$ is the n th digit of σ (where $\sigma(0)$ is the “0th” digit) and $\sigma \upharpoonright n = \sigma(0) \cdots \sigma(n-1)$.

Typically, a MARTINGALE (on the fair-coin probability measure) is defined as a function $M: 2^{<\omega} \rightarrow [0, \infty)$ such that the following property holds for each $\sigma \in 2^{<\omega}$: $M(\sigma) = \frac{1}{2}(M(\sigma 0) + M(\sigma 1))$. Such a martingale can be thought of as a betting strategy on coin flips: the gambler starts with the value $M(\varepsilon)$ as her capital (where ε is the empty string) and bets on fair coin flips. Assuming the string σ represents the sequence of coin flips she has seen so far, $M(\sigma 0)$ is the resulting capital she has if the next flip comes up tails, and $M(\sigma 1)$ if heads. A martingale M is said to be COMPUTABLE if the value $M(\sigma)$ is uniformly computable from each σ .

A martingale M is said to SUCCEED on a string $x \in 2^\omega$ if $\limsup_{n \rightarrow \infty} M(x \upharpoonright n) = \infty$ (where $x \upharpoonright n$ is the first n bits of x), i.e. the gambler wins arbitrary large amounts of money using the martingale M while betting on the sequence x of flips. By Kolmogorov’s theorem (see [14, Theorem 6.3.3]), such a martingale can only succeed on a measure-zero set of points. A string $x \in 2^\omega$ is said to be COMPUTABLY RANDOM (on the fair-coin probability measure) if there does not exist a computable martingale M which succeeds on x .

Definition 2.1. A finite Borel measure μ on 2^ω is a COMPUTABLE MEASURE if the measure $\mu([\sigma]^\prec)$ of each basic open set is computable from σ . Further, if $\mu(2^\omega) = 1$, then we say μ is a COMPUTABLE PROBABILITY MEASURE (on 2^ω) and $(2^\omega, \mu)$ is a COMPUTABLE PROBABILITY SPACE (on 2^ω).

In this paper, measure always means a finite Borel measure. When convenient, I will drop the brackets and write $\mu(\sigma)$ instead. By the Carathéodory extension theorem, one may uniquely represent a computable measure as a computable function $\mu: 2^{<\omega} \rightarrow [0, \infty)$ such that

$$\mu(\sigma 0) + \mu(\sigma 1) = \mu(\sigma)$$

for all $\sigma \in 2^{<\omega}$. I will use often confuse a computable measure on 2^ω with its representation on $2^{<\omega}$.

The FAIR-COIN PROBABILITY MEASURE (or the LEBESGUE MEASURE on 2^ω) is the measure λ on 2^ω , defined by

$$\lambda(\sigma) = 2^{-|\sigma|}$$

where $|\sigma|$ is the length of σ . (The Greek letter λ will always be the fair-coin measure on 2^ω , except in a few examples where it is the Lebesgue measure on $[0, 1]^d$ or the uniform measure on 3^ω .)

One may easily generalize the definitions of martingale and computable randomness to a computable probability measure μ . The key idea is that the fairness condition still holds, but is now “weighted” by μ .

Definition 2.2. If μ is a computable probability measure on 2^ω , then a MARTINGALE M (with respect to the measure μ) is a partial function $M: 2^{<\omega} \rightarrow [0, \infty)$ such that the following two conditions hold:

- (1) (Fairness condition) For all $\sigma \in 2^{<\omega}$

$$M(\sigma 0)\mu(\sigma 0) + M(\sigma 1)\mu(\sigma 1) = M(\sigma)\mu(\sigma).$$

- (2) (Impossibility condition) $M(\sigma)$ is defined when $\mu(\sigma) > 0$.

We say M is a COMPUTABLE MARTINGALE if $M(\sigma)$ is uniformly computable from σ (assuming $\mu(\sigma) > 0$).

Definition 2.3. Given a computable probability space $(2^\omega, \mu)$, a martingale M on $(2^\omega, \mu)$ and $x \in 2^\omega$, we say M SUCCEEDS on x if and only if $\limsup_{n \rightarrow \infty} M(x \upharpoonright n) = \infty$. Further, given $x \in 2^\omega$, if x is not in any measure-zero basic open set and there does not exist a computable martingale M on $(2^\omega, \mu)$ which succeeds on x , then we say x is COMPUTABLY RANDOM with respect to the measure μ .

Remark 2.4. The above definitions have been given before by Bienvenu and Merkle [6], and Definition 2.2 is an instance of the more general concept of martingale in probability theory (see for example Williams [40]).

The impossibility condition of Definition 2.2 follows from the slogan in probability theory that a measure-zero (or impossible) event can be ignored. A measure μ such that every open set has measure greater than zero is called a **STRICTLY-POSITIVE** measure. (Bienvenu and Merkle use the term “nowhere vanishing.”) Hence, the impossibility condition is not necessary when μ is strictly positive.

If $(2^\omega, \mu)$ is a strictly-positive probability space, then it is an easy folklore result that there is a bijection between computable martingales M and computable measures ν given by

$$\nu(\sigma) = M(\sigma)\mu(\sigma) \quad \text{and} \quad M(\sigma) = \nu(\sigma)/\mu(\sigma).$$

Even in the case where μ is not strictly positive, the impossibility condition guarantees that these equations can be used to define a computable measure from a computable martingale and vice-versa (under the conditions that $undefined \cdot 0 = 0$ and $x/0 = undefined$ for all x). Further, ν is always computable from M . Indeed, compute $\nu(\sigma)$ by recursion on the length of σ as follows. Since $\mu(\varepsilon) = 1$, $\nu(\varepsilon)$ is computable. To compute, say, $\nu(\sigma 0)$ from $\nu(\sigma)$, use

$$\nu(\sigma 0) = \begin{cases} M(\sigma 0)\mu(\sigma 0) & \text{if } \mu(\sigma 0) > 0 \\ \nu(\sigma) - M(\sigma 1)\mu(\sigma 1) & \text{if } \mu(\sigma 1) > 0 \\ 0 & \text{otherwise} \end{cases}.$$

This is computable, since in the case that $\mu(\sigma) = \mu(\sigma 0) = \mu(\sigma 1) = 0$, the bounds $0 \leq \nu(\sigma 0) \leq \nu(\sigma)$ “squeeze” $\nu(\sigma 0)$ to 0. Conversely, M can be computed from ν by waiting until $\mu(\sigma) > 0$, else $M(\sigma)$ is never defined.

Remark 2.5. It is possible to eliminate the impossibility condition altogether by considering martingales defined on the extended real numbers, i.e. $M: 2^{<\omega} \rightarrow [0, \infty]$. (Use the usual measure-theoretic convention that $\infty \cdot 0 = 0$.) Consider the martingale M_0 defined by $M_0(\sigma) = \lambda(\sigma)/\mu(\sigma)$ where λ is the fair-coin measure. Since, $\lambda(\sigma) > 0$ for all σ , we have that M_0 is computable on the extended real numbers. Notice $M_0(\sigma) = \infty$ if and only if $\mu(\sigma) = 0$, hence one can “forget” the infinite values to get a computable finite-valued martingale M_1 as in Definition 2.2. For any $x \in 2^\omega$, if M_0 succeeds on x then either $\mu(x \upharpoonright n) = 0$ for some n or $M_1(x)$ succeeds on x . In either case, x is not computably random. Conversely, if $x \in 2^\omega$ is not computably random, either M_0 succeeds on x or there is some finite-valued martingale M as in Definition 2 which succeeds on x . In the later case, $N = M + M_0$ is a martingale computable on the extended real numbers which also succeeds on x . However, this paper will only use the finite-valued martingales as in Definition 2.2.

I leave as an open question whether computable randomness can be defined on non-strictly positive measures without the impossibility condition and without infinite values.

Question 2.6. Let μ be a computable probability measure on 2^ω , and assume x is not computably random on μ . Is there necessary a computable martingale $M: 2^{<\omega} \rightarrow [0, \infty)$ with respect to μ which is total, finite-valued and succeeds on x ?

See Downey and Hirschfeldt [14, Section 7.1] and Nies [32, Chapter 7] for more information on computable randomness for $(2^\omega, \lambda)$.

3. COMPUTABLE PROBABILITY SPACES AND ALGORITHMIC RANDOMNESS

In this section I give some background on computable analysis, computable probability spaces, and algorithmic randomness.

3.1. Computable analysis and computable probability spaces. Here I present computable Polish spaces and computable probability spaces. For a more detailed exposition of the same material see Hoyrup and Rojas [22]. This paper assumes some familiarity with basic computability theory and computable analysis, as in Pour El and Richards [34], Weihrauch [39], or Brattka et al. [11].

Definition 3.1. A **COMPUTABLE POLISH SPACE** (or **COMPUTABLE METRIC SPACE**) is a triple (X, d, S) such that

- (1) X is a complete metric space with metric $d: X \times X \rightarrow [0, \infty)$.
- (2) $S = \{a_i\}_{i \in \mathbb{N}}$ is a countable dense subset of X (the **SIMPLE POINTS** of \mathcal{X}).
- (3) The distance $d(a_i, a_j)$ is computable uniformly from i and j .

A point $x \in X$ is said to be **COMPUTABLE** if there is a computable **CAUCHY-NAME** $h \in \mathbb{N}^\omega$ for x , i.e. h is a computable sequence of natural numbers such that $d(a_{h(k)}, x) \leq 2^{-k}$ for all k .

The BASIC OPEN BALLS are sets of the form $B(a, r) = \{x \in X \mid d(x, a) < r\}$ where $a \in S$ and $r > 0$ is rational. The Σ_1^0 sets (EFFECTIVELY OPEN SETS) are computable unions of basic open balls; Π_1^0 sets (EFFECTIVELY CLOSED SETS) are the complements of Σ_1^0 sets; Σ_2^0 sets are computable unions of Π_1^0 sets; and Π_2^0 sets are computable intersections of Σ_1^0 sets. A function $f: \mathcal{X} \rightarrow \mathbb{R}$ is COMPUTABLE (-LY CONTINUOUS) if for each Σ_1^0 set U in \mathbb{R} , the set $f^{-1}(U)$ is Σ_1^0 in \mathcal{X} (uniformly in U), or equivalently, there is an algorithm which sends every Cauchy-name of x to a Cauchy-name of $f(x)$. A function $f: \mathcal{X} \rightarrow [0, \infty]$ is LOWER SEMICOMPUTABLE if it is the supremum of a computable sequence of computable functions $f_n: \mathcal{X} \rightarrow [0, \infty)$.

A real x is said to be LOWER (UPPER) SEMICOMPUTABLE if $\{q \in \mathbb{Q} \mid q < x\}$ (respectively $\{q \in \mathbb{Q} \mid q > x\}$) is a c.e. set.

Definition 3.2. If $\mathcal{X} = (X, d, S)$ is a computable Polish space, then a Borel measure μ is a COMPUTABLE MEASURE on \mathcal{X} if the value $\mu(X)$ is computable, and for each Σ_1^0 set U , the value $\mu(U)$ is lower semicomputable uniformly from the code for U . A COMPUTABLE PROBABILITY SPACE is a pair (\mathcal{X}, μ) where \mathcal{X} is a computable Polish space, μ is a computable measure on \mathcal{X} , and $\mu(\mathcal{X}) = 1$.

While this definition of computable probability space may seem ad hoc, it turns out to be equivalent to a number of other definitions. In particular, the computable probability measures on \mathcal{X} are exactly the computable points in the space of probability measures under the Prokhorov metric. Also, a probability space is computable precisely if the integral operator is a computable operator on computable functions $f: \mathcal{X} \rightarrow [0, 1]$. See Hoyrup and Rojas [22] and Schröder [36] for details.

I will often confuse a metric space or a probability space with its set of points, e.g. writing $x \in \mathcal{X}$ or $x \in (\mathcal{X}, \mu)$ to mean that $x \in X$ where $\mathcal{X} = (X, d, S)$.

3.2. Algorithmic randomness. In this section I give background on algorithmic randomness. Namely, I present three types of tests for Martin-Löf and Schnorr randomness. In Section 5, I will generalize these tests to computable randomness, building off the work of Merkle, Mihailović and Slaman [28] (which is similar to that of Downey, Griffiths and LaForte [13]). I also present Kurtz randomness.

Throughout this section, let (\mathcal{X}, μ) be a computable probability space.

Definition 3.3. A MARTIN-LÖF TEST (with respect to (\mathcal{X}, μ)) is a computable sequence of Σ_1^0 sets (U_n) such that $\mu(U_n) \leq 2^{-n}$ for all n . A SCHNORR TEST is a Martin-Löf test such that $\mu(U_n)$ is also uniformly computable from n . We say x is COVERED BY the test (U_n) if $x \in \bigcap_n U_n$.

Definition 3.4. We say $x \in \mathcal{X}$ is MARTIN-LÖF RANDOM (with respect to (\mathcal{X}, μ)) if there is no Martin-Löf test which covers x . We say x is SCHNORR RANDOM if there is no Schnorr test which covers x . We say x is KURTZ RANDOM (or WEAK RANDOM) if x is not in any null Π_1^0 set (or equivalently a null Σ_2^0 set).

It is easy to see that for all computable probability spaces

$$\text{Martin-Löf} \rightarrow \text{Schnorr} \rightarrow \text{Kurtz}$$

It is also well-known (see [14, 32]) on $(2^\omega, \lambda)$ that

$$(3.1) \quad \text{Martin-Löf} \rightarrow \text{Computable} \rightarrow \text{Schnorr} \rightarrow \text{Kurtz}$$

In the next section, after defining computable randomness for computable probability spaces, I will show (3.1) holds for all computable probability spaces.

In analysis it is common to adopt the slogan “anything that happens on a measure-zero set is negligible.” In this paper it will be useful to adopt the slogan “anything that happens on a measure-zero Σ_2^0 set is negligible,” or in other words, “we do not care about points that are not Kurtz random.” (The reason for this choice will become apparent and is due to the close relationship between Kurtz randomness and a.e. computability. Section 7 contains more discussion.)

Next, I mention two other useful tests.

Definition 3.5. A VITALI TEST (or SOLOVAY TEST) is a sequence of Σ_1^0 sets (U_n) such that $\sum_n \mu(U_n) < \infty$. We say x is VITALI COVERED by (U_n) if $x \in U_n$ for infinitely many n . An INTEGRAL TEST is a lower semicomputable function $g: \mathcal{X} \rightarrow [0, \infty]$ such that $\int g d\mu < \infty$.

Theorem 3.6. For $x \in \mathcal{X}$, the following are equivalent.

- (1) x is Martin-Löf random (respectively Schnorr random).

- (2) x is not Vitali covered by any Vitali test (respectively any Vitali test (U_n) such that $\sum_n \mu(U_n)$ is computable).
- (3) $g(x) < \infty$ for all integral tests g (respectively for all integral tests g such that $\int g d\mu$ is computable).

Remark 3.7. The term Vitali test was coined recently by Nies. For a history of the tests for Schnorr and Martin-Löf randomness see Downey and Hirschfeld [14]. The integral test characterization for Schnorr randomness is due to Miyabe [30] and was also independently communicated to me by Hoyrup and Rojas.

I will give Vitali and integral test characterizations of computable randomness in Section 5.

There are also martingale characterizations of Martin-Löf and Schnorr randomness for 2^ω , but they will not be needed.

4. ALMOST-EVERYWHERE DECIDABLE CELL DECOMPOSITIONS

The main thesis of this paper is that “bit-wise” definitions of randomness for 2^ω , such as computable randomness, can be extended to arbitrary computable probability spaces by replacing the basic open sets $[\sigma]^\prec$ on 2^ω with an indexed family $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$ of a.e. decidable sets. This is the thesis of Hoyrup and Rojas [22]. My method is based off of theirs, although the presentation and definitions differ on a few key points.

Recall that a set $A \subseteq \mathcal{X}$ is DECIDABLE if both A and its complement $\mathcal{X} \setminus A$ are Σ_1^0 sets (equivalently A is both Σ_1^0 and Π_1^0). The intuitive idea is that from the code for any $x \in \mathcal{X}$, one may effectively decide if x is in A or its complement. On 2^ω , the cylinder sets $[\sigma]^\prec$ are decidable. Unfortunately, a space such as $\mathcal{X} = [0, 1]$ has no non-trivial clopen sets, and therefore no non-trivial decidable sets. However, using the idea that null measure sets can be ignored, we can use “almost-everywhere decidable sets” instead.

Definition 4.1 (Hoyrup and Rojas [22]). Let (\mathcal{X}, μ) be a computable probability space. A pair $U, V \subseteq \mathcal{X}$ is a μ -A.E. DECIDABLE PAIR if

- (1) U and V are both Σ_1^0 sets,
- (2) $U \cap V = \emptyset$, and
- (3) $\mu(U \cup V) = 1$.

A set A is a μ -A.E. DECIDABLE SET if there is a μ -a.e. decidable pair U, V such that $U \subseteq A \subseteq \mathcal{X} \setminus V$. The code for the μ -a.e. decidable set A is the pair of codes for the Σ_1^0 sets U and V .

Hoyrup and Rojas [22] also required that $U \cup V$ be dense for technical reasons. We will relax this condition, working under the principle that one can safely ignore null open sets. They also use the terminology “almost decidable set”.

Definition 4.1 is an effectivization of μ -CONTINUITY SET, i.e. a set with μ -null boundary. Notice, the set $\mathcal{X} \setminus (U \cup V)$ includes the topological boundary, but since we do not require $U \cup V$ to be dense, it may also include null open sets.

Not every Σ_1^0 set is a.e. decidable; for example, take a dense open set with measure less than one. However, any basic open ball $B(a, r)$ is a.e. decidable provided that $\{x \mid d(a, x) = r\}$ has null measure. (Again, if we require the boundary to be nowhere dense, the situation is more subtle. See the discussion in Hoyrup and Rojas [22].) Further, the closed ball $\overline{B}(a, r)$ is also a.e. decidable with the same code. Any two a.e. decidable sets with the same code will be considered the same for our purposes. Hence, I will occasionally say $x \in A$ (respectively $x \notin A$), when I mean $x \in U$ (respectively $x \notin V$) for the corresponding a.e. decidable pair (U, V) .

Also notice that if A and B are a.e. decidable, then the Boolean operations $\mathcal{X} \setminus A$, $A \cap B$ and $A \cup B$ are a.e. decidable with codes computable from the codes for A and B .

Definition 4.2 (Inspired by Hoyrup and Rojas [22]). Let (\mathcal{X}, μ) be a computable probability space, and let $\mathcal{A} = (A_i)$ be a uniformly computable sequence of a.e. decidable sets. Let \mathcal{B} be the closure of \mathcal{A} under finite Boolean combinations. We say \mathcal{A} is an (A.E. DECIDABLE) GENERATOR of (\mathcal{X}, μ) if given a Σ_1^0 set $U \subseteq \mathcal{X}$ one can find (effectively from the code of U) a c.e. family $\{B_j\}$ of sets in \mathcal{B} (where $\{B_j\}$ is possibly finite or empty) such that $U = \sum_j B_j$ a.e.

Notice each generator generates the Borel sigma-algebra of \mathcal{X} up to a μ -null set. Hoyrup and Rojas [22] show that not only does such a generator \mathcal{A} exist for each (\mathcal{X}, μ) , but it can be taken to be a basis of the topology, hence they call \mathcal{A} a “basis of almost decidable sets”. I will not require that \mathcal{A} is a basis.

Theorem 4.3 (Hoyrup and Rojas [22]). *Let (\mathcal{X}, μ) be a computable probability space. There exists an a.e. decidable generator \mathcal{A} of (\mathcal{X}, μ) . Further, \mathcal{A} is computable from (\mathcal{X}, μ) .*

The main idea of the proof for Theorem 4.3 is to start with the collection of basic open balls centered at simple points with rational radii. While, these may not have null boundary, a basic diagonalization argument (similar to the proof of the Baire category theory, see [10]) can be used to calculate a set of radii approaching zero for each simple point such that the resulting ball is a.e. decidable. Similar arguments have been given by Bosserhoff [9] and Gács [17]. The technique is related to Bishop's theory of profiles [8, Section 6.4] and to "derandomization" arguments (see Freer and Roy [15] for example).

From a generator we can decompose \mathcal{X} into a.e. decidable cells. This is the indexed family $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$ mentioned in the introduction.

Definition 4.4. Let $\mathcal{A} = (A_i)$ be an a.e. decidable generator of (\mathcal{X}, μ) . Recall each A_i is coded by an a.e. decidable pair (U_i, V_i) where $U_i \subseteq A_i \subseteq \mathcal{X} \setminus V_i$. For $\sigma \in 2^\omega$ of length s define $[\sigma]_{\mathcal{A}} = A_0^{\sigma(0)} \cap A_1^{\sigma(1)} \cap \dots \cap A_{s-1}^{\sigma(s-1)}$ where for each i , $A_i^0 = U_i$ and $A_i^1 = V_i$. When possible, define $x \upharpoonright_{\mathcal{A}} n$ as the unique σ of length n such that $x \in [\sigma]_{\mathcal{A}}$. Also when possible, define the \mathcal{A} -NAME of x as the string $\text{name}_{\mathcal{A}}(x) = \lim_{n \rightarrow \infty} x \upharpoonright_{\mathcal{A}} n$. A point without an \mathcal{A} -name will be called an UNREPRESENTED POINT. Each $[\sigma]_{\mathcal{A}}$ will be called a CELL, and the collection of $\{[\sigma]_{\mathcal{A}}\}_{\sigma \in 2^{<\omega}}$ will be called an (A.E. DECIDABLE) CELL DECOMPOSITION of (\mathcal{X}, μ) .

The choice of notation allows one to quickly translate between Cantor space and the space (\mathcal{X}, μ) . Gács [17] and others refer to the cell $[x \upharpoonright_{\mathcal{A}} n]_{\mathcal{A}}$ as the n -CELL of x and writes it as $\Gamma_n(x)$.

Remark 4.5. There are two types of "bad points", unrepresented points and points $x \in [\sigma]_{\mathcal{A}}$ where $\mu([\sigma]_{\mathcal{A}}) = 0$. The set of "bad points" is a null Σ_2^0 set, so each "bad point" is not even Kurtz random! One may also go further, and for each generator \mathcal{A} compute another \mathcal{A}' such that $[\sigma]_{\mathcal{A}} = [\sigma]_{\mathcal{A}'}$ a.e., but $\mu([\sigma]_{\mathcal{A}}) = 0$ if and only if $[\sigma]_{\mathcal{A}'} = \emptyset$. Then all the "bad points" would be unrepresented points.

Example 4.6. Consider a computable measure μ on 2^ω . Let $A_i = \{x \in 2^\omega \mid x(i) = 1\}$ where $x(i)$ is the i th bit of x . Then $\mathcal{A} = (A_i)$ is a generator of $(2^\omega, \mu)$. Further $[\sigma]_{\mathcal{A}} = [\sigma]^\prec$, $x \upharpoonright_{\mathcal{A}} n = x \upharpoonright n$, and $\text{name}_{\mathcal{A}}(x) = x$. Call \mathcal{A} the NATURAL GENERATOR of $(2^\omega, \mu)$, and $\{[\sigma]^\prec\}_{\sigma \in 2^{<\omega}}$ the NATURAL CELL DECOMPOSITION.

In this next proposition, recall that a set $S \subseteq 2^{<\omega}$ is PREFIX-FREE if there is no pair $\tau, \sigma \in S$ such that $\tau \prec \sigma$.

Proposition 4.7. *Let (\mathcal{X}, μ) be a computable probability space with generator \mathcal{A} and $\{[\sigma]_{\mathcal{A}}\}_{\sigma \in 2^{<\omega}}$ the corresponding cell decomposition. Then for each Σ_1^0 set $U \subseteq \mathcal{X}$ there is a c.e. set $\{\sigma_i\}$ (computable from U) such that $U = \bigcup_i [\sigma_i]_{\mathcal{A}}$ a.e. Further, $\{\sigma_i\}$ can be assumed to be prefix-free and such that $\mu([\sigma_i]_{\mathcal{A}}) > 0$ for all i .*

Proof. Straight-forward from Definition 4.1. □

It is clear that a generator \mathcal{A} is computable from its cell decomposition $\{[\sigma]_{\mathcal{A}}\}_{\sigma \in 2^{<\omega}}$, namely let

$$A_i = \bigcup_{\{\sigma : \sigma(i)=1\}} [\sigma]_{\mathcal{A}}.$$

Hence we will often confuse a generator and its cell decomposition writing both as \mathcal{A} . Further, this next proposition gives the criterion for when an indexed family $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$ forms an a.e. decidable cell decomposition.

Proposition 4.8. *Let (\mathcal{X}, μ) be a computable probability space. Let $\mathcal{A} = \{A_\sigma\}_{\sigma \in 2^{<\omega}}$ be a computably indexed family of Σ_1^0 sets such that*

- (1) *for all $\sigma \in 2^\omega$, $A_{\sigma 0} \cap A_{\sigma 1} = \emptyset$ and $A_{\sigma 0} \cup A_{\sigma 1} = A_\sigma$ a.e.*
- (2) *$\mu(A_\varepsilon) = 1$, and*
- (3) *for each Σ_1^0 set $U \subseteq \mathcal{X}$ there is a c.e. set $\{\sigma_i\}$ (computable from U) such that $U = \bigcup_i [\sigma_i]_{\mathcal{A}}$ a.e.*

Then \mathcal{A} is an a.e. decidable cell decomposition where $[\sigma]_{\mathcal{A}} = A_\sigma$ a.e. for all $\sigma \in 2^{<\omega}$.

Proof. Straight-forward from Definition 4.1 and Definition 4.4. □

Each computable probability space (\mathcal{X}, μ) is uniquely represented by a cell decomposition \mathcal{A} and the values $\mu([\sigma]_{\mathcal{A}})$.

The main difference between the method here and that of Gács [17] and Hoyrup and Rojas [22] is that they pick a canonical cell decomposition for each (\mathcal{X}, μ) . Also they assume every point $x \in \mathcal{X}$ is in some cell, and that no two points have the same \mathcal{A} -name. I, instead, work with all cell decompositions simultaneously and do not require as strong of properties. This will allow me to give a correspondence between cell decompositions and isomorphisms in Section 7.

5. COMPUTABLE RANDOMNESS ON COMPUTABLE PROBABILITY SPACES

In this section I define computable randomness on a computable probability space. As a first step, I have already done this for spaces $(2^\omega, \mu)$. The second step will be to define computable randomness with respect to a particular cell decomposition of the space. Finally, the last step is Theorem 5.7, where I will show the definition is invariant under the choice of cell decomposition.

There are two characterizations of computable randomness on $(2^\omega, \lambda)$ that use Martin-Löf tests. The first was due to Downey, Griffiths, and LaForte [13]. However, I will use another due to Merkle, Mihailović, and Slaman [28].

Definition 5.1 (Merkle et al. [28]). On $(2^\omega, \lambda)$ a Martin-Löf test (U_n) is called a **BOUNDED MARTIN-LÖF TEST** if there is a computable measure $\nu: 2^{<\omega} \rightarrow [0, \infty)$ such that for all $n \in \mathbb{N}$ and $\sigma \in 2^{<\omega}$

$$\mu(U_n \cap [\sigma]^\prec) \leq 2^{-n} \nu(\sigma).$$

We say that the test (U_n) is **BOUNDED BY** the measure ν .

Theorem 5.2 (Merkle et al. [28]). On $(2^\omega, \lambda)$, a string $x \in 2^\omega$ is computably random if and only if x is not covered by any bounded Martin-Löf test.

The next theorem and definition give five equivalent tests for computable randomness (with respect to a cell decomposition \mathcal{A}). (I also give a machine characterization of computable randomness in Section 6.) The integral test and Vitali cover test are new for computable randomness, although they are implicit in the proof of Theorem 5.2.

Theorem 5.3. Let \mathcal{A} be a cell decomposition of the computable probability space (\mathcal{X}, μ) . If $x \in \mathcal{X}$ is neither an unrepresented point nor in a null cell, then the following are equivalent.

- (1) (Martingale test) There is a computable martingale $M: 2^{<\omega} \rightarrow [0, \infty)$ satisfying

$$M(\sigma 0) \mu([\sigma 0]_{\mathcal{A}}) + M(\sigma 1) \mu([\sigma 1]_{\mathcal{A}}) = M(\sigma) \mu([\sigma]_{\mathcal{A}})$$

$$M(\sigma) \text{ is defined} \iff \mu([\sigma]_{\mathcal{A}}) > 0$$

for all $\sigma \in 2^{<\omega}$ such that $\limsup_{n \rightarrow \infty} M(x \upharpoonright_{\mathcal{A}} n) = \infty$.

- (2) (Martingale test with savings property, see for example [14, Proposition 2.3.8]) There is a computable martingale $N: 2^{<\omega} \rightarrow [0, \infty)$ satisfying the conditions of (1) and a partial-computable “savings function” $f: 2^{<\omega} \rightarrow [0, \infty)$ satisfying

$$f(\sigma) \leq N(\sigma) \leq f(\sigma) + 1$$

$$\sigma \preceq \tau \implies f(\sigma) \leq f(\tau)$$

$$f(\sigma) \text{ is defined} \iff \mu([\sigma]_{\mathcal{A}}) > 0$$

for all $\sigma, \tau \in 2^{<\omega}$ such that $\lim_{n \rightarrow \infty} N(x \upharpoonright_{\mathcal{A}} n) = \infty$.

- (3) (Integral test) There is a computable measure $\nu: 2^{<\omega} \rightarrow [0, \infty)$ and a lower semicomputable function $g: \mathcal{X} \rightarrow [0, \infty]$ satisfying

$$\int_{[\sigma]_{\mathcal{A}}} g d\mu \leq \nu(\sigma)$$

for all $\sigma \in 2^{<\omega}$ such that $g(x) = \infty$.

- (4) (Bounded Martin-Löf test) There is a computable measure $\nu: 2^{<\omega} \rightarrow [0, \infty)$ and a Martin-Löf test (U_n) satisfying

$$\mu(U_n \cap [\sigma]_{\mathcal{A}}) \leq 2^{-n} \nu(\sigma).$$

for all $n \in \mathbb{N}$ and $\sigma \in 2^{<\omega}$ such that (U_n) covers x .

(5) (*Vitali cover test*) There is a computable measure $\nu: 2^{<\omega} \rightarrow [0, \infty)$ and a Vitali cover (V_n) satisfying

$$\sum_n \mu(V_n \cap [\sigma]_{\mathcal{A}}) \leq \nu(\sigma)$$

for all $n \in \mathbb{N}$ and $\sigma \in 2^{<\omega}$ such that (V_n) Vitali covers x .

For (3) through (5), the measure ν may be assumed to be a probability measure and satisfy the following absolute-continuity condition,

$$(5.1) \quad \nu(\sigma) \leq \int_{[\sigma]_{\mathcal{A}}} h d\mu$$

for some integrable function h .

Further, each test is uniformly computable from any other.

Definition 5.4. Let \mathcal{A} be a cell decomposition of the space (X, μ) . Say $x \in X$ is **COMPUTABLY RANDOM** (with respect to \mathcal{A}) if x is neither an unrepresented point nor in a null cell, and x does not satisfy any of the equivalent conditions (1–5) of Theorem 5.3.

Before proving the theorem, here is a technical lemma. It will be needed for the savings property in (2).

Lemma 5.5 (Technical lemma). *Let (a_n) be a sequence of positive real numbers. Define (b_n) and (c_n) recursively as follows: $b_0 = a_0$, $c_0 = b_0 - 1$,*

$$b_{n+1} = c_n + \frac{a_{n+1}}{a_n}(b_n - c_n)$$

and $c_{n+1} = \max(c_n, b_{n+1} - 1)$. If $\limsup_n a_n = \infty$, then $\lim_n b_n = \infty$.

Proof. Let (n_i) be indices such that $c_{n_i} = b_{n_i} - 1$ listed in order. By induction on $n \in [n_i, n_{i+1} - 1]$ we have $c_n = b_{n_i} - 1$ and

$$b_{n+1} = b_{n_i} + \frac{a_{n+1}}{a_{n_i}} - 1.$$

Since $\limsup_n a_n = \infty$, there exists some $m > n_i$ such $b_m - 1 \geq b_{n_i} - 1 = c_{n_i}$. The first such m is n_{i+1} . This is also the first m such that $a_m \geq a_{n_i}$. Therefore (n_i) is a infinite series, $a_{n_{i+1}} \geq a_{n_i}$, $\lim_i a_{n_i} = \infty$, and

$$b_{n_{i+1}} = b_{n_i} + \frac{a_{n_{i+1}}}{a_{n_i}} - 1.$$

We have that $c_{n_i} \geq \log(a_{n_i})$ (natural logarithm) by the identity $x - 1 \geq \log(x)$ and by induction:

$$\begin{aligned} c_{n_{i+1}} &= b_{n_{i+1}} - 1 = (b_{n_i} - 1) + \left(\frac{a_{n_{i+1}}}{a_{n_i}} - 1 \right) \\ &\geq \log(a_{n_i}) + \log\left(\frac{a_{n_{i+1}}}{a_{n_i}} \right) = \log(a_{n_{i+1}}) + 1. \end{aligned}$$

Hence $\lim_i c_{n_i} \geq \lim_i \log(a_{n_i}) = \infty$, and since c_n is nondecreasing, $\lim_n b_n \geq \lim_n c_n = \infty$. \square

Proof of Theorem 5.3. (1) implies (2): The idea is to bet with the martingale M as usual, except at each stage set some of the winnings aside into a savings account $f(\sigma)$ and bet only with the remaining capital. Formally, define N and f recursively as follows. (One may assume $M(\sigma) > 0$ for all σ by adding 1 to $M(\sigma)$.) Start with $N(\varepsilon) = M(\varepsilon)$ and $f(\varepsilon) = N(\varepsilon) - 1$. At σ , for $i = 0, 1$ let

$$N(\sigma i) = f(\sigma) + \frac{M(\sigma i)}{M(\sigma)}(N(\sigma) - f(\sigma))$$

and $f(\sigma i) = \max(f(\sigma), N(\sigma i) - 1)$. By the technical lemma above, $\lim_n N(x \upharpoonright_{\mathcal{A}} n) = \infty$.

(2) implies (3): Let $\nu(\sigma) = N(\sigma)\mu([\sigma]_{\mathcal{A}})$ and $g(x) = \sup_{s \rightarrow \infty} f(x \upharpoonright_{\mathcal{A}} s)$. Then $\int_{[\sigma]_{\mathcal{A}}} g d\mu \leq \nu(\sigma) \leq \int_{[\sigma]_{\mathcal{A}}} (g + 1) d\mu$, which also shows ν satisfies the absolute-continuity condition of formula (5.1). If $N(\varepsilon)$ is scaled to be 1, then ν is a probability measure.

(3) implies (1): Let $M(\sigma) = \nu(\sigma)/\mu([\sigma]_{\mathcal{A}})$. Then $M(x \upharpoonright_{\mathcal{A}} k) \geq \frac{\int_{[x \upharpoonright_{\mathcal{A}} k]_{\mathcal{A}}} g d\mu}{\mu([x \upharpoonright_{\mathcal{A}} k]_{\mathcal{A}})}$, which converges to ∞ .

(3) implies (4): Let $U_n = \{x \mid g(x) > 2^n\}$. By Markov's inequality, $\mu(U_n \cap [\sigma]_{\mathcal{A}}) \leq \int_{[\sigma]_{\mathcal{A}}} g d\mu \leq \nu(\sigma)$.

(4) implies (5): Let $V_n = U_n$.

(5) implies (3): Let $g = \sum_n \mathbf{1}_{V_n}$. \square

In this next proposition, I show the standard randomness implications (as in formula (3.1)) still hold.

Proposition 5.6. *Let (\mathcal{X}, μ) be a computable probability space. If $x \in \mathcal{X}$ is Martin-Löf random, then x is computably random (with respect to every cell decomposition \mathcal{A}). If $x \in \mathcal{X}$ is computably random (with respect to a cell decomposition \mathcal{A}), then x is Schnorr random, and hence Kurtz random.*

Proof. The statement on Martin-Löf randomness follows from the bounded Martin-Löf test (Theorem 5.3 (4)).

For the last statement, assume x is not Schnorr random. If x is an unrepresented point or in a null cell, then x is not computably random by Definition 5.4. Else, there is some Vitali cover (V_n) where $\sum_n \mu(V_n)$ is computable and (V_n) Vitali-covers x . Define $\nu: 2^{<\omega} \rightarrow [0, \infty)$ as $\nu(\sigma) = \sum_n \mu(V_n \cap [\sigma]_{\mathcal{A}})$. Then clearly, $\mu(V_n \cap [\sigma]_{\mathcal{A}}) \leq \nu(\sigma)$ for all n and σ . By the Vitali cover test (Theorem 5.3 (5)), it is enough to show that ν is a computable measure. It is straightforward to verify that $\nu(\sigma 0) + \nu(\sigma 1) = \nu(\sigma)$. As for the computability of ν ; notice $\nu(\sigma)$ is lower semicomputable for each σ since μ is a computable probability measure (see Definition 3.2). Then since $\nu(\varepsilon) = \sum_n \mu(V_n)$ is computable, ν is a computable measure. \square

Theorem 5.7. *The definition for computable randomness does not depend on the choice of cell decomposition.*

Proof. Before giving the details, here is the main idea. It suffices to convert a test with respect to one cell decomposition \mathcal{A} to another test which covers the same points, but is with respect to a different cell decomposition \mathcal{B} . In order to do this, take a bounding measure ν with respect to \mathcal{A} (which is really a measure on 2^ω) and transfer it to an actual measure π on \mathcal{X} . Then transfer π back to a bounding measure κ with respect to \mathcal{B} . In order to guarantee that this will work, we will assume ν satisfies the absolute-continuity condition of formula 5.1, which ensures that π exists and is absolutely continuous with respect to μ .

Now I give the details. Assume $x \in X$ is not computably random with respect to the cell decomposition \mathcal{A} of the space. Let \mathcal{B} be another cell decomposition. If x is an unrepresented point or in a null cell, then x is not a Kurtz random point of (\mathcal{X}, μ) , and by Proposition 5.6, x is not computably random with respect to \mathcal{B} .

So assume x is neither an unrepresented point nor in a null cell. By condition (4) of Theorem 5.3 there is some Martin-Löf test (U_n) bounded by a probability measure ν such that (U_n) covers x . Further, ν can be assumed to satisfy the absolute-continuity condition in formula (5.1).

Claim. There is a computable probability measure π on \mathcal{X} defined by $\pi([\sigma]_{\mathcal{A}}) = \nu(\sigma)$ which is absolutely continuous with respect to μ , i.e. every μ -null set is a π -null set.

Proof of claim. This is basically the Carathéodory extension theorem. The collection $\{[\sigma]_{\mathcal{A}}\}_{\sigma \in 2^{<\omega}}$ is essentially a semi-ring. A semi-ring contains \emptyset , is closed under intersections, and for each A, B in the semi-ring, there are pairwise disjoint sets C_1, \dots, C_n in the semi-ring such that $A \setminus B = C_1 \cup \dots \cup C_n$. To make this collection a semi-ring which generates the Borel sigma-algebra, add every μ -null set and every set which is μ -a.e. equal to $[\sigma]_{\mathcal{A}}$ for some σ .

Define $\pi([\sigma]_{\mathcal{A}}) = \nu(\sigma)$ and $\pi(\emptyset) = 0$ and similarly for the μ -a.e. equivalent sets. (This is well defined since if $[\sigma]_{\mathcal{A}} = [\tau]_{\mathcal{A}}$ μ -a.e. then by the absolute continuity condition, $\nu(\sigma) = \nu(\tau)$, and similarly if $\mu([\sigma]_{\mathcal{A}}) = 0$, then $\nu(\sigma) = 0$.) Now, it is enough to show π is a pre-measure, specifically that it satisfies countable additivity. Assume for some pairwise disjoint family $\{A_i\}$ and some B , both in the semi-ring, that $B = \bigcup_i A_i$. If B is μ -null, then each A_i is as well. By the definition of π on μ -null sets, we have $\pi(B) = 0 = \sum_i \pi(A_i)$. If B is not μ -null, then $B = [\tau]_{\mathcal{A}}$ μ -a.e. for some τ and each A_i of positive μ -measure is μ -a.e. equal to $[\sigma_i]_{\mathcal{A}}$ for some $\sigma_i \succeq \tau$. For each k , let $C_k = [\tau]_{\mathcal{A}} \setminus \bigcup_{i=0}^{k-1} [\sigma_i]_{\mathcal{A}}$, which is a finite union of basic open sets in 2^ω . Let D_k be the same union as C_k but replacing each $[\sigma_i]_{\mathcal{A}}$ with $[\sigma]_{\mathcal{A}}$. Then by the absolute continuity condition,

$$\pi(B) - \sum_{i=0}^{k-1} \pi(A_i) = \nu(\tau) - \sum_{i=0}^{k-1} \nu(\sigma_i) = \nu(C_k) = \int_{D_k} h d\mu$$

Since $[\tau]_{\mathcal{A}} = \bigcup_i [\sigma_i]_{\mathcal{A}}$ μ -a.e., the right-hand-side goes to zero as $k \rightarrow \infty$. So π is a pre-measure and may be extended to a measure by the Carathéodory extension theorem.

Similarly by approximation, π satisfies $\pi(A) \leq \int_A h d\mu$ for all Borel sets A and hence is absolutely continuous with respect to μ .

To see π is a computable probability measure on \mathcal{X} , take a Σ_1^0 set U . By Proposition 4.7, there is a c.e., prefix-free set $\{\sigma_i\}$ of finite strings such that $U = \bigcup_i [\sigma_i]_{\mathcal{A}}$ μ -a.e. (and so π -a.e. by absolute continuity). As

this union is disjoint, $\pi(U) = \sum_i \pi([\sigma_i]_{\mathcal{A}}) = \sum_i \nu(\sigma_i)$ μ -a.e. and so $\pi(U)$ is lower-semicomputable. Since $\pi(\mathcal{X}) = 1$, π is a computable probability measure. This proves the claim. \square

Let π be as in the claim. Since π is absolutely continuous with respect to μ , any a.e. decidable set of μ is an a.e. decidable set of π . In particular, the measures $\pi([\tau]_{\mathcal{B}})$ are computable from τ . Now transfer π back to a measure $\kappa: 2^{<\omega} \rightarrow [0, \infty)$ using $\kappa(\sigma) = \pi([\sigma]_{\mathcal{B}})$. This is a computable probability measure since $\pi([\sigma]_{\mathcal{B}})$ is computable.

Last, we show the Martin-Löf test (U_n) is bounded by κ with respect to the cell decomposition \mathcal{B} . To see this, fix $\tau \in 2^{<\omega}$ and take the c.e., prefix-free set $\{\sigma_i\}$ of finite strings such that $[\tau]_{\mathcal{B}} = \bigcup_i [\sigma_i]_{\mathcal{A}}$ μ -a.e. (and so π -a.e.). Then $\kappa(\tau) = \sum_i \nu(\sigma_i)$, and for each n ,

$$\mu(U_n \cap [\tau]_{\mathcal{B}}) = \sum_i \mu(U_n \cap [\sigma_i]_{\mathcal{A}}) \leq \sum_i 2^{-n} \nu(\sigma_i) = 2^{-n} \kappa(\tau). \quad \square$$

Theorem 5.3 is just a sample of the many equivalent definitions for computable randomness. I conjecture that the other known characterizations of computable randomness, see for example Downey and Hirschfeld [14, Section 7.1], can be extended to arbitrary computable Polish spaces using the techniques above. As well, other test characterizations for Martin-Löf randomness can be extended to computable randomness by “bounding the test” with a computable measure or martingale. (See Section 6 for an example using machines.) Further, the proof of Theorem 5.7 shows that the bounding measure ν can be assumed to be a measure on \mathcal{X} , instead of 2^ω , under the additional condition that \mathcal{A} is a cell decomposition for both (\mathcal{X}, μ) and (\mathcal{X}, ν) . Similarly, we could modify the martingale test to assume M is a martingale on (\mathcal{X}, μ) (in the sense of probability theory) with an appropriate filtration.

Actually, the above ideas can be used to show any L^1 -bounded a.e. computable martingale (in the sense of probability theory) converges on computable randoms if the filtration converges to the Borel sigma-algebra (or even a “computable” sigma-algebra) and the L^1 -bound is computable. This can be extended to (the Schnorr layerwise-computable representatives of) L^1 -computable martingales as well. The proof is beyond the scope of this paper and will be published separately.

In Section 11, I give ideas on how computable randomness can be defined on an even broader class of spaces, and also on non-computable probability spaces. I end this section by showing that Definition 5.4 is consistent with the usual definitions of computable randomness on 2^ω , Σ^ω , and $[0, 1]$.

Example 5.8. Consider a computable probability measure μ on 2^ω . It is easy to see that computable randomness in the sense of Definition 5.4 with respect to the natural cell decomposition is equivalent to computable randomness on 2^ω as defined in Definition 2.3. Since Definition 5.4 is invariant under the choice of cell decomposition (Theorem 5.7), the two definitions agree on $(2^\omega, \mu)$.

Example 5.9. Consider a computable probability measure μ on Σ^ω where $\Sigma = \{a_0, \dots, a_{k-1}\}$ is a finite alphabet. It is natural to define a martingale $M: \Sigma^\omega \rightarrow [0, \infty)$ as one satisfying the fairness condition

$$M(\sigma a_0)\mu(\sigma a_0) + \dots + M(\sigma a_{k-1})\mu(\sigma a_{k-1}) = M(\sigma)\mu(\sigma)$$

for all $\sigma \in \Sigma^{<\omega}$ (along with the impossibility condition from Definition 2.2). A little thought reveals that by systematically grouping and upgrouping cylinder sets M can be turned into a binary martingale which succeeds on the same points. For example, given a martingale M on 3^ω , one may first split $[\sigma]^\prec$ into $[\sigma 0]^\prec$ and $A_\sigma = [\sigma 1]^\prec \cup [\sigma 2]^\prec$. Define,

$$M(A_\sigma) = \frac{M(\sigma 1)\mu(\sigma 1) + M(\sigma 2)\mu(\sigma 2)}{\mu([\sigma 1]^\prec \cup [\sigma 2]^\prec)}$$

and notice the fairness condition is still satisfied,

$$M(\sigma 0)\mu(\sigma 0) + M(A_\sigma)\mu(A_\sigma) = M(\sigma)\mu(\sigma).$$

In the next step, one may split A_σ into $[\sigma 1]^\prec$ and $[\sigma 2]^\prec$ to give

$$M(\sigma 1)\mu(\sigma 1) + M(\sigma 2)\mu(\sigma 2) = M(A_\sigma)\mu(A_\sigma).$$

This grouping and ungrouping of cylinder sets forms a (binary) cell decomposition \mathcal{A} on $(3^\omega, \mu)$. If M was first given the savings property, this new martingale succeeds on the same points. It follows that $x \in 3^\omega$ is computably random in the natural sense if and only if it is computably random as in Definition 5.4.

Example 5.10. Let $([0, 1], \lambda)$ be the space $[0, 1]$ with the Lebesgue measure. Let $A_i = \{x \in [0, 1] \mid \text{the } i\text{th binary digit of } x \text{ is } 1\}$. Then $\mathcal{A} = (A_i)$ is a generator of $([0, 1], \lambda)$ and $[\sigma]_{\mathcal{A}} = [0.\sigma, 0.\sigma + 2^{-|\sigma|})$ a.e. A little thought reveals that $x \in ([0, 1], \lambda)$ is computably random (in the sense of Definition 5.4) if and only if the binary expansion of x is computably random in $(2^\omega, \lambda)$ with the fair-coin measure. This is the standard definition of computable randomness on $([0, 1], \lambda)$. Further, using a base b other than binary gives a different generator, for example let $A_{bi+j} = \{x \in [0, 1] \mid \text{the } i\text{th } b\text{-ary digit of } x \text{ is } j\}$ where $0 \leq j < b$. Yet, the computably random points remain the same. Hence computable randomness on $([0, 1], \lambda)$ is base invariant [12, 37]. (The proof of Theorem 5.7 has similarities to the proof of Brattka, Miller and Nies [12], but as mentioned in the introduction, there are also key differences.) Also see Example 7.11.

More examples are given at the end of Section 7.

6. MACHINE CHARACTERIZATIONS OF COMPUTABLE AND SCHNORR RANDOMNESS

In this section I give machine characterizations of computable and Schnorr randomness for computable probability spaces. This has already been done for Martin-Löf randomness.

Recall the following definition and fact.

Definition 6.1. A machine M is a partial-computable function $M: 2^{<\omega} \rightarrow 2^{<\omega}$. A machine is PREFIX-FREE if $\text{dom } M$ is prefix-free. The prefix-free Kolmogorov complexity of σ relative to a machine M is

$$K_M(\sigma) = \inf \{|\tau| \mid \tau \in 2^{<\omega} \text{ and } M(\tau) = \sigma\}.$$

(There is a non-prefix-free version of complexity as well.)

Theorem 6.2 (Schnorr (see [14, Theorem 6.2.3])). *A string $x \in (2^\omega, \lambda)$ is Martin-Löf random if and only if for all prefix-free machines M ,*

$$(6.1) \quad \limsup_{n \rightarrow \infty} (n - K_M(x \upharpoonright n)) < \infty.$$

Schnorr's theorem has been extended to both Schnorr and computable randomness.

Definition 6.3. For a machine M define the semimeasure $\text{meas}_M: 2^{<\omega} \rightarrow [0, \infty)$ as

$$\text{meas}_M(\sigma) = \sum_{\substack{\tau \in \text{dom } M \\ M(\tau) \succeq \sigma}} 2^{-|\tau|}.$$

A machine M is a COMPUTABLE-MEASURE MACHINE if $\text{meas}_M(\varepsilon)$ is computable. A machine M is a BOUNDED MACHINE if there is some computable-measure ν such that $\text{meas}_M(\sigma) \leq \nu(\sigma)$ for all $\sigma \in 2^{<\omega}$.

Downey, Griffiths, and LaForte [13] showed that $x \in (2^\omega, \lambda)$ is Schnorr random precisely if formula (6.1) holds for all prefix-free, computable-measure machines. Mihailović (see [14, Theorem 7.1.25]) showed that $x \in (2^\omega, \lambda)$ is computably random precisely if formula (6.1) holds for all prefix-free, bounded machines.

Schnorr's theorem was extended to all computable probability measures on Cantor space by Gács [16]. Namely, replace formula (6.1) with

$$\limsup_{n \rightarrow \infty} (-\log_2 \mu([x \upharpoonright n]^\prec) - K_M(x \upharpoonright n)) < \infty.$$

If $\mu([x \upharpoonright n]) = 0$ for any n then we say this inequality is false. Hoyrup and Rojas [22] extended this to any computable probability space. Here, I do the same for Schnorr and computable randomness (I include Martin-Löf randomness for completeness).

Theorem 6.4. *Let (\mathcal{X}, μ) be a computable probability space and $x \in \mathcal{X}$.*

(1) *$x \in \mathcal{X}$ is Martin-Löf random precisely if*

$$(6.2) \quad \limsup_{n \rightarrow \infty} (-\log_2 \mu([x \upharpoonright_{\mathcal{A}} n]_{\mathcal{A}}) - K_M(x \upharpoonright_{\mathcal{A}} n)) < \infty.$$

holds for all prefix-free machines M . (Again, we say formula (6.2) is false if $\mu([x \upharpoonright n]) = 0$ for any n .)

(2) *$x \in \mathcal{X}$ is computably random precisely if formula (6.2) holds for all prefix-free, computable-measure machines M .*

(3) *$x \in \mathcal{X}$ is Schnorr random precisely if formula (6.2) holds for all prefix-free, bounded machines M .*

Further, (1) through (3) hold even if M is not assumed to be prefix-free, but only that $\text{meas}_M(\varepsilon) \leq 1$.

Proof. Slightly modify the proofs of Theorems 6.2.3, 7.1.25, and 7.1.15 in Downey and Hirschfeld [14], respectively. \square

7. COMPUTABLE RANDOMNESS AND ISOMORPHISMS

In this section I give another piece of evidence that the definition of computable randomness in this paper is robust, namely that the computably random points are preserved under isomorphisms between computable probability spaces. I also show a one-to-one correspondence between cell decompositions of a computable measure space and isomorphisms from that space to the Cantor space.

Definition 7.1. Let (\mathcal{X}, μ) and (\mathcal{Y}, ν) be computable probability spaces.

- (1) A partial map $T: \mathcal{X} \rightarrow \mathcal{Y}$ is said to be **PARTIAL COMPUTABLE** if there is a partial-computable function $F: \mathbb{N}^\omega \rightarrow \mathbb{N}^\omega$ which given a Cauchy-name for $x \in \mathcal{X}$ returns the Cauchy-name for $T(x)$, and further, the domain of T is maximal for this h , i.e. $x \in \text{dom}(T)$ if and only if for all $a, b \in \mathbb{N}^\omega$ which are Cauchy-names for x , then $a, b \in \text{dom}(F)$ and both $F(a)$ and $F(b)$ are Cauchy-names for the same point in \mathcal{Y} .
- (2) A partial map $T: (\mathcal{X}, \mu) \rightarrow \mathcal{Y}$ is said to be **A.E. COMPUTABLE** if it is partial computable with a measure-one domain.
- (3) (Hoyrup and Rojas [22]) A partial map $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is said to be an **(A.E. COMPUTABLE) MORPHISM** if it is a.e. computable and measure preserving, i.e. $\mu(T^{-1}(A)) = \nu(A)$ for all measurable $A \subseteq \mathcal{Y}$.
- (4) (Hoyrup and Rojas [22]) A pair of partial maps $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ and $S: (\mathcal{Y}, \nu) \rightarrow (\mathcal{X}, \mu)$ are said to be an **(A.E. COMPUTABLE) ISOMORPHISM** if both maps are (a.e. computable) morphisms such that $(S \circ T)(x) = x$ for μ -a.e. $x \in \mathcal{X}$ and $(T \circ S)(y) = y$ for ν -a.e. $y \in \mathcal{Y}$. We also say $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is an isomorphism if such an S exists.

Note that this definition of isomorphism differs slightly from that of Hoyrup and Rojas [22]. They require that the domain must also be dense.

The definition of partial-computable map above basically says that the domain of T is determined by its algorithm and not some artificial restriction on the domain. This next proposition shows that this is equivalent to saying that the domain is Π_2^0 .

Proposition 7.2. *A partial map $T: \mathcal{X} \rightarrow \mathcal{Y}$ is partial computable if and only if the domain of T is a Π_2^0 set and T is computable on its domain.*

Proof. The proof of the first direction is straightforward. (For example, given $F: \mathbb{N}^\omega \rightarrow \mathbb{N}^\omega$, then $\text{dom}(F)$ is Π_2^0 in \mathbb{N}^ω [39, Theorem 2.2.4]. Also, the set of \mathcal{X} -Cauchy-names is Π_1^0 and the set of pairs (a, b) such that $a \approx_{\mathcal{X}} b$ (i.e. a and b are Cauchy-names for the same point in \mathcal{X}) and $h(a) \not\approx_{\mathcal{Y}} h(b)$ is Δ_2^0 .)

For the other direction, let D be the Π_2^0 domain. Then $D = \bigcap_n U_n$ where (U_n) is a computable sequence of Σ_1^0 sets. Let $F: \mathbb{N}^\omega \rightarrow \mathbb{N}^\omega$ be the partial-computable map from Cauchy-names to Cauchy-names that computes T . Modify $F(a)$ to return an n th approximation only if a “looks like” a Cauchy-name for some $x \in U_n$. \square

This next corollary says a.e. computable maps are defined on Kurtz randoms. Further, Kurtz randomness can be characterized by a.e. computable maps, and a.e. computable maps are determined by their values on Kurtz randoms. (For a different characterization of Kurtz randomness using a.e. computable functionals, see Hertling and Yongge [19].)

Corollary 7.3. *Let (\mathcal{X}, μ) be a computable measure space and \mathcal{Y} a computable Polish space. For $x \in \mathcal{X}$, x is Kurtz random if and only if it is in the domain of every a.e. computable map $T: (\mathcal{X}, \mu) \rightarrow \mathcal{Y}$. Further, two a.e. computable maps are a.e. equal if and only if they agree on Kurtz randoms.*

Proof. For the first part, if x is Kurtz random, it avoids all null Σ_2^0 sets, and by Proposition 7.2 is in the domain of every a.e. computable map. Conversely, x is not Kurtz random, it is in some null Σ_2^0 set A . But the partial map $T: (\mathcal{X}, \mu) \rightarrow \mathcal{Y}$ with domain $\mathcal{X} \setminus A$ and $T(x) = 1$ for $x \in \mathcal{X} \setminus A$ is a.e. computable by Proposition 7.2.

For the second part, let $T, S: (\mathcal{X}, \mu) \rightarrow \mathcal{Y}$ be a.e. computable maps that are a.e. equal. The set $\{x \in \mathcal{X} \mid T(x) \neq S(x)\}$ is a null Σ_2^0 set in \mathcal{X} . Conversely, if $T(x) = S(x)$ for all Kurtz randoms x , then $T = S$ a.e. \square

Remark 7.4 (Preimages of Σ_1^0 sets). The preimage of a Σ_1^0 set under an computable map is still Σ_1^0 . Unfortunately, the preimage of a Σ_1^0 set under an *partial computable map* is not always Σ_1^0 . However, it is equal to the intersection of a Σ_1^0 set and the domain of the map. (We leave the details to the reader.) As an abuse of notation, if $T: \mathcal{X} \rightarrow \mathcal{Y}$ is a partial-computable map and $V \subseteq \mathcal{Y}$ is Σ_1^0 , we will define $T^{-1}(V)$ to be a Σ_1^0 set $U \subseteq \mathcal{X}$ such that for all $x \in \mathcal{X}$, $x \in U \cap \text{dom}(T) \Leftrightarrow T(x) \in V$. (We leave it to the reader to verify that such a U can be computed uniformly from the codes for T and V .) Also, if $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is a morphism, it is easy to see that $\mu(U) = \nu(V)$. We can similarly define the preimage of a $\Pi_1^0, \Sigma_2^0, \Pi_2^0$ set to remain in the same point class. Last, if $B \subseteq \mathcal{Y}$ is a.e. decidable with a.e. decidable pair (V_0, V_1) , then define $T^{-1}(B)$ to be the a.e. decidable set A given by the pair $(T^{-1}(V_0), T^{-1}(V_1))$.

This next proposition shows that for many common notions of randomness are preserved by morphisms, and the set of randoms is preserved under isomorphisms.

Proposition 7.5. *If $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is a morphism and $x \in \mathcal{X}$ is Martin-Löf random, then $T(x)$ is Martin-Löf random. The same is true of Kurtz and Schnorr randomness. Hence, if T is an isomorphism, then x is Martin-Löf (respectively Kurtz, Schnorr) random if and only if $T(x)$ is.*

Proof. Assume $T(x)$ is not Martin-Löf random in (\mathcal{Y}, ν) . Then there is a Martin-Löf test (U_n) in (\mathcal{Y}, ν) which covers $T(x)$. Let $V_n = T^{-1}(U_n)$ for each n . By Remark 7.4 (V_n) is a Martin-Löf test in (\mathcal{X}, μ) which covers x . Hence x is not Martin-Löf random in (\mathcal{X}, μ) .

Kurtz and Schnorr randomness follow similarly, namely the inverse image of a test is still a test. \square

(Bienvenu and Porter have pointed out to me the following partial converse to Proposition 7.5, which was first proved by Shen—see [7]. If $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is a morphism and y is Martin-Löf random for (\mathcal{Y}, ν) , then there is some x that is Martin-Löf random for (\mathcal{X}, μ) such that $T(x) = y$.)

In Corollary 9.7, we will see that computable randomness is not preserved by morphisms. However, just looking at the previous proof gives a clue. There is another criterion to the tests for computable randomness besides complexity and measure, namely the cell decompositions of the space. The “inverse image” of cell decomposition may not be a cell decomposition.

However, if T is an isomorphism the situation is much better. Indeed, these next three propositions show a correspondence between isomorphisms and cell decompositions. We say two cell decompositions \mathcal{A} and \mathcal{B} of a computable probability space (\mathcal{X}, μ) are almost-everywhere equal if $[\sigma]_{\mathcal{A}} = [\sigma]_{\mathcal{B}}$ a.e. for all $\sigma \in 2^{<\omega}$. Recall, two isomorphisms are almost-everywhere equal if they are pointwise a.e. equal.

Proposition 7.6 (Isomorphisms to cell decompositions). *If $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ is an isomorphism and \mathcal{B} is a cell decomposition of (\mathcal{Y}, ν) , then there is an a.e. unique cell decomposition \mathcal{A} (which we will notate as $T^{-1}(\mathcal{B})$) such that $\text{name}_{\mathcal{A}}(x) = \text{name}_{\mathcal{B}}(T(x))$ for μ -a.e. x . This cell decomposition \mathcal{A} is given by $[\sigma]_{\mathcal{A}} = T^{-1}([\sigma]_{\mathcal{B}})$. In particular, every isomorphism $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \nu)$ induces a cell decomposition \mathcal{A} such that $\text{name}_{\mathcal{A}}(x) = T(x)$ for μ -a.e. x .*

Proof. We will show $[\sigma]_{\mathcal{A}} = T^{-1}([\sigma]_{\mathcal{B}})$ defines a cell decomposition \mathcal{A} . By Remark 7.4, $T^{-1}([\sigma]_{\mathcal{B}})$ is Σ_1^0 uniformly from σ . Clearly, $\mu([\varepsilon]_{\mathcal{A}}) = 1$, $[\sigma 0]_{\mathcal{A}} \cap [\sigma 1]_{\mathcal{A}} = \emptyset$, and $[\sigma 0]_{\mathcal{A}} \cup [\sigma 1]_{\mathcal{A}} = [\sigma]_{\mathcal{A}}$ μ -a.e. Finally, take a Σ_1^0 set $U \subseteq \mathcal{X}$. By Proposition 4.8, it is enough to show there is some c.e. set $\{\sigma_i\}$ such that $U = \bigcup_i [\sigma_i]_{\mathcal{A}}$ μ -a.e. Let S be the inverse isomorphism to T . Then define $V = S^{-1}(U)$. By Remark 7.4, V is Σ_1^0 in \mathcal{Y} and $T^{-1}(V) = U$ μ -a.e. By Proposition 4.7 there is some c.e. set $\{\sigma_i\}$ such that $V = \bigcup_i [\sigma_i]_{\mathcal{B}}$ ν -a.e. and therefore $U = T^{-1}(V) = \bigcup_i T^{-1}([\sigma_i]_{\mathcal{B}}) = \bigcup_i [\sigma_i]_{\mathcal{A}}$ μ -a.e. Therefore, $[\sigma]_{\mathcal{A}} = T^{-1}([\sigma]_{\mathcal{B}})$ defines a cell decomposition \mathcal{A} .

For μ -a.e. x , $x \in \text{dom}(T) \cap \text{dom}(\text{name}_{\mathcal{A}})$. Then for all n , $x \in [x \upharpoonright_{\mathcal{A}} n]_{\mathcal{A}} = T^{-1}([x \upharpoonright_{\mathcal{B}} n]_{\mathcal{B}})$. By Remark 7.4, $T(x) \in [x \upharpoonright_{\mathcal{B}} n]_{\mathcal{B}}$. Therefore $\text{name}_{\mathcal{B}}(T(x)) = \text{name}_{\mathcal{A}}(x)$.

For $\mathcal{Y} = 2^\omega$, let \mathcal{B} be the natural cell decomposition of $(2^\omega, \nu)$, then $[\sigma]_{\mathcal{B}} = [\sigma]^{<}$ for all $\sigma \in 2^{<\omega}$. Therefore for μ -a.e. x , $\text{name}_{\mathcal{A}}(x) = \text{name}_{\mathcal{B}}(T(x)) = T(x)$.

To show the cell decomposition \mathcal{A} is unique, assume \mathcal{A}' is another cell decomposition such that for μ -a.e. x , the \mathcal{A} -name and \mathcal{A}' -name of x are both the \mathcal{B} name of $T(x)$. Then $[\sigma]_{\mathcal{A}} = [\sigma]_{\mathcal{A}'}$ μ -a.e. for all $\sigma \in 2^{<\omega}$. \square

Proposition 7.7 (Cell decompositions to isomorphisms). *Let (\mathcal{X}, μ) be a computable probability space with cell decomposition \mathcal{A} . There is a unique computable probability space $(2^\omega, \mu_{\mathcal{A}})$ such that $\text{name}_{\mathcal{A}}: (\mathcal{X}, \mu) \rightarrow (2^\omega, \mu_{\mathcal{A}})$ is an isomorphism. Namely, $\mu_{\mathcal{A}}(\sigma) = \mu([\sigma]_{\mathcal{A}})$.*

Proof. If such a measure $\mu_{\mathcal{A}}$ exists, it must be unique. Indeed, since $\text{name}_{\mathcal{A}}$ is then measure-preserving, $\mu_{\mathcal{A}}$ must satisfy $\mu_{\mathcal{A}}(\sigma) = \mu(\text{name}_{\mathcal{A}}^{-1}([\sigma]^{\prec})) = \mu([\sigma]_{\mathcal{A}})$, which uniquely defines $\mu_{\mathcal{A}}$.

It remains to show the map $\text{name}_{\mathcal{A}}: (\mathcal{X}, \mu) \rightarrow (2^{\omega}, \mu_{\mathcal{A}})$ which maps x to its \mathcal{A} -name is an isomorphism. Clearly, $\mu_{\mathcal{A}}$ is a computable measure since $\mu([\sigma]_{\mathcal{A}})$ is computable. The map $\text{name}_{\mathcal{A}}$ which takes x to its \mathcal{A} -name is measure preserving for cylinder sets and therefore for all sets by approximation. The map from x to $x \upharpoonright_{\mathcal{A}} n$ is a.e. computable. Indeed, wait for x to show up in one of the sets $[\sigma]_{\mathcal{A}}$ where $|\sigma| = n$. Hence the map from x to its \mathcal{A} -name is also a.e. computable. So $\text{name}_{\mathcal{A}}$ is a morphism. (As an extra verification, clearly $\text{dom}(\text{name}_{\mathcal{A}})$ is a Π_2^0 measure-one set.)

The inverse of $\text{name}_{\mathcal{A}}$ will be the map S from (a measure-one set of) \mathcal{A} -names $y \in 2^{\omega}$ to points $x \in \mathcal{X}$ such that $\text{name}_{\mathcal{A}}(x) = y$. The algorithm for S will be similar to the algorithm given by the proof of the Baire category theorem (see [10]). Pick $y \in 2^{\omega}$. We compute $S(y)$ by a back-and-forth argument. Assume $\tau \prec y$. Recall, $[\tau]_{\mathcal{A}}$ is Σ_1^0 . We can enumerate a sequence of pairs (a_i, k_i) where each a_i is a simple point of \mathcal{X} and each $k_i > |\tau|$ such that $[\tau]_{\mathcal{A}} = \bigcup_i B(a_i, 2^{-k_i})$. Further, by Proposition 4.7, we have that for each i , there is a c.e. set $\{\sigma_j^i\}$ such that $B(a_i, 2^{-k_i}) = \bigcup_j [\sigma_j^i]_{\mathcal{A}}$ μ -a.e. (We may assume $|\sigma_j^i| > |\tau|$ for all i, j .) Given y , compute the Cauchy-name of $S(y)$ as follows. Start with $\tau_1 = y \upharpoonright 1$. Then search for $\sigma_j^i \prec y$. If we find one, let $b_1 = a_i$ be the first approximation. Now continue with $\tau_2 = \sigma_j^i$, and so on. This gives a Cauchy-name (b_n) . The algorithm will fail if at some stage it cannot find any $\sigma_j^i \prec y$. But then $y \in [\tau]^{\prec} \setminus \bigcup_i \bigcup_j [\sigma_j^i]^{\prec}$, which by the definition of $\mu_{\mathcal{A}}$, is a $\mu_{\mathcal{A}}$ -measure-zero set since $[\tau]_{\mathcal{A}} = \bigcup_i \bigcup_j [\sigma_j^i]_{\mathcal{A}}$ μ -a.e. Hence S is a.e. computable.

By the back-and-forth algorithm, $\text{name}_{\mathcal{A}}(S(y)) = y$ for all $y \in \text{dom}(S)$. To show $S(\text{name}_{\mathcal{A}}(x)) = x$ a.e., assume $x \in \text{dom}(\text{name}_{\mathcal{A}})$. Consider the back-and-forth sequence created by the algorithm: $[\tau_n]_{\mathcal{A}} \supseteq B(b_n, 2^{-k_n}) \supseteq [\tau_{n+1}]_{\mathcal{A}} \supseteq \dots$. For all n , we have $\tau_n \prec \text{name}_{\mathcal{A}}(x)$, then $x \in [\tau_n]_{\mathcal{A}}$ for all n . So $x = \lim_{n \rightarrow \infty} b_n = S(\text{name}_{\mathcal{A}}(x))$. Since $S^{-1}([\sigma]_{\mathcal{A}}) = S^{-1}(\text{name}_{\mathcal{A}}^{-1}([\sigma]^{\prec})) = [\sigma]^{\prec}$ $\mu_{\mathcal{A}}$ -a.e., S is a measure-preserving map, and hence a morphism. Therefore, $\text{name}_{\mathcal{A}}$ is an isomorphism. \square

These last two propositions show that there is a one-to-one correspondence between cell decompositions \mathcal{A} of a space (\mathcal{X}, μ) and isomorphisms of the form $T: (\mathcal{X}, \mu) \rightarrow (2^{\omega}, \nu)$. This next proposition shows a further one-to-one correspondence between isomorphisms $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ and $S: (2^{\omega}, \mu_{\mathcal{A}}) \rightarrow (2^{\omega}, \nu_{\mathcal{B}})$.

Proposition 7.8 (Pairs of cell decompositions to isomorphisms). *Let (\mathcal{X}, μ) and (\mathcal{Y}, ν) be computable probability spaces with cell decompositions \mathcal{A} and \mathcal{B} . Let $\mu_{\mathcal{A}}$ be as in Proposition 7.7, and similarly for $\nu_{\mathcal{B}}$. Then for every isomorphism $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ there is an a.e. unique isomorphism $S: (2^{\omega}, \mu_{\mathcal{A}}) \rightarrow (2^{\omega}, \nu_{\mathcal{B}})$ and vice versa, such that S maps $\text{name}_{\mathcal{A}}(x)$ to $\text{name}_{\mathcal{B}}(T(x))$ for μ -a.e. $x \in \mathcal{X}$. In other words the following diagram commutes for μ -a.e. $x \in \mathcal{X}$.*

$$\begin{array}{ccc} (\mathcal{X}, \mu) & \xrightarrow{\text{name}_{\mathcal{A}}} & (2^{\omega}, \mu_{\mathcal{A}}) \\ \downarrow T & & \downarrow S \\ (\mathcal{Y}, \nu) & \xrightarrow{\text{name}_{\mathcal{B}}} & (2^{\omega}, \nu_{\mathcal{B}}) \end{array}$$

Further we have the following.

- (1) If (\mathcal{X}, μ) equals (\mathcal{Y}, ν) , then T is the identity isomorphism precisely when S is the isomorphism which maps $\text{name}_{\mathcal{A}}(x)$ to $\text{name}_{\mathcal{B}}(x)$.
- (2) Conversely, S is the identity isomorphism (and hence $\mu_{\mathcal{A}}$ equals $\nu_{\mathcal{B}}$) precisely when $\mathcal{A} = T^{-1}(\mathcal{B})$ (as in Proposition 7.6).

Proof. Given T , let $S = \text{name}_{\mathcal{A}}^{-1} \circ T \circ \text{name}_{\mathcal{B}}$, and similarly to get T from S . Then the diagram clearly commutes. A.e. uniqueness follows since the maps are isomorphisms.

If $\mathcal{A} = T^{-1}(\mathcal{B})$ is induced by T , then by Proposition 7.6, $\text{name}_{\mathcal{A}}(x) = \text{name}_{\mathcal{B}}(T(x))$ which makes S the identity map. But since S is an isomorphism, $\mu_{\mathcal{A}}$ and $\nu_{\mathcal{B}}$ must be the same measure.

The rest follows from “diagram chasing”. \square

Now we can show computable randomness is preserved by isomorphisms.

Theorem 7.9. *Isomorphisms preserve computable randomness. Namely, given an isomorphism $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$, then $x \in \mathcal{X}$ is computably random if and only if $T(x)$ is.*

Proof. Assume $T(x)$ is not computably random. Fix an isomorphism $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$. Let \mathcal{B} be a cell decomposition of (\mathcal{Y}, ν) . Take a bounded Martin-Löf test (U_n) on (\mathcal{Y}, ν) with bounding measure κ with respect to \mathcal{B} which covers $T(x)$. By Proposition 7.6 there is a cell decomposition $\mathcal{A} = T^{-1}(\mathcal{B})$ on (\mathcal{X}, μ) such that $[\sigma]_{\mathcal{A}} = T^{-1}([\sigma]_{\mathcal{B}})$ for all $\sigma \in 2^{<\omega}$. Define $V_n = T^{-1}(U_n)$. Then (V_n) is a bounded Martin-Löf test on (\mathcal{X}, μ) bounded by the same measure κ with respect to \mathcal{A} . Indeed,

$$\mu(V_n \cap [\sigma]_{\mathcal{A}}) = \nu(U_n \cap [\sigma]_{\mathcal{B}}) \leq 2^{-n} \kappa(\sigma).$$

Also, (V_n) covers x , hence x is not computably random. \square

Using Theorem 7.9, we can explore computable randomness on various spaces.

Example 7.10 (Computably random vectors). Let $([0, 1]^d, \lambda)$ be the cube $[0, 1]^d$ with the Lebesgue measure. The following is a natural isomorphism from $([0, 1]^d, \lambda)$ to $(2^\omega, \lambda)$. First, represent $(x_1, \dots, x_d) \in [0, 1]^d$ by the binary sequence of each component; then interleave the binary sequences. By Theorem 7.9, (x_1, \dots, x_d) is computably random if and only if the sequence of interleaved binary sequences is computably random. (This definition of computable randomness on $[0, 1]^d$ was proposed by Brattka, Miller and Nies [12].)

Example 7.11 (Base invariance). Let λ_3 be the uniform measure on 3^ω . Consider the natural isomorphism $T_{2,3}: (2^\omega, \lambda) \rightarrow (3^\omega, \lambda_3)$ which identifies the binary and ternary expansions of a real. This is an a.e. computable isomorphism, so $x \in [0, 1]$ is computably random if and only if $T_{2,3}(x)$ is computably random. We say a randomness notion (defined on (b^ω, λ_b) for all $b \geq 2$, see Example 5.9) is **BASE INVARIANT** if this property holds for all base pairs b_1, b_2 .

Example 7.12 (Computably random closed set). Consider the space $\mathcal{F}(2^\omega)$ of closed sets of 2^ω . This space has a topology called the Fell topology. The subspace $\mathcal{F}(2^\omega) \setminus \{\emptyset\}$ can naturally be identified with trees on $\{0, 1\}$ with no dead branches. Barmpalias et al. [3] gave a natural construction of these trees from ternary strings in 3^ω . Axon [2] showed the corresponding map $T: 3^\omega \rightarrow \mathcal{F}(2^\omega) \setminus \{\emptyset\}$ is a homeomorphism between 3^ω and the Fell topology restricted to $\mathcal{F}(2^\omega) \setminus \{\emptyset\}$. Hence $\mathcal{F}(2^\omega) \setminus \{\emptyset\}$ can be represented as a computable Polish space, and the probability measure on $\mathcal{F}(2^\omega) \setminus \{\emptyset\}$ induced by T can be represented as a computable probability measure. Since T is an a.e. computable isomorphism, the computably random closed sets of this space are then the ones whose corresponding trees are constructed from computably random ternary strings in 3^ω .

Example 7.13 (Computably random structures). The last example can be extended to a number of random structures—infinite random graphs, Markov processes, random walks, random matrices, Galton-Watson processes, etc. The main idea is as follows. Assume (Ω, P) is a computable probability space (the sample space), \mathcal{X} is the space of structures, and $T: (\Omega, P) \rightarrow \mathcal{X}$ is an a.e. computable map (a random structure). This induces a measure μ on \mathcal{X} (the distribution of T). If, moreover, T is an a.e. computable isomorphism between (Ω, P) and (\mathcal{X}, μ) , then the computably random structures of (\mathcal{X}, μ) are exactly the objects constructed from computably random points in (Ω, P) .

In this next theorem, an **ATOM** (or **POINT-MASS**) is a point with positive measure. An **ATOMLESS** probability space is one without atoms.

Theorem 7.14 (Hoyrup and Rojas [22]). *If (\mathcal{X}, μ) is an atomless computable probability space, then there is a isomorphism $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$. Further, T is computable from (\mathcal{X}, μ) .*

Corollary 7.15. *If (\mathcal{X}, μ) is an atomless computable probability space, then $x \in \mathcal{X}$ is computably random if and only if $T(x)$ is computably random for any (and all) isomorphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$.*

Proof. Follows from Theorems 7.9 and 7.14. \square

Example 7.16 (Computably random Brownian motion). Consider the space $C([0, 1])$ of continuous functions from $[0, 1]$ to \mathbb{R} endowed with the Wiener probability measure W (i.e. the measure of Brownian motion). The space $C([0, 1])$ with the uniform norm is a computable Polish space (where the simple points are the rational piecewise linear functions). The measure W is an atomless computable probability measure. Let $T: (2^\omega, \lambda) \rightarrow (C([0, 1]), W)$ be the isomorphism from Theorem 7.9. (Kjos-Hanssen and Nerode [24] construct a similar map φ directly for Brownian motion.) By Corollary 7.15, the computably random Brownian motions (i.e. the computably random points of $(C([0, 1]), W)$) are exactly the forward image of the computable random sequences under the map T .

Corollary 7.17. *Given a measure (\mathcal{X}, μ) with cell decomposition \mathcal{A} , $x \in X$ is computably random if and only if $\text{name}_{\mathcal{A}}(x)$ is computably random in $(2^\omega, \mu_{\mathcal{A}})$ where $\mu_{\mathcal{A}}(\sigma) = \mu([\sigma]_{\mathcal{A}})$.*

Proof. Use Proposition 7.7 and Theorem 7.9. □

8. GENERALIZING RANDOMNESS TO COMPUTABLE PROBABILITY SPACES

In this section, I explain the general method of this paper which generalizes a randomness notion from $(2^\omega, \lambda)$ to an arbitrary computable measure space.

Imagine we have an arbitrary randomness notion called X -RANDOMNESS defined on $(2^\omega, \lambda)$. (Here X is a place-holder for a name like “Schnorr” or “computable”; it has no relation to being random relative to an oracle.) The definition of X -random should either explicitly or implicitly assume we are working in the fair-coin measure. The method can be reduced to three steps.

Step 1: Generalize X -randomness to computable probability measures on 2^ω . This is self-explanatory, although not always trivial.

Step 2: Generalize X -randomness to computable probability spaces. There are three equivalent ways to do this for a computable probability space (\mathcal{X}, μ) .

- (1) Replace all instances of $[\sigma]^\prec$ with $[\sigma]_{\mathcal{A}}$, $x \upharpoonright n$ with $x \upharpoonright_{\mathcal{A}} n$, etc. in the definition of X -random from Step 1. Call this X^* -random with respect to \mathcal{A} . Then define $x \in \mathcal{X}$ to be X^* -random on (\mathcal{X}, μ) if it is X^* -random with respect to all cell decompositions \mathcal{A} (ignoring unrepresented points of \mathcal{A} and points in null cells—which are not even Kurtz random). (Compare with Definition 5.4.)
- (2) Define $x \in \mathcal{X}$ to be X^* -random on (\mathcal{X}, μ) if for each cell decomposition \mathcal{A} , $\text{name}_{\mathcal{A}}(x)$ is X -random on $(2^\omega, \mu_{\mathcal{A}})$, where $\mu_{\mathcal{A}}$ is given by $\mu_{\mathcal{A}}(\sigma) = \mu([\sigma]_{\mathcal{A}})$. (Compare with Corollary 7.17.)
- (3) Define $x \in \mathcal{X}$ to be X^* -random on (\mathcal{X}, μ) if for all isomorphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \nu)$ we have that $T(x)$ is X -random on $(2^\omega, \nu)$. (Compare with Theorem 7.9.)

The description of (1) is a bit vague, but when done correctly it is the most useful definition. The definition given by (1) should be equivalent to that given by (2) because (1) is essentially about \mathcal{A} -names. To see that (2) and (3) give the same definition, use Propositions 7.6 and 7.7, which show that isomorphisms to 2^ω are maps to \mathcal{A} -names and vice versa.

Step 3: Verify that the new definition is consistent with the original. It may be that on $(2^\omega, \lambda)$ the class of X^* -random points is strictly smaller than the class of the original X -random points. There are three equivalent techniques to show that X^* -randomness on 2^ω is equivalent to X -randomness. The three techniques are related to the three definitions from Step 2.

- (1) Show the definition of X^* -randomness is invariant under the choice of cell decomposition. (Compare with Theorem 5.7.)
- (2) Show that for every two cell decompositions \mathcal{A} and \mathcal{B} , the \mathcal{A} -name of x is X -random on $(2^\omega, \mu_{\mathcal{A}})$ if and only if the \mathcal{B} -name is X -random on $(2^\omega, \mu_{\mathcal{B}})$. (Compare with Corollary 7.17.)
- (3) Show that X -randomness is invariant under all isomorphisms from $(2^\omega, \mu)$ to $(2^\omega, \nu)$. (Compare with Theorem 7.9.)

Again, these three approaches are equivalent. Assuming the definition is stated correctly, (1) and (2) say the same thing.

To see that (3) implies (2), assume X -randomness is invariant under isomorphisms on 2^ω . Consider two cell decompositions \mathcal{A} and \mathcal{B} of the same space (\mathcal{X}, μ) . By Proposition 7.8(1), there is an isomorphism $S: (2^\omega, \mu_{\mathcal{A}}) \rightarrow (2^\omega, \mu_{\mathcal{B}})$ which maps \mathcal{A} -names to \mathcal{B} -names, i.e. this diagram commutes.

$$\begin{array}{ccc} (\mathcal{X}, \mu) & \xrightarrow{\text{name}_{\mathcal{A}}} & (2^\omega, \mu_{\mathcal{A}}) \\ & \searrow \text{name}_{\mathcal{B}} & \downarrow S \\ & & (2^\omega, \mu_{\mathcal{B}}) \end{array}$$

Since S preserves X -randomness, $\text{name}_{\mathcal{A}}(x)$ is X -random on $(2^\omega, \mu_{\mathcal{A}})$ if and only if $\text{name}_{\mathcal{B}}(x)$ is X -random on $(2^\omega, \mu_{\mathcal{B}})$.

To see that (2) implies (3), assume that (2) holds. Consider an isomorphism $S: (2^\omega, \mu) \rightarrow (2^\omega, \nu)$. Let (\mathcal{X}, κ) be any space isomorphic to $(2^\omega, \mu)$. Then (\mathcal{X}, κ) is also isomorphic to $(2^\omega, \nu)$. So there are isomorphisms T_1 and T_2 such that this diagram commutes.

$$\begin{array}{ccc} (\mathcal{X}, \kappa) & \xrightarrow{T_1} & (2^\omega, \mu) \\ & \searrow T_2 & \downarrow S \\ & & (2^\omega, \nu) \end{array}$$

By Proposition 7.6 there are two cell decompositions \mathcal{A} and \mathcal{B} on (\mathcal{X}, κ) such that $T_1 = \text{name}_{\mathcal{A}}$ and $(2^\omega, \mu) = (2^\omega, \kappa_{\mathcal{A}})$. The same holds for \mathcal{B} and ν . Then we have this commutative diagram.

$$\begin{array}{ccc} (\mathcal{X}, \kappa) & \xrightarrow{\text{name}_{\mathcal{A}}} & (2^\omega, \kappa_{\mathcal{A}}) \\ & \searrow \text{name}_{\mathcal{B}} & \downarrow S \\ & & (2^\omega, \kappa_{\mathcal{B}}) \end{array}$$

Consider any \mathbf{X} -random $y \in (2^\omega, \kappa_{\mathcal{A}})$. It is the \mathcal{A} -name of some $x \in (\mathcal{X}, \kappa)$, in other words $y = \text{name}_{\mathcal{A}}(x)$. By (2), we also have that $\text{name}_{\mathcal{B}}(x)$ is \mathbf{X} -random. So S preserves \mathbf{X} -randomness.

Notice that Step 3 implies that some randomness notions cannot be generalized without making the set of randoms smaller. This is because they are not invariant under isomorphisms between computable probability measures on 2^ω . Yet, even when the \mathbf{X}^* -randoms are a proper subclass of the \mathbf{X} -randoms, the \mathbf{X}^* randoms are an interesting class of randomness. In particular we have the following.

Proposition 8.1. *\mathbf{X}^* -randomness is invariant under isomorphisms.*

In some sense the \mathbf{X}^* -randoms are the largest such subclass of the \mathbf{X} -randoms. (One must be careful how to say this, since \mathbf{X} -randomness is only defined on measures $(2^\omega, \mu)$.)

Proof. Let $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{Y}, \nu)$ be an isomorphism and let $x \in (\mathcal{X}, \mu)$ be \mathbf{X}^* -random. Let \mathcal{B} be a arbitrary cell decomposition of (\mathcal{Y}, ν) . Since \mathcal{B} is arbitrary, it is enough to show that $\text{name}_{\mathcal{B}}(T(x))$ is \mathbf{X} -random in $(2^\omega, \nu_{\mathcal{B}})$. By Proposition 7.6 and Proposition 7.8 (2) we have a cell decomposition \mathcal{A} on (\mathcal{X}, μ) such that $(2^\omega, \mu_{\mathcal{A}}) = (2^\omega, \nu_{\mathcal{B}})$ and the following diagram commutes.

$$\begin{array}{ccc} (\mathcal{X}, \mu) & \xrightarrow{\text{name}_{\mathcal{A}}} & (2^\omega, \mu_{\mathcal{A}}) \\ \downarrow T & & \parallel \\ (\mathcal{Y}, \nu) & \xrightarrow{\text{name}_{\mathcal{B}}} & (2^\omega, \nu_{\mathcal{B}}) \end{array}$$

Since $x \in (\mathcal{X}, \mu)$ is \mathbf{X}^* -random, $\text{name}_{\mathcal{A}}(x)$ is \mathbf{X} -random in $(2^\omega, \mu_{\mathcal{A}}) = (2^\omega, \nu_{\mathcal{B}})$. Since the diagram commutes, $\text{name}_{\mathcal{B}}(T(x))$ is also \mathbf{X} -random in $(2^\omega, \nu_{\mathcal{B}})$. Since \mathcal{B} is arbitrary, x is \mathbf{X} -random. \square

In the case that (\mathcal{X}, μ) is an atomless computable probability measure, we could instead define $x \in \mathcal{X}$ to be \mathbf{X}^\star -random if $T(x)$ is random for all isomorphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$. We can then skip Step 1, and in Step 3 it is enough to check that \mathbf{X} -randomness is invariant under automorphisms of $(2^\omega, \lambda)$. Similarly, \mathbf{X}^\star -randomness would be invariant under isomorphisms.

9. BETTING STRATEGIES AND KOLMOGOROV-LOVELAND RANDOMNESS

In the next two sections I consider how the method of Section 8 can be applied to Kolmogorov-Loveland randomness, which is also defined through a betting strategy on the bits of the string.

Call a betting strategy on bits **NONMONOTONIC** if the gambler can decide at each stage which coin flip to bet on. For example, maybe the gambler first bets on the 5th bit. If it is 0, then he bets on the 3rd bit; if it is

1, he bets on the 8th bit. (Here, and throughout this paper we still assume the gambler cannot bet more than what is in his capital, i.e. he cannot take on debt.) A string $x \in 2^\omega$ is **KOLMOGOROV-LOVELAND RANDOM** or **NONMONOTONICALLY RANDOM** (in $(2^\omega, \lambda)$) if there is no computable nonmonotonic betting strategy on the bits of the string which succeeds on x .

Indeed, this gives a lot more freedom to the gambler and leads to a strictly stronger notion than computable randomness. While it is easy to show that every Martin-Löf random is Kolmogorov-Loveland random, the converse is a difficult open question.

Question 9.1. *Is Kolmogorov-Loveland randomness the same as Martin-Löf randomness?*

On one hand, there are a number of results that show Kolmogorov-Loveland randomness is very similar to Martin-Löf randomness. On the other hand, it is not even known if Kolmogorov-Loveland randomness is base invariant, and it is commonly thought that Kolmogorov-Loveland randomness is strictly weaker than Martin-Löf randomness. For the most recent results on Kolmogorov-Loveland randomness see [14, Section 7.5], [32, Section 7.6], and [5, 23, 29].

In this section I will ask what type of randomness one gets by applying the method of Section 8 to Kolmogorov-Loveland randomness. The result is Martin-Löf randomness. However, this does not prove that Kolmogorov-Loveland randomness is the same as Martin-Löf randomness, since I leave as an open question whether Kolmogorov-Loveland randomness (naturally extended to all computable probability measures on 2^ω) is invariant under isomorphisms. The presentation of this section follows the three-step method of Section 8.

9.1. Step 1: Generalize to other computable probability measures μ on 2^ω . Kolmogorov-Loveland randomness can be naturally extended to computable probability measures on 2^ω . Namely, bet as usual, but adjust the payoffs to be fair. For example, if the gambler wagers 1 unit of money to bet that $x(4) = 1$ (i.e. the 4th bit is 1) after seeing that $x(2) = 1$ and $x(6) = 0$, then if he wins, the fair payoff is

$$\frac{\mu(x(4) = 0 \mid x(2) = 1, x(6) = 0)}{\mu(x(4) = 1 \mid x(2) = 1, x(6) = 0)}.$$

where $\mu(A \mid B) = \mu(A \cap B)/\mu(B)$ represents the conditional probability of A given B . If the gambler loses, he loses his unit of money.

(Note, we could also allow the gambler to bet on a bit he has already seen. Indeed, he will not win any money. This would, however, introduce “partial randomness” since the gambler could delay betting on a new bit. Nonetheless, Merkle [27] showed that partial Kolmogorov-Loveland randomness is the same as Kolmogorov-Loveland randomness.)

As with computable randomness, we must address division by zero. The gambler is not allowed to bet on a bit if it has probability zero of occurring (conditioned on the information already known). Instead we just declare the elements of such null cylinder sets to be not random.

9.2. Step 2: Generalize Kolmogorov-Loveland randomness to computable probability measures.

Pick a computable probability measure (\mathcal{X}, μ) with generator $\mathcal{A} = (A_n)$. Following the second step of the method in Section 8, the gambler bets on the bits of the \mathcal{A} -name of x . A little thought reveals that what the gambler is doing when she bets that the n th bit of the \mathcal{A} -name is 1 is betting that $x \in A_n$. For any generator \mathcal{A} , if we add more a.e. decidable sets to \mathcal{A} , it is still a generator. Further, since we are not necessary betting on all the sets in \mathcal{A} , we do not even need to assume \mathcal{A} is anything more than a collection of a.e. decidable sets. (This is the key difference between computable randomness.)

Hence, we may think of the betting strategy as follows. The gambler chooses some a.e. decidable set A and bets that $x \in A$ (or x has property A). (Again, the gambler must know that $\mu(A) > 0$ before betting on it.) Then if she wins, she gets a fair payout, and if she loses, she loses her bet. Call such a strategy a **COMPUTABLE BETTING STRATEGY**. Call the resulting randomness **BETTING RANDOMNESS**. (A more formal definition is given in Remark 9.4.)

I argue that betting randomness is the most general randomness notion that can be described by a finitary fair-game betting scenario with a “computable betting strategy.” Indeed, consider these three basic properties of such a game:

- (1) The gambler must be able determine (almost surely) some property of x that she is betting on, and this determination must be made with only the information about x that she has gained during the game.
- (2) A bookmaker must be able to determine (almost surely) if this property holds of x or not.
- (3) If the gambler wins, the bookmaker must be able to determine (almost surely) the fair payoff amount.

The only way to satisfy (2) is if the property is a.e. decidable. Then (3) follows since a.e. decidable sets have finite descriptions and their measures are computable. To satisfy (1), the gambler must be able to compute the a.e. decidable set only knowing the results of her previous bets. This is exactly the computable betting strategy defined above.¹

Now recall Schnorr's Critique that Martin-Löf randomness does not have a "computable-enough" definition. The definition Schnorr had in mind was a betting scenario. In particular, Schnorr gave a martingale characterization of Martin-Löf randomness that is the same as that of computable randomness, except the martingales are only lower semicomputable [35] (see also [14, 32]). If Martin-Löf randomness equals Kolmogorov-Loveland randomness, then some believe that this will give a negative answer to Schnorr's Critique; namely, we will have found a computable betting strategy that describes Martin-Löf randomness. While, there is some debate as to what Schnorr meant by his critique (and whether he still agrees with it), we think the following is a worthwhile question.

Can Martin-Löf randomness be characterized using a finitary fair-game betting scenario with a "computable betting strategy"?

The answer turns out to be yes. As this next theorem shows, betting randomness is equivalent to Martin-Löf randomness. Hitchcock and Lutz [21] defined a generalization of martingales (as in the type used to define computable randomness on 2^ω) called martingale processes. In the terminology of this paper, a MARTINGALE PROCESS is basically a computable betting strategy on 2^ω with the fair-coin measure which bets on decidable sets (i.e. finite unions of basic open sets). Merkle, Mihailović and Slaman [28] showed that Martin-Löf randomness is equivalent to the randomness characterized by martingale processes. The proof of this next theorem is basically the Merkle et al. proof.²

Theorem 9.2. *Betting randomness and Martin-Löf randomness are the same.*

Proof. Fix a computable probability space (\mathcal{X}, μ) . To show Martin-Löf randomness implies betting randomness, we use a standard argument which was employed by Hitchcock and Lutz [21] for martingale processes. Assume $x \in \mathcal{X}$ is not betting random. Namely, there is some computable betting strategy \mathcal{B} which succeeds on x . Without loss of generality, the starting capital of \mathcal{B} may be assumed to be 1. Let $U_n = \{x \in \mathcal{X} \mid \mathcal{B} \text{ wins at least } 2^n \text{ on } x\}$. Each U_n is uniformly Σ_1^0 in n , and by a standard result in martingale theory $\mu(U_n) \leq C2^{-n}$ where $C = 1$ is the starting capital.³ Hence (U_n) is a Martin-Löf test which covers x , and x is not Martin-Löf random.

For the converse, the argument is basically the Merkle, Mihailović and Slaman [28] proof for martingale processes.

First, let us prove a fact. Assume a gambler starts with a capital of 1 and $U \subset \mathcal{X}$ is some Σ_1^0 set such that $\mu(U) \leq 1/2$. Then there is a computable way that the gambler can bet on an unknown $x \in \mathcal{X}$ such that he doubles his capital (to 2) if $x \in U$ (actually, some Σ_1^0 set a.e. equal to U). The strategy is as follows. Choose a cell decomposition \mathcal{A} of (\mathcal{X}, μ) . Since U is Σ_1^0 , by Proposition 4.7 there is a c.e., prefix-free set $\{\sigma_i\}$ of finite strings such that $U = \bigcup_i [\sigma_i]_{\mathcal{A}}$ a.e. We may assume $\mu([\sigma_i]_{\mathcal{A}}) > 0$ for all i . To start, the gambler bets on the set $[\sigma_0]_{\mathcal{A}}$ with a wager such that if he wins, his capital is 2. If he wins, he is done. If he loses, then he bets on the set $[\sigma_1]_{\mathcal{A}}$, and so on. Since the set $\{\sigma_i\}$ may be finite, the gambler may not have a set to

¹In the three properties we did not consider the possibility of betting on a collection of three or more pair-wise disjoint events simultaneously. This is not an issue since one may break up the betting and bet on each event individually (see Example 5.9). There is also a more general possibility of having a computable or a.e. computable wager function over the space \mathcal{X} . This can be made formal using the martingales in probability theory, but it turns out that it does not change the randomness characterized by such a strategy. By an unpublished result of Ed Dean [personal communication], any L^1 -bounded layerwise-computable martingale converges on Martin-Löf randomness (which, as we will see, is equivalent to betting randomness).

²Downey and Hirschfeld [14, footnote on p. 269] also remark that the Merkle et al. result gives a possible answer to Schnorr's critique.

³This follows from Kolmogorov's inequality (proved by Ville, see [14, Theorem 6.3.3 and Lemma 6.3.15(ii)]) which is a straight-forward application of Doob's submartingale inequality (see for example [40, Section 16.4]).

bet on at certain stages. This is not an issue, since he may just bet on the whole space. This is functionally equivalent to not betting at all since he wins no money.

The only difficulty now is showing that his capital remains nonnegative. Merkle et al. leave this an exercise for the reader; I give an intuitive argument. It is well-known in probability theory that in a betting strategy one can combine bets for the same effect. (Formally, this is the martingale stopping theorem—see [40].) Hence instead of separately betting on $[\sigma_0]_{\mathcal{A}}, \dots, [\sigma_k]_{\mathcal{A}}$ the gambler will have the same capital as if he just bet on the union $[\sigma_0]_{\mathcal{A}} \cup \dots \cup [\sigma_k]_{\mathcal{A}}$. In the later case, the proper wager would be.

$$\frac{\mu([\sigma_0]_{\mathcal{A}} \cup \dots \cup [\sigma_k]_{\mathcal{A}})}{\mu(\mathcal{X} \setminus ([\sigma_0]_{\mathcal{A}} \cup \dots \cup [\sigma_k]_{\mathcal{A}}))} \leq 1,$$

The inequality follows from

$$\mu([\sigma_0]_{\mathcal{A}} \cup \dots \cup [\sigma_k]_{\mathcal{A}}) \leq 1/2 \leq \mu(\mathcal{X} \setminus ([\sigma_0]_{\mathcal{A}} \cup \dots \cup [\sigma_k]_{\mathcal{A}})).$$

Hence the gambler never wagers (and so loses) more than his starting capital of 1.

Now, assume $z \in \mathcal{X}$ is not Martin-Löf random. Let (U_k) be a Martin-Löf test which covers z . We may assume (U_n) is decreasing. The betting strategy will be as follows which bets on some $x \in \mathcal{X}$. Since $\mu(U_1) < 1/2$ we can start with the computable betting strategy above which will reach a capital of 2 if $x \in U_1$. (Recall, we are not actually betting on U_1 , but the a.e. equal set $\bigcup_i [\sigma_i]_{\mathcal{A}}$. This is not an issue, since the difference is a null Σ_2^0 set. If x is in the difference, then x is not computably random, and so not betting random.)

Now, if the capital of 2 is never reached then $x \notin U_1$ and x is random. However, if the capital of 2 is reached (in a finite number of steps) then we know that $x \in [\sigma]_{\mathcal{A}}$ for some $\sigma = \sigma_i$ (and no other). Further, by the assumptions in the above construction, $\mu([\sigma]_{\mathcal{A}}) > 2^{-k}$ for some k . Then we can repeat the first step, but now we bet that $x \in U_{k+1}$ and attempt to double our capital to 4. Since $\mu(U_{k+1} \mid [\sigma]_{\mathcal{A}}) \leq 1/2$, the capital will remain positive.

Continuing this strategy for capitals of 8, 16, 32, \dots we have a computable betting strategy. If this strategy succeeds on x , then $x \in U_k$ for infinitely many k . Hence x is covered by (U_k) and is not Martin-Löf random. \square

Remark 9.3. Since there is a universal Martin-Löf test (U_k) , there is a universal computable betting strategy. (The null Σ_2^0 set of exceptions can be handled by being more careful. Choose \mathcal{A} to be basis for the topology, and combine the null cells $[\sigma]_{\mathcal{A}}$ with non-null cells $[\tau]_{\mathcal{A}}$.) However, note that this universal strategy is very different from that of Kolmogorov-Loveland randomness. This is the motivation for the next section.

It is also possible to characterize Martin-Löf randomness by computable randomness. First I give a more formal definition of computable betting strategy.

Remark 9.4. Represent a computable betting strategy as follows. There is a computably indexed family of a.e. decidable sets $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$. These represent the sets being bet on after the wins/loses characterized by $\sigma \in 2^{<\omega}$. From this, we have a computably indexed family $\{B_\sigma\}_{\sigma \in 2^{<\omega}}$ defined recursively by $B_\epsilon = \mathcal{X}$, $B_{\sigma 1} = B_\sigma \cap A_\sigma$ and $B_{\sigma 0} = B_\sigma \cap (\mathcal{X} \setminus A_\sigma)$. This represents the known information after the wins/loses characterized by $\sigma \in 2^{<\omega}$. It is easy to see that $B_{\sigma 0} \cap B_{\sigma 1} = \emptyset$ and $B_{\sigma 0} \cup B_{\sigma 1} = B_\sigma$ a.e. Then a computable betting strategy can be represented as a partial computable martingale $M: 2^{<\omega} \rightarrow [0, \infty)$ such that

$$M(\sigma 0)\mu(B_{\sigma 0}) + M(\sigma 1)\mu(B_{\sigma 1}) = M(\sigma)\mu(B_\sigma)$$

and $M(\sigma)$ is defined if and only if $\mu(B_{\sigma 0}) > 0$. Again, $M(\sigma)$ represents the capital after a state of σ wins/losses. Say the strategy SUCCEEDS on x if there is some strictly-increasing chain $\sigma_0 \prec \sigma_1 \prec \sigma_2 \prec \dots$ from $2^{<\omega}$ such that $\limsup_{n \rightarrow \infty} M(\sigma_n) = \infty$ and $x \in B_{\sigma_n}$ for all n . Then $x \in \mathcal{X}$ is betting random if there does not exist some $\{A_\sigma\}_{\sigma \in 2^{<\omega}}$ and M as above which succeed on x .

Lemma 9.5. *Fix a computable probability space (\mathcal{X}, μ) . For each computable betting strategy there is a computable probability measure ν on 2^ω , a morphism $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \nu)$, and a computable martingale M on $(2^\omega, \nu)$ such that if this betting strategy succeeds on x , then the martingale M succeeds on $T(x)$. Hence $T(x)$ is not computably random on $(2^\omega, \nu)$.*

Proof. Fix a computable betting strategy. Let $M: 2^{<\omega} \rightarrow [0, \infty)$ and $\{B_\sigma\}_{\sigma \in 2^{<\omega}}$ be the as in Remark 9.4. Then define $(2^\omega, \nu)$ by $\nu(\sigma) = \mu(B_\sigma)$. Also, let $T(x)$ map x to the $y \in 2^\omega$ such that $x \in B_{y|n}$ for all n . Then

T is a morphism, M also represents a martingale on $(2^\omega, \nu)$, and if the betting strategy succeeds on x then M succeeds on $T(x)$. \square

We now have the following characterizations of Martin-Löf randomness.

Corollary 9.6. *For a computable probability space (\mathcal{X}, μ) , the following are equivalent for $x \in \mathcal{X}$.*

- (1) x is Martin-Löf random.
- (2) No computable betting strategy succeeds on x (i.e. x is betting random).
- (3) For all isomorphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \nu)$, $T(x)$ is “Kolmogorov-Loveland random” on $(2^\omega, \nu)$ (i.e. the randomness from Section 9.1).
- (4) For all morphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \nu)$, $T(x)$ is computably random on $(2^\omega, \nu)$.

Proof. The equivalence of (1) and (2) is Theorem 9.2. (1) implies both (3) and (4) since morphisms preserve Martin-Löf randomness (Proposition 7.5).

(4) implies (2): Use Lemma 9.5. Assume x is not betting random. Then there is some morphism T such that $T(x)$ is not computably random.

(3) implies (2): Recall that the definition of betting randomness came from applying the method of Section 8 to Kolmogorov-Loveland randomness. By method (3) of Step 2 in Section 8, x is betting random if and only if (3) holds. (An alternate proof would be to modify Lemma 9.5.) \square

Corollary 9.7. *Computable randomness is not preserved under morphisms. (See comments after Proposition 7.5.)*

Proof. It is well-known that there is an $x \in 2^\omega$ which is computably random on $(2^\omega, \lambda)$ but not Martin-Löf random (see [14, 32]). Then by Corollary 9.6, there is some morphism T such that $T(x)$ is not computably random. \square

Corollary 9.7 was also proved by Bienvenu and Porter [7].

9.3. Step 3: Is the new definition consistent with the former? To show that Martin-Löf randomness equals Kolmogorov-Loveland randomness, we would need to show that “Kolmogorov-Loveland randomness” for all computable probability measures on Cantor space (as in Section 9.1) is preserved by isomorphisms. However, it is not even known if Kolmogorov-Loveland randomness on $(2^\omega, \lambda)$ is base invariant (see Examples 5.10 and 7.11), so I leave this as an open question.

Question 9.8. *Is Kolmogorov-Loveland randomness, as in Section 9.1, preserved under isomorphisms?*

10. ENDOMORPHISM RANDOMNESS

The generalization of Kolmogorov-Loveland randomness given in the last section was, in some respects, not very satisfying. In particular, the definition of Kolmogorov-Loveland randomness on $(2^\omega, \lambda)$ assumes each event being bet on is independent of all the previous events, and further has conditional probability $1/2$. Therefore, at the “end” of the gambling session, regardless of how much the gambler has won or lost, he knows what x is up to a measure-zero set (where x is the string being bet on). This is in contrast to the universal betting strategy given in the proof of Theorem 9.2 (see Remark 9.3), which only narrows x down to a positive measure set when x is Martin-Löf random.

In this section, I now give a new type of randomness which behaves more like Kolmogorov-Loveland randomness. This randomness notion can be defined using both morphisms and betting strategies.

Definition 10.1. Let (\mathcal{X}, μ) be a computable probability space. An ENDOMORPHISM on (\mathcal{X}, μ) is a morphism from (\mathcal{X}, μ) to itself. Say $x \in \mathcal{X}$ is ENDOMORPHISM RANDOM if for all endomorphisms $T: (\mathcal{X}, \mu) \rightarrow (\mathcal{X}, \mu)$, we have that $T(x)$ is computably random.

Notice the above definition is the same as that given in Corollary 9.6 (4), except that “morphism” is replaced with “endomorphism”.

If the space is atomless, we have an alternate characterization.

Proposition 10.2. *Let (\mathcal{X}, μ) be a computable probability space with no atoms. Then $x \in \mathcal{X}$ is endomorphism random if and only if for all morphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$, $T(x)$ is computably random.*

Proof. Use that there is an isomorphism from (\mathcal{X}, μ) to $(2^\omega, \lambda)$ (Theorem 7.14) and that isomorphisms preserve computable randomness (Theorem 7.9). \square

Also, we can define endomorphism randomness using computable betting strategies as in the previous section.

Definition 10.3. Let (\mathcal{X}, μ) be an atomless computable probability space. Consider a computable betting strategy \mathcal{B} . Let $\{A_\sigma\}_{\sigma \in 2^{<\omega}}, \{B_\sigma\}_{\sigma \in 2^{<\omega}}$ be as in Remark 9.4. Call the betting strategy \mathcal{B} **BALANCED** if it only bets on events with conditional probability $\frac{1}{2}$, conditioned on B_σ (the information known by the gambler at after the wins/loses given by σ). In other words, $\mu(A_\sigma | B_\sigma) = 1/2$. Call the betting strategy \mathcal{B} **EXHAUSTIVE** if $\mu(B_{\sigma_n}) \rightarrow 0$ for any strictly increasing chain $\sigma_0 < \sigma_1 < \dots$. In other words the measure of the information known about x approaches 0.

Theorem 10.4. Let (\mathcal{X}, μ) be an atomless computable probability space and $x \in \mathcal{X}$. The following are equivalent.

- (1) x is endomorphism random.
- (2) There does not exist a balanced computable betting strategy which succeeds on x .
- (3) There does not exist an exhaustive computable betting strategy which succeeds on x .

Proof. (3) implies (2) since balanced betting strategies are exhaustive. For (2) implies (1), assume x is not endomorphism random. Then there is some morphism $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$ such that $T(x)$ is not computably random. Hence there is a computable martingale M which succeeds on $T(x)$. We can also assume this martingale is rational valued, so it is clear what bit is being bet on. This martingale on $(2^\omega, \lambda)$ can be pulled back to a computable betting strategy on (\mathcal{X}, μ) (use the proof of Lemma 9.5, except in reverse). This betting strategy is balanced since M is a balanced “dyadic” martingale.

For (1) implies (3), assume there is some computable, exhaustive betting strategy which succeeds on x . Then from this strategy we can construct a morphism $S: (\mathcal{X}, \mu) \rightarrow ([0, 1], \lambda)$ recursively as follows. Each B_σ will be mapped to an open interval (a, b) of length $\mu(B_\sigma)$. First, map $S(B_\epsilon) = (0, 1)$. For the recursion step, assume $S(B_\sigma) = (a, b)$ of length $\mu(B_\sigma)$. Set $S(B_{\sigma 0}) = (a, a + \mu(B_{\sigma 0}))$ and $S(B_{\sigma 1}) = (a + \mu(B_{\sigma 0}), b)$. This function S is well-defined and computable since the betting strategy is exhaustive. Also, S is clearly measure-preserving, so it is a morphism. Then using the usual isomorphism from $([0, 1], \lambda)$ to $(2^\omega, \lambda)$, we can assume S is a morphism to $(2^\omega, \lambda)$. Moreover, the set of images $S(B_\sigma)$ describes a cell decomposition \mathcal{A} of $(2^\omega, \lambda)$, and the betting strategy can be pushed forward to give a martingale on $(2^\omega, \lambda)$ with respect to \mathcal{A} (similar to the proof of Lemma 9.5). \square

Now we can relate endomorphism randomness to Kolmogorov-Loveland randomness.

Corollary 10.5. On $(2^\omega, \lambda)$, endomorphism randoms are Kolmogorov-Loveland randoms.

Proof. Every nonmonotonic, computable betting strategy on bits is a balanced betting strategy. Hence every Kolmogorov-Loveland random is endomorphism random. \square

Corollary 10.6. Let (\mathcal{X}, μ) be a computable probability space with no atoms. Then $x \in \mathcal{X}$ is endomorphism random if and only if for all morphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$, $T(x)$ is Kolmogorov-Loveland random.

Proof. If x is endomorphism random on $(2^\omega, \lambda)$, then so is $T(x)$. By Corollary 10.5, $T(x)$ is Kolmogorov-Loveland random. If $T(x)$ is Kolmogorov-Loveland random for all morphisms $T: (\mathcal{X}, \mu) \rightarrow (2^\omega, \lambda)$, then $T(x)$ is computably random for all such T . Therefore, x is endomorphism random. \square

Corollary 10.7. Computable randomness is not preserved by endomorphisms.

Proof. It is well-known that there exists a computable random $x \in (2^\omega, \lambda)$ which is not Kolmogorov-Loveland random (see [14, 32]). Then x is not endomorphism random, and is not preserved by some endomorphism. \square

Also, clearly each betting random (i.e. each Martin-Löf random) is an endomorphism random.

I will add one more randomness notion. Say $x \in 2^\omega$ is **AUTOMORPHISM RANDOM** (on $(2^\omega, \lambda)$) if for all automorphisms $T: (2^\omega, \lambda) \rightarrow (2^\omega, \lambda)$, $T(x)$ is Kolmogorov-Loveland random. It is clear that on $(2^\omega, \lambda)$ we have.

(10.1) Martin-Löf \rightarrow Endomorphism

\rightarrow Automorphism \rightarrow Kolmogorov-Loveland

We now have a more refined version of Question 9.1.

Question 10.8. *Do any of the implications in formula (10.1) reverse?*⁴

11. FURTHER DIRECTIONS

Throughout this paper I was working with a.e. computable objects: a.e. decidable sets, a.e. decidable cell decompositions, a.e. computable morphisms, and Kurtz randomness—which as I showed, can be defined by a.e. computability. Recall a.e. decidable sets are only sets of μ -continuity, and a.e. computable morphisms are only a.e. continuous maps.

The “next level” is to consider the computable Polish spaces of measurable sets and measurable maps. The a.e. decidable sets and a.e. computable maps are dense in these spaces. Hence, in the definitions, one may replace a.e. decidable sets, a.e. decidable cell decompositions, a.e. computable morphisms, and Kurtz randomness with effectively measurable sets, decompositions into effectively measurable cells, effectively measurable measure-preserving maps, and Schnorr randomness. (This is closely related to the work of Pathak, Simpson and Rojas [33]; Miyabe [30]; Hoyrup and Rojas [personal communication]; and the author on “Schnorr layerwise-computability” and convergence for Schnorr randomness.) Indeed, the results of this paper remain true, even with those changes. However, some proofs change and I will give the results in a later paper.

An even more general extension would be to ignore the metric space structure all together. Any standard probability space can be described uniquely by the measures of an intersection-closed class of sets, or a π -system, which generates the sigma-algebra of the measure. From this, one can obtain a cell decomposition. In the case of a computable probability space (Definition 3.2), each a.e. decidable generator closed under intersections is a π -system. The definition of computable randomness on such a general space would be the analog of the definition in this paper.

In particular, this would allow one to define computable randomness on effective topological spaces with measure [20]. In this case the π -system is the topological basis. This also allows one to define Schnorr, Martin-Löf, and weak-2 randomness as well, namely replace, say, Σ_1^0 sets in the definition with effective unions of sets in the π -system. This agrees with most definitions of, say, Martin-Löf randomness in the literature.⁵

Using π -systems also allows one to define “abstract” measure spaces without points. The computable randoms then become “abstract points” given by generic ultrafilters on the Boolean algebra of measurable sets *a la* Solovay forcing.

Another possible generalization is to non-computable probability spaces (on computable Polish spaces). This has been done by Levin [25] and extended by others (see [17, 4]) for Martin-Löf randomness in a natural way using UNIFORM TESTS which are total computable functions from measures to tests. Possibly a similar approach would work for computable randomness. For example, on 2^ω , a uniform test for computable randomness would be a total computable map $\mu \mapsto \nu$ where ν is the bounding measure for μ . This map is enough to define a uniform martingale test for each μ given by $\nu(\sigma)/\mu(\sigma)$. (I showed in Section 2 that this martingale is uniformly computable.) Uniform tests for Schnorr and computable randomness have been used by Miyabe [31].

Also, what other applications for a.e. decidable sets are there in effective probability theory? The method of Section 8 basically allows one to treat every computable probability space as the Cantor space. It is already known that the indicator functions of a.e. decidable sets can be used to define L^1 -computable functions [30].

However, when it comes to defining classes of points, the method of Section 8 is specifically for defining *random* points since such a definition must be a subclass of the Kurtz randoms. Under certain circumstances, however, one may be able to use related methods to generalize other definitions. For example, is the following a generalization of K-triviality to arbitrary computable probability spaces? Let $K = K_M$ where M is a

⁴Recently, and independently of my work, Tomislav Petrovic has claimed that there are two balanced betting strategies on $(2^\omega, \lambda)$ such that if a real x is not Martin-Löf random, then at least one of the two strategies succeeds on x . In particular, Petrovic’s result, which is in preparation, would imply that endomorphism randomness equals Martin-Löf randomness. Further, via the proof of Theorem 10.4, this result would extend to every atomless computable probability space.

⁵There are some authors [2, 20] that define Martin-Löf randomness via open covers, even for non-regular topological spaces. This will not necessarily produce a measure-one set of random points, where as my method will. All these methods agree for spaces with an effective regularity condition.

universal prefix-free machine. Recall, a string $x \in 2^\omega$ is K-TRIVIAL (on $(2^\omega, \lambda)$) if there is some b such that

$$\forall n \ K(x \upharpoonright n) \leq K(n) + b$$

where $K(n) = K(0^n)$ and 0^n is the string of 0's of length n . Taking a clue from Section 6, call a point $x \in (\mathcal{X}, \mu)$ K-TRIVIAL if there is some cell decomposition \mathcal{A} and some b such that for all n ,

$$K(x \upharpoonright_{\mathcal{A}} n) < K(-\log \mu([x \upharpoonright_{\mathcal{A}} n]_{\mathcal{A}})) + b.$$

(Here we assume $K(\infty) = \infty$.) Does the \mathcal{A} -name or Cauchy-name of x satisfy the other nice degree theoretic properties of K-triviality, such as being low-for- (\mathcal{X}, μ) -random? (Here I say a Turing degree \mathbf{d} is low-for- (\mathcal{X}, μ) -random if when used as an oracle, \mathbf{d} does not change the class of Martin-Löf randoms in (\mathcal{X}, μ) . Say a point $x \in (\mathcal{X}, \mu)$ is low-for- (\mathcal{X}, μ) -random if its Turing degree is.)

If it is a robust definition, how does it relate to the definition of Melnikov and Nies [26] generalizing K-triviality to computable Polish spaces (as opposed to probability spaces)? I conjecture that their definition is equivalent to being low-for- (\mathcal{X}, μ) -random on every computable probability measure μ of \mathcal{X} .

Last, isomorphisms and morphisms offer a useful tool to classify randomness notions. One may ask what randomness notions (defined for all computable probability measures on 2^ω) are invariant under morphisms or isomorphisms? By Proposition 7.5, Martin-Löf, Schnorr, and Kurtz randomness are invariant under morphisms. (This can easily be extended to n -randomness, weak n -randomness, and difference randomness. See [14, 32] for definitions.) However, by Proposition 9.6 (4), there is no randomness notion between Martin-Löf randomness and computable randomness that is invariant under morphisms. Is there such a randomness notion between Schnorr randomness and Martin-Löf randomness? Further, by Theorem 7.9 computable randomness is invariant under isomorphisms. André Nies pointed out to me that this is not true of partial computable randomness since it is not invariant under permutations [5]. In general what can be said of full-measure, isomorphism-invariant sets of a computable probability space (\mathcal{X}, μ) ? The notions of randomness connected to computable analysis will most likely be the ones that are invariant under isomorphisms.⁶

11.1. Acknowledgments. I would like to thank Mathieu Hoyrup and Cristóbal Rojas for suggesting to me the use of isomorphisms to define computable randomness outside Cantor space. I would also like to thank Jeremy Avigad, Laurent Bienvenu, Peter Gács, and Mathieu Hoyrup for helpful comments on a draft of this paper, as well as Joseph Miller and André Nies for helpful comments on this research.

REFERENCES

- [1] Jeremy Avigad. Uniform distribution and algorithmic randomness. *Journal of Symbolic Logic*, 78(1):334–344, 2013.
- [2] Logan M. Axon. *Algorithmically random closed sets and probability*. ProQuest LLC, Ann Arbor, MI, 2010. Thesis (Ph.D.)—University of Notre Dame.
- [3] George Barmpalias, Paul Brodhead, Douglas Cenzer, Seyyed Dashti, and Rebecca Weber. Algorithmic randomness of closed sets. *J. Logic Comput.*, 17(6):1041–1062, 2007.
- [4] Laurent Bienvenu, Peter Gács, Mathieu Hoyrup, Cristóbal Rojas, and A. Shen. Algorithmic tests and randomness with respect to a class of measures. *Proceedings of the Steklov Institute of Mathematics*, 274:41–102, 2011.
- [5] Laurent Bienvenu, Rupert Hölzl, Thorsten Kräling, and Wolfgang Merkle. Separations of non-monotonic randomness notions. *J. Logic and Computation*, 2010.
- [6] Laurent Bienvenu and Wolfgang Merkle. Constructive equivalence relations on computable probability measures. *Ann. Pure Appl. Logic*, 160(3):238–254, 2009.
- [7] Laurent Bienvenu and Christopher Porter. Strong reductions in effective randomness. *Theoret. Comput. Sci.*, 459:55–68, 2012.
- [8] Errett Bishop and Douglas Bridges. *Constructive analysis*. 279. Springer-Verlag, 1985.
- [9] Volker Bosserhoff. Notions of probabilistic computability on represented spaces. *J. Universal Computer Science*, 14(6):956–995, 2008.
- [10] Vasco Brattka. Computable versions of Baire’s category theorem. In *Mathematical foundations of computer science, 2001 (Mariánské Lázně)*, volume 2136 of *Lecture Notes in Comput. Sci.*, pages 224–235. Springer, Berlin, 2001.
- [11] Vasco Brattka, Peter Hertling, and Klaus Weihrauch. A tutorial on computable analysis. In *New computational paradigms*, pages 425–491. Springer, New York, 2008.
- [12] Vasco Brattka, Joseph S. Miller, and André Nies. Randomness and differentiability. *Submitted*.
- [13] Rodney G Downey, Evan Griffiths, and Geoffrey LaForte. On Schnorr and computable randomness, martingales, and machines. *Math. Logic Quart.*, 50(6):613–627, 2004.
- [14] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010.

⁶There is at least one exception to this rule. Avigad [1] discovered that the randomness notion, called UD randomness, characterized by a theorem of Weyl is incomparable with Kurtz randomness; and therefore, it is not even preserved by automorphisms.

- [15] Cameron E. Freer and Daniel M. Roy. Computable de Finetti measures. *Annals of Pure and Applied Logic*, 2011.
- [16] Péter Gács. Exact expressions for some randomness tests. *Z. Math. Logik Grundlag. Math.*, 26(5):385–394, 1980.
- [17] Peter Gács. Uniform test of algorithmic randomness over a general space. *Theoret. Comput. Sci.*, 341(1-3):91–137, 2005.
- [18] Peter Gács, Mathieu Hoyrup, and Cristóbal Rojas. Randomness on computable probability spaces—a dynamical point of view. *Theory Comput. Syst.*, 48(3):465–485, 2011.
- [19] Peter Hertling and Yongge Wang. Invariance properties of random sequences. *J.UCS*, 3(11):1241–1249 (electronic), 1997.
- [20] Peter Hertling and Klaus Weihrauch. Random elements in effective topological spaces with measure. *Inform. and Comput.*, 181(1):32–56, 2003.
- [21] John M. Hitchcock and Jack H. Lutz. Why computational complexity requires stricter martingales. *Theory Comput. Syst.*, 39(2):277–296, 2006.
- [22] Mathieu Hoyrup and Cristóbal Rojas. Computability of probability measures and Martin-Löf randomness over metric spaces. *Inform. and Comput.*, 207(7):830–847, 2009.
- [23] Bart Kastermans and Steffen Lempp. Comparing notions of randomness. *Theoret. Comput. Sci.*, 411(3):602–616, 2010.
- [24] Bjørn Kjos-Hanssen and Anil Nerode. Effective dimension of points visited by Brownian motion. *Theoret. Comput. Sci.*, 410(4-5):347–354, 2009.
- [25] L. A. Levin. Uniform tests for randomness. *Dokl. Akad. Nauk SSSR*, 227(1):33–35, 1976.
- [26] Alexander Melnikov and André Nies. K-triviality in computable metric spaces. *Proceedings of the American Mathematical Society*, 141(8):2885–2899, 2013.
- [27] Wolfgang Merkle. The Kolmogorov-Loveland stochastic sequences are not closed under selecting subsequences. *J. Symbolic Logic*, 68(4):1362–1376, 2003.
- [28] Wolfgang Merkle, Nenad Mihailović, and Theodore A. Slaman. Some results on effective randomness. *Theory Comput. Syst.*, 39(5):707–721, 2006.
- [29] Wolfgang Merkle, Joseph S. Miller, André Nies, Jan Reimann, and Frank Stephan. Kolmogorov-Loveland randomness and stochasticity. *Ann. Pure Appl. Logic*, 138(1-3):183–210, 2006.
- [30] Kenshi Miyabe. L^1 -computability, layerwise computability and Solovay reducibility. *Computability*. To appear.
- [31] Kenshi Miyabe. Truth-table Schnorr randomness and truth-table reducible randomness. *MLQ Math. Log. Q.*, 57(3):323–338, 2011.
- [32] André Nies. *Computability and randomness*, volume 51 of *Oxford Logic Guides*. Oxford University Press, Oxford, 2009.
- [33] Noopur Pathak, Cristóbal Rojas, and Stephen G. Simpson. Schnorr randomness and the Lebesgue differentiation theorem. *Proceedings of the American Mathematical Society*, To appear.
- [34] Marian B. Pour-El and J. Ian Richards. *Computability in analysis and physics*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1989.
- [35] Claus-Peter Schnorr. A unified approach to the definition of random sequences. *Math. Systems Theory*, 5:246–258, 1971.
- [36] Matthias Schröder. Admissible representations for probability measures. *MLQ Math. Log. Q.*, 53(4-5):431–445, 2007.
- [37] Javier Gonzalo Silveira. Invariancia por cambio de base de la aleatoriedad computable y la aleatoriedad con recursos acotados. Master’s thesis, University of Buenos Aires, 2011.
- [38] V. V. V’yugin. Ergodic theorems for individual random sequences. *Theoret. Comput. Sci.*, 207(2):343–361, 1998.
- [39] Klaus Weihrauch. *Computable analysis: An introduction*. Texts in Theoretical Computer Science. An EATCS Series. Springer-Verlag, Berlin, 2000.
- [40] David Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991.

ALGORITHMIC RANDOMNESS, MARTINGALES AND DIFFERENTIABILITY

ABSTRACT. In this paper, a number of almost-everywhere convergence theorems are looked at using computable analysis and algorithmic randomness. These include various martingale convergence theorems and almost-everywhere differentiability theorems. General conditions are given for when the rate of convergence is computable and for when convergence takes place on the Schnorr random points. Examples are provided to show that these almost-everywhere convergence theorems characterize Schnorr randomness.

CONTENTS

1. Introduction	30
1.1. Summary of results	31
1.2. A comment on the martingale results	33
1.3. A comment on measurable functions in computable analysis	34
1.4. Outline of the paper	34
1.5. Acknowledgments	35
2. Background	35
2.1. Notation	35
2.2. Computable analysis	35
2.3. Schnorr randomness	36
3. Functions and convergence in measure theory	36
3.1. Integrable functions, measurable functions, and measurable sets	37
3.2. Effective modes of convergence	39
3.3. Convergence on Schnorr randoms	40
3.4. Properties of effectively measurable functions	41
4. Differentiability	43
4.1. The dyadic Lebesgue differentiation theorem	43
4.2. The Lebesgue differentiation theorem	45
4.3. Corollaries to the Lebesgue differentiation theorem	47
5. Martingales in computable analysis	49
5.1. Conditional expectation	49
5.2. L^1 -computable martingales	50
6. The Lévy 0-1 law and uniformly integrable martingales	51
6.1. Some martingale convergence theorems	51
7. More martingale convergence results	53
7.1. Martingale convergence results	53
8. Submartingales and supermartingales	56
9. More differentiability results	57
9.1. Signed measures and Radon-Nikodym derivatives	57
9.2. Functions of bounded variation	60
10. The ergodic theorem	61
11. Backwards martingales and their applications	62
12. Characterizing Schnorr randomness	66
12.1. Monotone convergence, the Lebesgue differentiation theorem, absolutely continuous functions and measures, and uniformly integrable martingales	66
12.2. Singular martingales, functions of bounded variation, and measures	67
12.3. Backwards martingales, the strong law of large numbers, de Finetti's theorem, and the ergodic theorem	69
12.4. Convergence of test functions to 0	70
Appendix A. Proofs from Section 3.	70

A.1. Useful facts	70
A.2. Integrable functions, measurable functions, and measurable sets	71
A.3. Effective modes of convergence	71
A.4. Convergence on Schnorr randomness	73
A.5. Properties of effectively measurable functions	75
References	79

1. INTRODUCTION

The subjects of analysis and probability contain many convergence theorems of the following form.

A.E. Convergence Theorem. *If a sequence of functions $(f_n)_{n \in \mathbb{N}}$ satisfies some property P , then (f_n) converges to some integrable function f almost everywhere as $n \rightarrow \infty$. (Alternatively, $(f_r)_{r>0}$ converges to f as $r \rightarrow 0$.)*

Consider the following closely related examples.

Example 1.1. (Lebesgue differentiation theorem) If $g: [0, 1] \rightarrow \mathbb{R}$ is integrable, then $\frac{1}{2r} \int_{x-r}^{x+r} g(y) dy \rightarrow g(x)$ for almost every x as $r \rightarrow 0$.

Example 1.2. (Lebesgue's theorem) If $f: [0, 1] \rightarrow \mathbb{R}$ is a function of bounded variation function, then f is differentiable almost everywhere. (In this case, $f_r(x) = \frac{g(x+r) - g(x-r)}{2r}$.)

Example 1.3. (Doob's martingale convergence theorem) If (M_n) is a martingale and $\|M_n\|_{L^1} < \infty$, then M_n converges almost everywhere to an integrable function.

Example 1.4. (Ergodic theorem) If g is integrable, and T is a measure preserving transformation, then $\frac{1}{n} \sum_{k < n} g(T^k(x))$ converges almost everywhere. If T is ergodic, then $\frac{1}{n} \sum_{k < n} g(T^k(x)) \rightarrow \int g(x) dx$ for almost every x as $n \rightarrow \infty$.

For all the above theorems, it is natural to ask the following computability questions:

Question 1. *Is the rate of convergence effective (in the parameters of the theorem)?*

It is well known what it means for a sequence of functions to converge effectively in normed spaces like L^1 and L^2 . A similar characterization can be given for almost everywhere convergence: a sequence of functions (f_n) converges to f with a effective rate of almost everywhere convergence if given $\varepsilon > 0$ and $\delta > 0$, we can compute some $m \in \mathbb{N}$ such that $|f_n(x) - f(x)| < \varepsilon$ for all $n \geq m$ and all x except on a set of size less than δ .

Some a.e. convergence theorems have effective rates of convergence. For example, Avigad, Gerhardy, and Towsner [2] showed that the rate of almost everywhere convergence in the ergodic theorem is computable from T and g when T is ergodic. I will show a similar result for the Lebesgue differentiation theorem.

However, not all the theorems have computable rates of convergence. This is the case for Lebesgue's theorem, Doob's martingale convergence theorem, and the ergodic theorem (in the nonergodic setting). However, when certain additional conditions are assumed, one can then compute a rate of convergence.

Question 2. *If the rate of convergence is not effective, what are additional conditions that guarantee an effective rate of convergence.*

For example, Avigad et al. [2] showed that the rate of convergence in the ergodic theorem is computable from g , T and the limit g^* . (Note, it is not trivial to compute the rate of convergence from the limit of a series. For example, it is easy to construct a computable sequence of constant functions which converge to 0, but do not do so effectively.) In the L^2 -case, Avigad et al. [2] showed the rate of convergence is computable from g , T , and the L^2 -norm of g .

In this paper I will give similar results for Lebesgue's theorem, Doob's martingale convergence theorem, and others. All the results follow the pattern in this observation.¹

¹It is important to note that Observation 1 is not itself a theorem or metatheorem. Indeed, there are (contrived) cases where it fails to hold—let f_n be some computable sequence of constant functions converging to zero with a noncomputable rate of convergence.

Observation 1. *For most a.e. convergence theorems, a rate of almost everywhere convergence is computable from the sequence (f_n) , the limit f , and the bounds $\inf_n \|f_n\|_{L^1}$, $\sup_n \|f_n\|_{L^1}$.*

In many cases, such as in the ergodic theorem, $\inf_n \|f_n\|_{L^1}$ and $\sup_n \|f_n\|_{L^1}$ are computable from the sequence (f_n) and the limit f , and therefore they are not explicitly needed. In other cases, such as the Lebesgue differentiation theorem, all three extra conditions are naturally computable from the parameters of the theorem (which is why the rate of convergence in the Lebesgue differentiation theorem is computable without additional assumptions). Further, if we work in L^2 instead of L^1 , we do not need the limit f , just its L^2 -norm $\|f\|_{L^2}$.

Question 3. *At which points does the sequence converge (under various computability conditions)?*

For example, if we consider the Lebesgue differentiation theorem, we can ask at which x does $\frac{1}{2r} \int_{x-r}^{x+r} f(y) dy$ converge for all f computable in the L^1 -norm. Notice the set of such x is measure one, since there are only countably-many f computable in the L^1 -norm.

This question was first asked by Pathak [39] using the tools of algorithmic randomness. Algorithmic randomness classifies measure-one sets of points that behave randomly with respect to “computable tests”. Pathak showed that convergence happens on all Martin-Löf random x . She left it as an open question whether this could be strengthened to a larger class of points. In this paper, I will show that it can be strengthened to Schnorr randomness, and that this is the best possible. In other words, the Lebesgue differentiation theorem characterizes Schnorr randomness. This same result was independently and concurrently discovered by Pathak, Rojas, and Simpson [40].

Similar investigations have been made into randomness and the ergodic theorem [50, 37, 25, 21, 16, 3, 17], randomness and Lebesgue’s theorem [10, 8, 18], and randomness and martingale convergence [47]. In this paper, I expand on these results, specifically looking at Schnorr randomness. Indeed, I ask this converse to Question 3.

Question 4. *Which conditions guarantee convergence on Schnorr randoms?*

It turns out the answers to Questions 1 and 2 provide an answer, when using this informal observation—which will be made formal in Lemma 3.19.

Observation 2. *Effective a.e. convergence implies convergence on Schnorr randoms.*

This will allow us to “kill two birds with one stone”, by focusing on questions in computable analysis (Questions 1 and 2), we can answer questions in algorithmic randomness (Questions 3 and 4) for free. However, to show that one cannot strengthen Schnorr randomness to a larger class of random points, we will need an example for each theorem showing that if x is not Schnorr random, then there are computable parameters for which convergence does not happen on x . I provide a number of such examples.

1.1. Summary of results. The results of this paper are diverse and the paper is organized by the tools and lemmas needed to prove the theorems. Table 1 is a summary of all the known convergence theorems which characterize Schnorr randomness. The first column is a short description of the convergence theorem. The second column is a reference to the result showing that the sequence in question converges on all Schnorr randoms. The third column is a reference to the result showing that if a point is not Schnorr random, then there exists such a sequence which fails to converge on that point. If a cell is blank, that direction is subsumed by a stronger result in another row. A “?” means this direction is still an open question (and so that row may not really be a characterization of Schnorr randomness). Some of the results are due to others, or were independently discovered. I provide footnotes in these cases.

Table 1: Characterizations of Schnorr randomness by a.e. convergence theorems. (See Section 1.1 for an explanation of this table.)

Convergence of martingales: $M_n \rightarrow M_\infty$		
(M_n) is L^1 -comp.; M_∞ is L^1 -comp.; $\sup_n \ M_n\ _{L^1}$ is computable	Thm. 7.11	
(M_n) is uniformly integrable, L^1 -comp.; M_∞ is L^1 -computable	Lem. 6.5	
(M_n) is nonnegative, singular ($M_\infty = 0$), L^1 -computable	Lem. 7.4	
(M_n) is nonnegative, singular ($M_\infty = 0$), computable		Thm. 12.9

(M_n) is L^2 -computable; $\sup_n \ M_n\ _{L^2} = \ M_\infty\ _{L^2}$ is computable	Cor. 6.8	
(M_n) is nonneg., unif. int., dyadic, computable; $\ M_\infty\ _{L^2}$ is computable		Thm. 12.7

Convergence of super/submartingales: $M_n \rightarrow M_\infty$

(M_n) is L^1 -comp., super/submart.; $\lim_n \ M_n\ _{L^1}$ is comp.; M_∞ is L^1 -comp.	? ²	
(M_n) is nonnegative, L^1 -comp., supermart.; M_∞ is L^1 -comp.	Thm. 8.1	
(M_n) is nonnegative, L^1 -comp., supermart.; $M_\infty = 0$	Lem. 7.4	Thm. 12.9
(M_n) is nonneg., L^1 -comp., submart.; M_∞ is L^1 -comp.; $\sup_n \ M_n\ _{L^1}$ is comp.	Thm. 8.5	Thm. 12.6

Convergence of reverse martingales: $M_{-n} \rightarrow M_{-\infty}$

(M_{-n}) is L^1 -computable; $M_{-\infty}$ is L^1 -computable	Thm. 11.2	
(M_{-n}) is L^2 -computable; $\ M_{-\infty}\ _{L^2}$ is L^2 -computable	Thm. 11.2	
(M_{-n}) is bounded, a.e. computable; M_∞ is computable constant		Cor. 12.17

Lebesgue differentiation theorem: $\int_{B(x,r)} f(y) dy / \lambda(B(x,r)) \rightarrow f(x)$

f is L^1 -computable	Thm. 4.10, [40] ³	Thm. 12.3
f is L^2 -computable		
f is bounded, L^1 -computable		[40, 18]

Lebesgue density theorem: $\lambda(A \cap B(x,r)) / \lambda(B(x,r)) \rightarrow \mathbf{1}_A(x)$

A is effectively measurable	Cor. 4.15	[40, 18] ⁴
A is effectively closed; $\lambda(A)$ is computable	Cor. 4.15	?

Differentiability of functions f (with derivative Df and total variation $V(f)$)

f is comp. on dense set, bounded var.; Df is L^1 -comp.; $V(f)$ is comp.	Thm. 9.19	
f is comp., bounded variation; Df is L^1 -comp.; $V(f)$ is comp.	Cor. 9.20	
f is absolutely continuous; Df is L^1 -comp.	Cor. 4.17, [18] ⁵	
f is computable, absolutely continuous; $\ Df\ _{L^2}$ is computable	Cor. 6.9	
f is increasing, effectively absolutely continuous, comp.; $\ Df\ _{L^2}$ is comp.		Cor. 12.5 ⁶
f is increasing, Lipschitz, effectively absolutely continuous		[18]
f is computable, increasing, singular ($Df = 0$ a.e.)		Cor. 12.15
f is comp. on dense set, increasing, singular, only contains jumps		Cor. 12.12

$\nu(B(x,r)) / \lambda(B(x,r)) \rightarrow \frac{d\nu}{d\lambda}(x)$ for signed measures ν

ν is computable; $\frac{d\nu}{d\lambda}$ is L^1 -comp.; $\ \nu\ _{TV}$ is comp.	Thm. 9.12	
ν is absolutely continuous, computable, positive; $\frac{d\nu}{d\lambda}$ is L^1 -comp.	Cor. 4.20	Cor. 12.4
ν is continuous, singular ($\frac{d\nu}{d\lambda} = 0$ a.e.), computable, positive		Cor. 12.14
ν is atomic, singular ($\frac{d\nu}{d\lambda} = 0$ a.e.), computable, positive		Cor. 12.11

Ergodic theorem: $\frac{1}{n} \sum_{k < n} f \circ T^k \rightarrow f^*$

f is L^1 -comp.; T is effectively measurable; f^* is L^1 -comp.	Thm. 10.2 ⁷	
f is L^2 -comp.; T is effectively measurable; $\ f^*\ _{L^2}$ is comp.	Thm. 10.2	
f is a.e. comp.; T is a.e. comp., ergodic		[21]

Monotone convergence thm: Convergence of (f_n) increasing

(f_n) is L^1 -comp.; $\ f_n\ _{L^1}$ is computable	Prop. 8.2	
(f_n) is L^2 -comp.; $\ f_n\ _{L^2}$ is computable	Prop. 8.2	

²See Problem 8.6.

³This was independently discovered by this author and by Pathak, Rojas, and Simpson [40].

⁴While neither paper makes this explicit, the example function f they each give for the Lebesgue differentiation theorem is 0,1 valued and therefore the indicator function of some effectively measurable set A .

⁵This result is a direct corollary of the effective Lebesgue differentiation theorem (Thm. 4.10, [40]) that was noticed by this author and Freer, Kjos-Hanssen, Nies, and Stephan [18].

⁶This also follows from the Lipschitz result of Freer, Kjos-Hanssen, Nies, and Stephan [18] in the next line.

⁷Theorem 10.2 is a summary of the results from [2, 21, 22, 40] with a few gaps filled in.

(f_n) is computable; $\|f_n\|_{L^2}$ is computable Thm. 12.2⁸

$\varphi_n \rightarrow f$ for test functions, dyadic averages, trigonometric polynomials

φ_n is fast Cauchy sequence of test functions in L^1 (or in measure)	Prop. 3.18, [40] ⁹	
φ_n is fast Cauchy sequence of test functions converging to 0 in L^2		Thm. 12.18
$f^{(n)}$ are dyadic averages; f is L^1 -computable	Prop. 4.6	Thm. 12.6
$\sigma_n(f)$ are trig. polynomials (from Fejér kernel); f is $L^1(\mathbb{T} \rightarrow \mathbb{C})$ -comp.	Cor. 4.22	?
SLLN, de Finetti's thm: Convergence of $\frac{1}{n} \sum_{k < n} X_k$ for integrable random variables (X_n)		
(X_n) is L^1 -comp., exchangeable (de Finetti's theorem)	Cor. 11.10	
(X_n) is L^1 -computable, i.i.d. (strong law of large numbers)	Cor. 11.6	Cor. 12.17

1.2. A comment on the martingale results. A significant portion of this paper concerns martingales. Informally, martingales are formalizations of gambling strategies—a martingale (M_n) is a sequence of random variables representing the capital of a gambling strategy at time n . They are widely used in probability theory and analysis, as well as in algorithmic randomness. In algorithmic randomness, martingales can be used to characterize a number of randomness notions, including Schnorr randomness, computable randomness, and Martin-Löf randomness (see [12, 38]). However, there is a difference between how martingales are treated in algorithmic randomness and how they are used in probability theory and analysis.

For one, in algorithmic randomness, the martingales are (usually) gambling strategies on coin flips. Such martingales, which I will call dyadic martingales, are a specific instance of the more general martingales used in probability theory. By considering, this larger class of martingales, we can ask new questions that could not be asked of dyadic martingales.

Another difference is that algorithmic randomness is more concerned with success than convergence. We say that a martingale (M_n) succeeds on x if $\limsup_n M_n(x) = \infty$, that is the strategy M_n wins arbitrarily large amounts of money on x . Of the three most common randomness notions—Schnorr randomness, computable randomness, and Martin-Löf randomness—only computable randomness has a well-known characterization in terms of martingale convergence instead of martingale success.¹⁰

For example, consider the following three characterizations of Schnorr randomness. The characterization in (2) is the classical martingale characterization of Schnorr randomness and (3) and (4) are new characterizations which follow from results in this paper (Theorem 7.11, Corollary 6.8, and Theorem 12.6). (Note, in (3) and (4) we could replace convergence with success and the characterization would still hold.)

Example 1.5. Recall, a computable dyadic martingale is a computable function $M: 2^{<\omega} \rightarrow \mathbb{R}$ such that $\frac{1}{2}M(\sigma 0) + \frac{1}{2}M(\sigma 1) = M(\sigma)$. Use the notation $M_n(x) = M(x \upharpoonright n)$. The following are equivalent.

- (1) $x \in 2^{\mathbb{N}}$ is Schnorr random (on the fair-coin measure).
- (2) (Classical) For all nonnegative computable dyadic martingales (M_n) and all computable unbounded functions $h: \mathbb{N} \rightarrow \mathbb{N}$, we have that $M_n(x) \leq h(n)$ for all but finitely-many n .
- (3) (New) For all nonnegative, computable dyadic martingales (M_n) such that $\lim_n M_n$ is L^1 -computable, we have that $M_n(x)$ converges.
- (4) (New) For all nonnegative, computable dyadic martingales (M_n) such that $\sup_n \|M_n\|_{L^2}$ is computable, we have that $M_n(x)$ converges.

However, the results in this paper go far beyond giving a new dyadic martingale characterization of Schnorr randomness. Not only does algorithmic randomness provide us with new tools to study computable analysis; computable analysis also provides us new tools to study algorithmic randomness. Martingales are one such

⁸This is closely related to the Schnorr integral tests of Miyabe [34].

⁹This is also a theorem of Pathak, Rojas, and Simpson [40] where the test functions are rational polynomials and convergence is in L^1 .

¹⁰There are not-widely-known published results, which when combined, give a martingale-convergence characterization of Martin-Löf randomness. Takahashi [47] showed that Doob's upcrossing inequality implies that computable martingales (that is, martingales in the more general sense of probability theory where (M_n) is a computable sequence of computable functions) converge on Martin-Löf randoms. Edward Dean [personal communication], independently, showed that layerwise-computable martingales converge on Martin-Löf randoms. Merkle, Mihailović, and Slaman [33] gave an example of a computable martingale (in the more general sense of probability theory) which only converges on Martin-Löf randoms.

tool. This paper makes significant use of martingales to prove results. One particular type of martingale not previously used in algorithmic randomness is the backwards martingale. To demonstrate their usefulness in algorithmic randomness, I use backwards martingales to prove a new variation of Kučera's theorem in Corollary 11.4: for every Schnorr random $x \in 2^{\mathbb{N}}$ and for every closed set C of positive computable measure, C contains some y which equals x , except that finitely many bits are permuted.

1.3. A comment on measurable functions in computable analysis. There is an inherent challenge when working with measurable functions in a computable setting. Measurable functions are not continuous and therefore it is difficult to describe them as maps in a computable manner. Moreover, a single function is best thought of as an equivalence class (under a.e. equivalence). It is challenging to talk about the value $f(x)$ when f is an equivalence class (an important issue when asking about which points an a.e. convergence theorem holds!).

Some authors have taken the easy approach and restricted their attention to computable functions or a.e. computable functions. However, in this paper, I will try to express the theorems in full generality. In order to do this, I will need a clear theory of effectively measurable functions. The space of measurable functions (modulo a.e. equivalence) is naturally described as a computable metric space under a suitable metric which characterizes convergence in measure. The class of effectively measurable functions includes the real-valued functions computable in the L^1 -norm as well as other functions (which may not even be integrable or real-valued).

In order to talk about the valuation of functions, each effectively measurable function f will have a representative \tilde{f} . This representative is well-defined (and well-behaved) on Schnorr random points. This representative approach is adapted from Pathak [39] (and is also used in Pathak, Rojas, and Simpson [40]). The same ideas are implicit in the reverse mathematics of the dominated convergence theorem [58, 1].

Other computable approaches to measurable sets and functions include [43, 4, 31, 20, 56, 55, 15, 24, 34]. These approaches are essentially the same as either the metric space approach or the representative approach used in this paper. The biggest difference is that some representative approaches—e.g. layerwise computability [24]—only define f on the Martin-Löf random points. However, it is possible to uniquely extend each such f to the Schnorr random points.

My hope is that Section 3 (on effectively measurable functions) not only serves the needs of this paper, but is of use to other researchers in the field.

1.4. Outline of the paper. In Section 2, I give background on computable analysis and Schnorr randomness.

In Section 3, I present a theory of measurable functions, integrable functions, and measurable sets. This also includes the important Lemma 3.19 that effective a.e. convergence implies convergence on Schnorr randoms. Most of the proofs have been moved to Appendix A.

In Section 4, I prove an effective version of the Lebesgue differentiation theorem and discuss many of its corollaries. The proof relies on Kolmogorov's inequality for dyadic martingales.

In Section 5, I give a computable presentation of martingale theory, which will be needed for most of the rest of this paper.

In Section 6, I prove an effective version of the Lévy 0-1 law, which is a simpler version of Doob's martingale convergence theorem and an analog to the Lebesgue differentiation theorem.

In Section 7, I prove an effective version of the martingale convergence theorem. I also give another version for square integrable martingales.

In Section 8, I prove an effective version of the submartingale and supermartingale convergence theorems. I also give another version for square integrable martingales.

In Section 9, I return to differentiability, using effective martingale convergence to prove more differentiability results that extend the Lebesgue differentiation theorem and Lebesgue's theorem.

In Section 10, I survey some results in ergodic theory, filling in gaps in the published literature.

In Section 11, I discuss backwards martingales and some of their applications, including a variation of Kučera's theorem, the strong law of large numbers, and de Finetti's theorem. I also, compare them with ergodic averages.

I intend to follow up this paper with a sequel, exploring martingale convergence and differentiability when the limit is not computable. Such cases characterize computable randomness, Martin-Löf randomness, and weak-2 randomness.

1.5. Acknowledgments. I would like to thank André Nies, Jeremy Avigad, and Bjørn Kjos-Hanssen for many helpful corrections on earlier drafts of this paper. I would also like to thank Laurent Bienvenu, Johanna Franklin, Mathieu Hoyrup, Kenshi Miyabe, Noopur Pathak, Cristóbal Rojas, and Stephan Simpson for helpful discussions on parts of this work.

2. BACKGROUND

In this section I give the necessary background in computable analysis, effective measure theory, effective probability theory, and Schnorr randomness.

2.1. Notation. Let $2^{<\omega}$ be the space of finite binary strings, $2^{\mathbb{N}}$ be the space of infinite binary strings, \emptyset_{string} be the empty string, $\sigma \prec \tau$ and $\sigma \prec x$ mean σ is a proper initial segment of $\tau \in 2^{<\omega}$ or $x \in 2^{\mathbb{N}}$, $[\sigma] = \{x \in 2^{\mathbb{N}} \mid \sigma \prec x\}$. Also for $\sigma \in 2^{<\omega}$ (or $x \in 2^{\mathbb{N}}$), let $\sigma(n)$ be the n th digit of σ (where $\sigma(0)$ is the “0th” digit) and $\sigma \upharpoonright n = \sigma(0) \cdots \sigma(n-1)$. A set of strings $\{\sigma_0, \sigma_1, \dots\}$ is prefix free if the no string in the set is a prefix of another (equivalently, the collection $\{[\sigma_0], [\sigma_1], \dots\}$ is pair-wise disjoint).

2.2. Computable analysis. Here I present some basics of computable analysis. For additional information on the basics see Pour El and Richards [41], Weihrauch [52], or Brattka et al. [7]. I assume the reader has some familiarity with basic computability theory on \mathbb{N} , $2^{\mathbb{N}}$, and $\mathbb{N}^{\mathbb{N}}$ as in [46]. It would also help to have some familiarity with the theory of computation on the reals.

Definition 2.1. Fix an enumeration of the rationals $\mathbb{Q} = \{q_i\}_{i \in \mathbb{N}}$ (such that addition and multiplication are computable). A real $x \in \mathbb{R}$ is COMPUTABLE if there is a computable function $h: \mathbb{N} \rightarrow \mathbb{N}$ such that for all $m > n$, we have $|q_{h(m)} - q_{h(n)}| \leq 2^{-n}$ and $x = \lim_{n \rightarrow \infty} q_{h(n)}$.

This can be generalized to an arbitrary complete metric space.

Definition 2.2. A COMPUTABLE (POLISH) METRIC SPACE is a triple $\mathbb{X} = (X, d, S)$ such that

- (1) X is a complete metric space with metric $d: X \times X \rightarrow [0, \infty)$.
- (2) $S = \{a_i\}_{i \in \mathbb{N}}$ is a countable dense subset of X (the SIMPLE POINTS of \mathbb{X}).
- (3) The distance $d(a_i, a_j)$ is computable uniformly from i and j .

A point $x \in X$ is said to be COMPUTABLE if there is a computable function $h: \mathbb{N} \rightarrow \mathbb{N}$ such that for all $m > n$, we have $d(a_{h(m)}, a_{h(n)}) \leq 2^{-n}$ and $x = \lim_{n \rightarrow \infty} a_{h(n)}$. The sequence $(a_{h(n)})$ is the CAUCHY-NAME for x .

Example 2.3. For the differentiability results, I will be using two spaces. The first is the unit cube $[0, 1]^d$ with the usual Euclidean distance. The second is the unit torus $\mathbb{T}^d := (\mathbb{R}/\mathbb{Z})^d$, which will be identified as the half open unit cube $[0, 1)^d$ with the Euclidean metric that wraps around each edge, i.e. given $x = (x_1, \dots, x_d), y = (y_1, \dots, y_d) \in [0, 1)^d$,

$$d(x, y) = \left(\sum_{i=1}^d \left(\min \{|x_i - y_i|, 1 - |x_i - y_i|\} \right)^2 \right)^{1/2}.$$

The simple points of \mathbb{T}^d and $[0, 1]^d$ are taken to be vectors with rational components. A little thought reveals that a vector $x \in [0, 1]^d$ (or $x \in \mathbb{T}^d$) is computable if and only if each coordinate is a computable real.

On a computable metric space $\mathbb{X} = (X, d, S)$, the BASIC OPEN BALLS are sets of the form $B(a, r) = \{x \in X \mid d(x, a) < r\}$ where $a \in S$ and $r > 0$ is rational. The Σ_1^0 sets (EFFECTIVELY OPEN SETS) are computable unions of basic open balls; Π_1^0 sets (EFFECTIVELY CLOSED SETS) are the complements of Σ_1^0 sets. A function $f: \mathbb{X} \rightarrow \mathbb{R}$ is COMPUTABLE (or EFFECTIVELY CONTINUOUS) if for each Σ_1^0 set U in \mathbb{R} , the set $f^{-1}(U)$ is Σ_1^0 in \mathbb{X} (uniformly in U), or equivalently, there is an algorithm which sends every Cauchy-name of x to a Cauchy-name of $f(x)$ (see [52]). A function $f: \mathbb{X} \rightarrow [0, \infty]$ is LOWER SEMICOMPUTABLE if it is the supremum of a computable sequence of computable functions $f_n: \mathbb{X} \rightarrow [0, \infty)$.

A real x is said to be LOWER (UPPER) SEMICOMPUTABLE if it is the supremum (resp. infimum) of a computable sequence of rationals.

Definition 2.4. If $\mathbb{X} = (X, d, S)$ is a computable metric space, then a Borel measure μ is a COMPUTABLE MEASURE on \mathbb{X} if $\int g d\mu$ is computable uniformly from g for all computable $g: \mathbb{X} \rightarrow [0, 1]$. A COMPUTABLE PROBABILITY SPACE is a pair (\mathbb{X}, μ) where \mathbb{X} is a computable metric space, μ is a computable measure on \mathbb{X} , and $\mu(X) = 1$.

There are a number of other equivalent definitions of computable measure, including the following characterization.

Proposition 2.5 ([45, 26]). *A measure μ on a computable metric space $\mathbb{X} = (X, d, S)$ is computable if and only if the value $\mu(X)$ is computable, and for each effectively open set $U \subseteq \mathbb{X}$, the measure $\mu(U)$ is lower semicomputable uniformly from U .*

Moreover, the computable probability measures on \mathbb{X} are exactly the computable points in the space of probability measures under the Prokhorov metric.

I will often blur the distinction between a metric space—or a probability space—with its set of points, e.g. writing $x \in \mathbb{X}$ or $x \in (\mathbb{X}, \mu)$ to mean that $x \in X$ where $\mathbb{X} = (X, d, S)$.

Example 2.6. The manifolds $[0, 1]^d$ and \mathbb{T}^d can be endowed with the LEBESGUE MEASURE λ . (The Lebesgue measure on \mathbb{T}^d is just the Lebesgue measure on $[0, 1]^d$.) Both are computable probability measures, and further (\mathbb{T}^d, λ) is translation-invariant. Similarly, on $2^{\mathbb{N}}$ let λ be the FAIR-COIN MEASURE, i.e. the measure such that $\lambda([\sigma]) = 2^{-|\sigma|}$.

Definition 2.7. Let $\mathbb{X} = (X, S, d)$ be a computable metric space.

- (1) $(\mathbb{X}, +, \cdot)$ is a COMPUTABLE (TOPOLOGICAL) VECTOR SPACE if \mathbb{X} is a vector space and with computable vector addition $+$ and scalar multiplication \cdot operations.
- (2) $(\mathbb{X}, \|\cdot\|, +, \cdot)$ is a COMPUTABLE BANACH SPACE if $(\mathbb{X}, +, \cdot)$ is a computable vector space and the metric d comes from a computable norm $\|\cdot\|$.
- (3) $(\mathbb{X}, \|\cdot\|, \langle \cdot, \cdot \rangle, +, \cdot)$ is a COMPUTABLE HILBERT SPACE if $(\mathbb{X}, \|\cdot\|, +, \cdot)$ is a computable Banach space with computable inner product $\langle \cdot, \cdot \rangle$.
- (4) $(\mathbb{X}, \wedge, \vee)$ is a COMPUTABLE (TOPOLOGICAL) LATTICE if \mathbb{X} is a lattice with computable meet \wedge and join \vee operations.
- (5) $(\mathbb{X}, \wedge, \vee, \neg, \perp, \top)$ is a COMPUTABLE (TOPOLOGICAL) BOOLEAN ALGEBRA if \mathbb{X} is a Boolean algebra with computable meet \wedge , join \vee , and complement \neg operations and computable bottom \perp and top \top elements.

Remark 2.8. There are a number of natural Banach spaces that are not computable, for example the space of signed Borel measures on $[0, 1]$. This is because they have no countable dense subset. However, we may still represent these spaces using a weaker topology as will be done in Section 9.1.

2.3. Schnorr randomness.

Definition 2.9. Let (\mathbb{X}, μ) be a computable probability space. A SCHNORR TEST (U_n) is a computable sequence of effectively open sets U_n such that $\mu(U_n) \leq 2^{-n}$ for all n and $\mu(U_n)$ is uniformly computable in n . For any $x \in \mathbb{X}$, say x is COVERED by (U_n) if $x \in \bigcap_n U_n$. Say $x \in \mathbb{X}$ is SCHNORR RANDOM if x is not covered by any Schnorr test.

Remark 2.10. We may assume a Schnorr test (U_n) is decreasing by taking an intersection. Similarly, we may also replace 2^{-n} by any computable sequence that decreases to 0 by taking a subsequence (see [12, 38]).

Example 2.11. Let $y_1, \dots, y_d \in [0, 1]$ (resp. \mathbb{T}). For each $1 \leq i \leq d$, let x_i be some binary expansion of y_i . It is easy to see that (y_1, \dots, y_d) is Schnorr random on $([0, 1]^d, \lambda)$ (resp. on (\mathbb{T}^d, λ)) if and only if $x_1 \oplus \dots \oplus x_n \in 2^{\mathbb{N}}$ is Schnorr random on $(2^{\mathbb{N}}, \lambda)$. (Recall, $x_1 \oplus x_2$ is the join operation on $2^{\mathbb{N}}$ defined by $(x_1 \oplus x_2)(2n) := x_1(n)$ and $(x_1 \oplus x_2)(2n+1) := x_2(n)$.)

3. FUNCTIONS AND CONVERGENCE IN MEASURE THEORY

This section provides background on measurable functions and convergence. It is quite important to the results in this paper. (For example, the frequently used Lemma 3.19 is the only fact the reader will need to know about Schnorr randomness in Sections 4 through 11.)

As mentioned in the introduction, there is a need for two approaches to working with measurable functions (and sets).¹¹

¹¹A third approach may come to mind: use Borel measurable functions and sets, ignoring a.e. equivalence. The difficulty with this approach is that even effectively open sets may not have a computable measure. The situation becomes more complex as one moves up the Borel hierarchy.

- (1) Use *equivalence classes* of almost-everywhere equivalent objects.
- (2) Use *specific functions and sets* that are defined and unique up to some *specific* measure-one set (which will turn out to be the set of Schnorr random points).

Table 2 compares the two approaches (in the setting of L^1 -computable functions).

Equivalence classes	Specific functions
f an L^1 -limit of fast Cauchy sequences	\tilde{f} a pointwise limit of fast Cauchy sequences
f unique a.e.	\tilde{f} unique on Schnorr randoms
f computable in the L^1 -norm	\tilde{f} “computable” on Schnorr randoms

TABLE 2. The two approaches to the computability of L^1 functions.

Besides giving definitions and basic facts, the main result of this section is Lemma 3.19, that a.e. convergence implies convergence on Schnorr randoms. (This fact has been hinted at in some of the work on convergence for Schnorr randoms, including Pathak, Rojas, and Simpson [40]. It was also known to Hoyrup and Rojas [personal communication] independently of this author.)

3.1. Integrable functions, measurable functions, and measurable sets. Let us start with real-valued functions on the space $(2^{\mathbb{N}}, \lambda)$.

Proposition 3.1. *On $(2^{\mathbb{N}}, \lambda)$ the following hold.*

- (1) (Functions) Consider the following spaces (of a.e. equivalence classes $[f]_{\sim}$) of Borel measurable functions. Let the TEST FUNCTIONS \mathcal{T} be those of the form

$$(3.1) \quad \varphi = \sum_{i=0}^{k-1} c_i \mathbf{1}_{[\sigma_i]} \quad (\sigma_0, \dots, \sigma_{k-1} \in 2^{<\omega}; c_0, \dots, c_{k-1} \in \mathbb{Q}).$$

Also consider the lattice given by

$$f \wedge g = \min(f, g) \quad \text{and} \quad f \vee g = \max(f, g).$$

- (a) The MEASURABLE FUNCTIONS $L^0(2^{\mathbb{N}}, \lambda)$ with the metric¹²

$$d_{meas}(f, g) = \int \min\{|f - g|, 1\} d\lambda$$

form a computable metric space, a computable vector space, and a computable lattice

$$(L^0(2^{\mathbb{N}}, \lambda), \mathcal{T}, d_{meas}, +, \cdot, \min, \max).$$

- (b) The INTEGRABLE FUNCTIONS $L^1(2^{\mathbb{N}}, \lambda)$ with norm

$$\|f\|_{L^1} = \int |f| d\lambda$$

form a computable Banach space and a computable lattice

$$(L^1(2^{\mathbb{N}}, \lambda), \mathcal{T}, \|\cdot\|_{L^1}, +, \cdot, \min, \max).$$

- (c) The SQUARE INTEGRABLE FUNCTIONS $L^2(2^{\mathbb{N}}, \lambda)$ with inner product and norm

$$\langle f, g \rangle = \int f, g d\lambda, \quad \|f\|_{L^2} = \left(\int |f|^2 d\lambda \right)^{1/2}$$

form a computable Hilbert space and a computable lattice

$$(L^2(2^{\mathbb{N}}, \lambda), \mathcal{T}, \|\cdot\|_{L^2}, \langle \cdot, \cdot \rangle_{L^2}, +, \cdot, \min, \max).$$

¹²As we shall see, this metric characterizes convergence in measure. It is equivalent to the Ky-Fan metric $d_{KF}(f, g) := \inf \{ \varepsilon > 0 \mid \mu(\{x \mid |f - g| \geq \varepsilon\}) \leq \varepsilon \}$. (Indeed, $(L^0(2^{\mathbb{N}}, \lambda), \mathcal{T}, d_{KF}, +, \cdot, \min, \max)$ is also a computable metric space, a computable vector space, and a computable lattice with the same computable points as $(L^0(2^{\mathbb{N}}, \lambda), \mathcal{T}, d_{meas})$.)

- (2) (Set spaces) Consider the following space (of a.e. equivalence classes $[A]_{\sim}$) of Borel measurable sets. Let the TEST SETS \mathcal{T} be those of the form

$$C = \bigcup_{i=0}^{k-1} [\sigma_i] \quad (\text{prefix-free } \sigma_0, \dots, \sigma_{k-1} \in 2^{<\omega}).$$

- (d) The MEASURABLE SETS $\mathcal{B}(2^{\mathbb{N}}, \lambda)$ with metric

$$d(A, B) = \lambda(A \triangle B)$$

form a computable metric space and a computable Boolean algebra

$$(\mathcal{B}(2^{\mathbb{N}}, \lambda), \mathcal{T}, d, \cup, \cap, \cdot^c, \emptyset, 2^{\mathbb{N}}).$$

Proof. straightforward. \square

Definition 3.2. The computable points of each of the above spaces are, respectively, called the EFFECTIVELY MEASURABLE FUNCTIONS (L_{comp}^0), the L^1 -COMPUTABLE FUNCTIONS (L_{comp}^1), the L^2 -COMPUTABLE FUNCTIONS (L_{comp}^2), and the EFFECTIVELY MEASURABLE SETS.

We may also consider measurable functions taking values in other computable metric spaces $\mathbb{Y} = (Y, S, d_{\mathbb{Y}})$.

Proposition 3.3. Let $\mathbb{Y} = (Y, S, d_{\mathbb{Y}})$ be a computable metric space. The space of measurable functions from $(2^{\mathbb{N}}, \lambda)$ to $\mathbb{Y} = (Y, S, d_{\mathbb{Y}})$ is a computable metric space under the metric

$$d_{meas}(f, g) = \int \min\{d_{\mathbb{Y}}, 1\} d\lambda$$

and TEST FUNCTIONS of the form

$$\varphi(x) = c_i \mathbf{1}_{[\sigma_i]} \quad \text{when } x \in [\sigma_i] \quad (\text{prefix-free } \sigma_0, \dots, \sigma_{k-1} \in 2^{<\omega}; c_0, \dots, c_{k-1} \in S).$$

The computable points in this space are called EFFECTIVELY MEASURABLE FUNCTIONS.

Proof. straightforward. \square

Remark 3.4. The space of measurable sets and the space of 0, 1-valued measurable functions (Proposition 3.3 with $\mathbb{Y} = \{0, 1\}$) are the same space. (More specifically, the map $A \mapsto \mathbf{1}_A$ is a bijective isometry where test sets are mapped to test functions.)

The above definitions extend to any computable probability space (\mathbb{X}, μ) . The only thing that changes is the choice of test functions. This requires a technical lemma.

Lemma 3.5 (Bosserhoff [6], Hoyrup and Rojas [26]). *For any computable metric space $\mathbb{X} = (X, \mathcal{S}, d)$ with computable probability measure μ , there is a computable sequence of pairs $\{(a_i, r_i)\}_{i \in \mathbb{N}}$ ($a_i \in \mathcal{S}$, $r_i \in \mathbb{R}$) representing a family of balls $\text{Basis}(\mathbb{X}, \mu) = \{B(a_i, r_i)\}_{i \in \mathbb{N}}$.*

- (1) Each $B(a_i, r_i)$ has a μ -null boundary. (Hence $\mu(B(a_i, r_i))$ is computable uniformly from i .)
- (2) $\text{Basis}(\mathbb{X}, \mu)$ is an effective basis of \mathbb{X} , i.e. for every effectively open set U , there is a computable sequence (i_k) of indices computable uniformly from (each name for) U such that $U = \bigcup_{k=0}^{\infty} B(a_{i_k}, r_{i_k})$.

Since the choice of basis is not unique, let $\text{Basis}(\mathbb{X}, \mu)$ denote a fixed choice of basis for each space (\mathbb{X}, μ) .

Definition 3.6. Say that $C \subseteq \mathbb{X}$ is a CELL of $\text{Basis}(\mathbb{X}, \mu)$ if $C = A_1 \cap \dots \cap A_{\ell} \cap B_1^c \cap \dots \cap B_k^c$ for $A_1, \dots, A_{\ell}, B_1, \dots, B_k \in \text{Basis}(\mathbb{X}, \mu)$. (Notice, using the enumeration of $\text{Basis}(\mathbb{X}, \mu)$ that each cell is coded by some $\sigma \in 2^{<\omega}$.)

Proposition 3.7. The measure of each cell of $\text{Basis}(\mathbb{X}, \mu)$ is computable from its code σ .

Proof. See Appendix A.2. \square

Definition 3.8. On (\mathbb{X}, μ) , the spaces of real-valued functions $L^0(\mathbb{X}, \mu)$, $L^1(\mathbb{X}, \mu)$, $L^2(\mathbb{X}, \mu)$ as well as the space of measurable sets and the space of \mathbb{Y} -valued measurable functions are defined as before, except that cylinder sets $[\sigma_i]$ are replaced with cells C_i of $\text{Basis}(\mathbb{X}, \mu)$. Replace the requirement that $\{\sigma_0, \dots, \sigma_{k-1}\}$ is prefix-free with the requirement that $\{C_0, \dots, C_{k-1}\}$ is pairwise-disjoint.

Remark 3.9. For the real-valued computable metric spaces L^0, L^1, L^2 , a number of other test functions have been used in the literature. The resulting computable metric spaces are equivalent.

- (1) On (\mathbb{X}, μ) : functions as in Definition 3.8 (for any choice of $\text{Basis}(\mathbb{X}, \mu)$).
- (2) On (\mathbb{X}, μ) : any computable family $\mathcal{A} = \{\varphi_n\}_{n \in \mathbb{N}}$ of bounded computable functions, such that $\varphi_n: \mathbb{X} \rightarrow \mathbb{R}$ is computable uniformly in n , there is a bound C_n uniformly computable in n such that $\|\varphi_n(x)\|_\infty \leq C_n$ for all n , and $\{\varphi_n\}$ is dense in $L^1(\mathbb{X}, \mu)$.
- (3) On (\mathbb{X}, μ) : the set \mathcal{E} of bounded computable Lipschitz functions in [20, Section 2].
- (4) On effectively compact (\mathbb{X}, μ) : any computable family $\mathcal{A} = \{\varphi_n\}_{n \in \mathbb{N}}$ of computable functions which is dense in $C(\mathbb{X})$.
- (5) On effectively compact (\mathbb{X}, μ) : the “polynomials” in [57, 59] closed under pointwise multiplication and addition. (This family is dense in $C(\mathbb{X})$ by the Stone-Weierstrass theorem).
- (6) On $(2^\mathbb{N}, \lambda)$: the test functions in equation (3.1).
- (7) On $([0, 1]^d, \lambda)$: polynomials with rational coefficients.
- (8) On $([0, 1]^d, \lambda)$, (\mathbb{T}^d, λ) : dyadic functions of the form

$$\varphi = \sum_{i=0}^{k-1} c_i \mathbf{1}_{D_i} \quad (c_i \in \mathbb{Q}, D_i \text{ is a dyadic set})$$

where the DYADIC SETS are those of the form

$$\left[\frac{i_1}{2^n}, \frac{i_1 + 1}{2^n} \right) \times \cdots \times \left[\frac{i_d}{2^n}, \frac{i_d + 1}{2^n} \right).$$

3.2. Effective modes of convergence. In measure theoretic probability, there are various modes of convergence for measurable functions. I have already mentioned convergence in the L^1 -, L^2 -norms and the metric d_{meas} .

Definition 3.10. Let (f_i) be a sequence of measurable \mathbb{Y} -valued functions and f a measurable \mathbb{Y} -valued function.

- (1) The sequence f_i converges to f ALMOST UNIFORMLY if there is a RATE OF ALMOST-UNIFORM CONVERGENCE $n(\varepsilon_1, \varepsilon_2)$ such that for all $\varepsilon_1, \varepsilon_2 > 0$,

$$\mu \left(\left\{ x \mid \sup_{i \geq n(\varepsilon_1, \varepsilon_2)} d_{\mathbb{Y}}(f_i(x), f(x)) > \varepsilon_1 \right\} \right) \leq \varepsilon_2.$$

- (2) The sequence f_i converges to f IN MEASURE if there is a RATE OF CONVERGENCE IN MEASURE $n(\varepsilon_1, \varepsilon_2)$ such that for all $\varepsilon_1, \varepsilon_2 > 0$,

$$\forall i \geq n(\varepsilon_1, \varepsilon_2) \quad \mu(\{x \mid d_{\mathbb{Y}}(f_i(x), f(x)) > \varepsilon_1\}) \leq \varepsilon_2.$$


These definition can be extended to continuously-indexed sequences (i.e. functions $(f_r)_{r>0}$ with $r \rightarrow 0$) in the usual manner.

Fact 3.11. *Convergence in measure is the same as convergence in the metric d_{meas} .*

Proof. I give an effective version in Proposition 3.15. (For a similar proof with the Ky-Fan metric, see [13].) \square

Fact 3.12 (Egorov’s theorem, see [13]). *On a probability space, almost uniform convergence and almost everywhere convergence are the same (assuming (f_n) is a discretely-indexed sequence of measurable functions taking values in a complete separable metric space).*

Fact 3.13 (Modes of convergence, see [13]). *On a probability space, the following implications (and their transitive closures) hold between the modes of convergence. (Note, L^2 and L^1 only apply to real-valued functions. The dotted arrow represents convergence on some subsequence.)*




The goal of this section is to give the effective analog of the above chart.

Definition 3.14. Let (f_i) and f be uniformly effectively measurable. Then $f_i \rightarrow f$ EFFECTIVELY ALMOST UNIFORMLY, EFFECTIVELY IN MEASURE, or EFFECTIVELY IN d_{meas} if the respective rate of convergence is computable.

Further if real-valued (f_i) and f are uniformly L^1 -computable (resp. L^2 -computable), then $f_i \rightarrow f$ EFFECTIVELY IN THE L^1 -NORM (resp. EFFECTIVELY IN THE L^2 -NORM) if the corresponding rate of convergence is computable.

Proposition 3.15 (Modes of effective convergence). *On a computable probability space (\mathbb{X}, μ) , the following implications are effective—in that a rate of convergence for the latter is computable from the former. (L^1 and L^2 only apply to real-valued functions.)*



- (1) The dotted arrow represents that if $f_i \rightarrow f$ with a geometric rate of convergence in the metric d_{meas} , e.g. $\forall j \geq i \ d_{meas}(f_j, f) \leq 2^{-i}$, then $f_i \rightarrow f$ effectively almost uniformly.
- (2) For the arrow going to “Schnorr”, see Lemma 3.19 below.

Proof. See Appendix A.3. □

Rather than use the term “effectively almost uniformly”, we will use the more common term EFFECTIVELY ALMOST EVERYWHERE (or EFFECTIVELY A.E.). This is justified by Egorov’s theorem (Fact 3.12).

The following limit properties are also useful.

Proposition 3.16. *Let (f_n) and f be uniformly effectively measurable real-valued functions.*

- (1) *If $f_n \rightarrow f$ effectively a.e. and $g_n \rightarrow g$ effectively a.e., then $f_n + g_n \rightarrow f + g$ effectively a.e..*
- (2) *If $f_n^j \rightarrow f^j$ effectively a.e.. ($j \in \{0, \dots, k-1\}$), and g is computable with a uniform modulus of continuity, then $g(f_n^0, \dots, f_n^{k-1}) \rightarrow g(f^0, \dots, f^{k-1})$ effectively a.e..*
- (3) *(Squeeze theorem) Assume $f_n \leq g_n \leq h_n$ a.e. and that $f_n \rightarrow g$ effectively a.e.. and $h_n \rightarrow g$ effectively a.e., then $g_n \rightarrow g$ effectively a.e..*

Further, in all cases the rates of convergence for the latter are computable from the former (in (2) use the modulus of continuity for g). Indeed, we do not need to assume the functions are effectively measurable, just that the rates of convergence are computable. The same results hold for continuous convergence, e.g. $f_r \rightarrow f$ as $r \rightarrow 0$.

3.3. Convergence on Schnorr randomness. Now we define representatives for each (equivalence class of an) effectively measurable function. The proofs are in Appendix A.4.

Recall that Cauchy-names are computable sequences of test functions with a geometric rate of convergence.

Definition 3.17. Let $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ be effectively measurable with Cauchy-name (φ_n) in the metric d_{meas} . Define

$$\tilde{f}(x) = \begin{cases} \lim_{n \rightarrow \infty} \varphi_n(x) & \text{if the limit exists} \\ \text{undefined} & \text{otherwise} \end{cases}.$$

If A is an effectively measurable set (and therefore $\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \{0, 1\}$ is effectively measurable), then define \tilde{A} as

$$x \in \tilde{A} \iff \tilde{\mathbf{1}}_A(x) = 1.$$

These definitions are justified as follows. Similar versions of this proposition are in Pathak [39] (L^1 -computable functions and Martin-Löf randomness) and Pathak, Rojas, and Simpson [40] (L^1 -computable functions and Schnorr randomness).

Proposition 3.18. *Suppose $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable with Cauchy-name (φ_n) (in the metric d_{meas} , L^1 -norm, or L^2 -norm).*

- (1) *(Existence) The limit $\lim_{n \rightarrow \infty} \varphi_n(x)$ exists on all Schnorr randoms x .*
- (2) *(Uniqueness) Given another Cauchy-name (ψ_n) for f ,*

$$\lim_{n \rightarrow \infty} \varphi_n(x) = \lim_{n \rightarrow \infty} \psi_n(x) \quad (\text{on Schnorr random } x).$$

In Theorem 12.19, I show that Schnorr randomness is the best possible for the previous theorem.

This next lemma is quite useful and for much of the paper is the only fact about Schnorr randomness needed.

Lemma 3.19 (Convergence Lemma). *Suppose that (f_k) and f are uniformly effectively measurable. If*

$$f_k \rightarrow f \quad (\text{effectively a.e.})$$

then

$$\tilde{f}_k(x) \rightarrow \tilde{f}(x) \quad (\text{for all Schnorr random } x).$$

3.4. Properties of effectively measurable functions. The proofs are in Appendix A.5.

Proposition 3.20. *The following implications hold for real-valued functions (and all the computations are uniform).*

- (1) $f \in L^2_{\text{comp}} \Rightarrow f \in L^1_{\text{comp}} \Rightarrow f \in L^0_{\text{comp}}$. *(The converses do not hold in general.)*
- (2) *If $0 \leq f \leq 1$, then $f \in L^2_{\text{comp}} \Leftrightarrow f \in L^1_{\text{comp}} \Leftrightarrow f \in L^0_{\text{comp}}$.*
- (3) $f \in L^1_{\text{comp}} \Leftrightarrow (f \in L^0_{\text{comp}} \text{ and } \|f\|_{L^1} \text{ is computable})$.
- (4) $f \in L^2_{\text{comp}} \Leftrightarrow (f \in L^0_{\text{comp}} \text{ and } \|f\|_{L^2} \text{ is computable})$.
- (5) *If $f \in L^1_{\text{comp}}$ then $\int f d\mu$ is computable.*
- (6) *If B is effectively measurable, then $\mu(B)$ is computable.*
- (7) *If $0 \leq g \leq 1$, $g \in L^1_{\text{comp}}$, and $f \in L^1_{\text{comp}}$, then $g \cdot f \in L^1_{\text{comp}}$.*

Proposition 3.21 (Effective Lusin's theorem). *Given an effectively measurable $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$, and some rational $\varepsilon \geq 0$, there are an effectively closed set K of computable measure $\mu(K) \geq 1 - \varepsilon$ and a computable function $g: K \rightarrow \mathbb{Y}$ such that $g = \tilde{f} \upharpoonright K$ on Schnorr randoms. (Further, g and K are computable uniformly from ε and any name for f .) Moreover, if $\mathbb{Y} = \mathbb{R}$, then $g: K \rightarrow \mathbb{Y}$ can be extended (uniformly from its name) to a total computable function $g: \mathbb{X} \rightarrow \mathbb{Y}$ such that $g = \tilde{f} \upharpoonright K$ on Schnorr randoms.*

Proposition 3.22 (Effective inner/outer regularity). *Given $A \subseteq (\mathbb{X}, \mu)$ effectively measurable, and some rational $\varepsilon > 0$, there are an effectively open set U and an effectively closed set C both of computable measure such that $C \subseteq \tilde{A} \subseteq U$ for Schnorr randoms, and such that $\mu(U) - \mu(C) \leq \varepsilon$. (The sets U, C and their measures $\mu(U), \mu(C)$ are uniformly computable from ε and any name for A .)*

This next result is the converse to the effective Lusin's theorem and shows that the representative functions of this paper are the same as the Schnorr layerwise computable functions of Miyabe [34], which are an extension of the layerwise computable functions of Hoyrup and Rojas [24]. Miyabe [34], proved the corresponding result for L^1 -computable functions.

Proposition 3.23 (Schnorr layerwise computability). *Consider a (pointwise-defined) measurable function $f: \mathbb{X} \rightarrow \mathbb{Y}$ that is SCHNORR LAYERWISE COMPUTABLE, that is, there is a computable sequence (C_n) of effectively closed sets of computable measure $\mu(C_n) \leq 2^{-n}$, such that $f \upharpoonright C_n$ is computable on C_n uniformly in n . Then there is an effectively measurable $g: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ such that $\tilde{g} = f$ on Schnorr randoms.*

In this next proposition, an ALMOST-EVERYWHERE COMPUTABLE FUNCTION $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is a partial computable function whose domain is measure one. (Here I mean “domain” to mean the points x for which the underlying computation computes a name for $f(x)$ from a name for x . To avoid ambiguity, I could alternately define an almost-everywhere computable function as a function $f: A \subseteq \mathbb{X} \rightarrow \mathbb{Y}$ which is computable on a measure-one Π^0_2 set A . See [42] for more discussion.)

Proposition 3.24 (Examples of effectively measurable functions and sets). *All of these functions $f: \mathbb{X} \rightarrow \mathbb{Y}$ and sets $A \subseteq \mathbb{X}$ are effectively measurable, and $\tilde{f} = f$ and $\tilde{A} = A$ on Schnorr randoms.*

- (1) *Test functions and test sets as in Propositions 3.1 and 3.3 and in Definition 3.8.*

- (2) *Computable functions and decidable sets (i.e., computable 0,1-valued functions).*
- (3) *Almost-everywhere computable functions $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and almost-everywhere decidable sets (i.e., almost everywhere computable 0,1-valued functions).*
- (4) *Nonnegative lower semicomputable functions $f: \mathbb{X} \rightarrow \mathbb{R}$ with a computable integral, effectively open sets $U \subseteq \mathbb{X}$ of computable measure, and effectively closed sets $C \subseteq \mathbb{X}$ of computable measure.*

Recall that for a measurable function $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$, the PUSH-FORWARD MEASURE of μ along f (denoted μ_*f) is the measure on \mathbb{Y} defined by $\int \varphi d\mu_*f = \int \varphi \circ f d\mu$ for bounded computable φ .

Proposition 3.25 (Push-forward measures). *If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable, then the push-forward measure (\mathbb{Y}, μ_*f) is a computable probability space (uniformly from (\mathbb{X}, μ) , \mathbb{Y} , and f).*

Proposition 3.26 (Preservation of Schnorr randomness). *If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable and x is Schnorr random, then $\tilde{f}(x)$ is Schnorr random on (\mathbb{Y}, μ_*f) .*

Proposition 3.27 (Composition and tuples).

- (1) (Composition) *Given $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and $g: (\mathbb{Y}, \mu_*f) \rightarrow \mathbb{Z}$ effectively measurable, the composition $g \circ f$ is effectively measurable (uniformly from f and g) and*

$$\widetilde{f \circ g} = \tilde{f} \circ \tilde{g} \quad (\text{on Schnorr randoms}).$$

- (2) (Tuples) *Given $f_n: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}_n$ effectively measurable (uniformly in n), the tuples*

$$(f_0, \dots, f_{k-1}): (\mathbb{X}, \mu) \rightarrow \mathbb{Y}_0 \times \dots \times \mathbb{Y}_{k-1}$$

and

$$(f_n)_{n \in \mathbb{N}}: (\mathbb{X}, \mu) \rightarrow \prod_{n \in \mathbb{N}} \mathbb{Y}_n$$

are effectively measurable (uniformly from (f_n)) and

$$(f_0, \dots, f_{k-1}) = (\tilde{f}_0, \dots, \tilde{f}_{k-1}) \quad \text{and} \quad (\tilde{f}_i)_{i \in \mathbb{N}} = (\tilde{f}_i)_{i \in \mathbb{N}} \quad (\text{on Schnorr randoms}).$$

These two combinations, along with the results about computable functions in Proposition 3.24, can be used to prove a number of useful facts.

Proposition 3.28 (Combinations of measurable functions).

- (1) (Computable pointwise operations). *All computable pointwise operations, including vector, lattice, and Boolean algebra operations preserve effective measurability. Moreover, given $f, g: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ and $A, B \subseteq (\mathbb{X}, \mu)$ effectively measurable, we have*

$$\begin{aligned} \widetilde{f+g} &= \tilde{f} + \tilde{g}, & \widetilde{af} &= a\tilde{f}, & \widetilde{f \cdot g} &= \tilde{f} \cdot \tilde{g} \\ \widetilde{\min(f, g)} &= \min(\tilde{f}, \tilde{g}), & \widetilde{\max(f, g)} &= \max(\tilde{f}, \tilde{g}), & \widetilde{|f|} &= |\tilde{f}| \\ \widetilde{A \cup B} &= \tilde{A} \cup \tilde{B}, & \widetilde{A \cap B} &= \tilde{A} \cap \tilde{B}, & \widetilde{A^c} &= \tilde{A}^c, & \widetilde{\mathbb{X}} &= \mathbb{X}, & \widetilde{\emptyset} &= \emptyset \end{aligned}$$

on Schnorr randoms, and

$$f \leq g \text{ a.e.} \quad \Leftrightarrow \quad \tilde{f} \leq \tilde{g} \quad (\text{on Schnorr randoms})$$

$$A \subseteq B \text{ a.e.} \quad \Leftrightarrow \quad \tilde{A} \subseteq \tilde{B} \quad (\text{on Schnorr randoms}).$$

- (2) (Inverse image) *Given $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and $B \subseteq (\mathbb{Y}, \mu_*f)$ effectively measurable then $f^{-1}(B)$ is effectively measurable and $\widetilde{f^{-1}(B)} = \tilde{f}^{-1}(\tilde{B})$ on Schnorr randoms.*
- (3) (Rotations) *Given $f: (\mathbb{T}^d, \lambda) \rightarrow \mathbb{R}$ effectively measurable, and a computable vector $t \in \mathbb{T}^d$, then $h(x) := f(x - t)$ is effectively measurable and $\tilde{h}(x) = \tilde{f}(x - t)$ on Schnorr randoms.*
- (4) (Indicator functions) *Given $A \subseteq (\mathbb{X}, \mu)$, A is effectively measurable if and only if $\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ is effectively measurable (equivalently, L^1 -computable by Proposition 3.20 (2)) and $x \in \tilde{A}$ if and only if $\tilde{\mathbf{1}}_A(x) = 1$ on Schnorr randoms. (Notice the codomain of $\mathbf{1}_A$ is \mathbb{R} here rather than $\{0, 1\}$ as in Definition 3.17.)*

Proposition 3.29. *The following implications hold for real-valued functions (and all the computations are uniform).*

- (1) If $f \in L^1_{\text{comp}}$ and A is effectively measurable, then $\int_A f d\mu$ is computable.
- (2) If \mathbb{X} is effectively compact (see [36])—as is $[0, 1]^d$, \mathbb{T}^d , and $2^{\mathbb{N}}$ —and $g: \mathbb{X} \rightarrow \mathbb{R}$ is computable, then g is L^1 -computable (since it has computable bounds).
- (3) If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable and $g \in L^1_{\text{comp}}(\mathbb{Y}, \mu_* f)$ (resp. $L^2_{\text{comp}}(\mathbb{Y}, \mu_* f)$), then $g \circ f \in L^1_{\text{comp}}(\mathbb{X}, \mu)$ (resp. $L^2_{\text{comp}}(\mathbb{X}, \mu)$).

Proposition 3.30. *Given a measurable map $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$, the following are equivalent.*

- (1) f is effectively measurable.
- (2) The push-forward measure $(\mathbb{Y}, \mu_* f)$ is computable and one (or all) of the following “pull-back” maps are computable:
 - (a) (L^1 functions) $g \in L^1(\mathbb{Y}, \mu_* f) \mapsto g \circ f \in L^1(\mathbb{X}, \mu)$.
 - (b) (L^2 functions) $g \in L^2(\mathbb{Y}, \mu_* f) \mapsto g \circ f \in L^2(\mathbb{X}, \mu)$.
 - (c) (Measurable sets) $B \subseteq (\mathbb{Y}, \mu_* f) \mapsto f^{-1}(B) \subseteq (\mathbb{X}, \mu)$.

4. DIFFERENTIABILITY

In this section I present effective versions of the Lebesgue differentiation theorem and its corollaries.

4.1. The dyadic Lebesgue differentiation theorem. Before considering the full Lebesgue differentiation theorem, let us consider the simpler dyadic version on the fair-coin measure $(2^{\mathbb{N}}, \lambda)$. This will contain most of the work for the version on $[0, 1]$.

Fact 4.1 (Dyadic Lebesgue differentiation theorem). *Given $f \in L^1(2^{\mathbb{N}}, \lambda)$,*

$$\frac{\int_{[x \upharpoonright k]} |f - f(x)| d\lambda}{\lambda([x \upharpoonright k])} \xrightarrow{k \rightarrow \infty} 0 \quad (\lambda\text{-a.e. } x \in 2^{\mathbb{N}}).$$

In particular

$$\frac{\int_{[x \upharpoonright k]} f d\lambda}{\lambda([x \upharpoonright k])} \xrightarrow{k \rightarrow \infty} f(x) \quad (\lambda\text{-a.e. } x \in 2^{\mathbb{N}}).$$

As a helpful notation, I will write

$$f^{(k)}(x) := \frac{\int_{[x \upharpoonright k]} f d\lambda}{\lambda([x \upharpoonright k])}.$$

Notice $f^{(k)} \in L^1(2^{\mathbb{N}}, \lambda)$ and that $f^{(k)}$ is constant on each cylinder set $[\sigma]$ where $\sigma \in 2^{k'}$ ($k' \geq k$). Further, we can use $f^{(k)}$ to approximate f in the L^1 -norm as follows.

Fact 4.2 (Lebesgue approximation theorem). *Given $f \in L^1(2^{\mathbb{N}}, \lambda)$,*

$$f^{(k)} \xrightarrow[k \rightarrow \infty]{L^1} f.$$

As we will see, Facts 4.1 and 4.2 are both instances of the more general Lévy 0-1 law (Fact 6.2).

Proposition 4.3 (Effective Lebesgue approximation theorem). *Suppose we are given $f \in L^1_{\text{comp}}(2^{\mathbb{N}}, \lambda)$. Then*

$$f^{(k)} \xrightarrow[k \rightarrow \infty]{L^1} f \quad (\text{effectively}).$$

Proof. We compute the rate of convergence $k(\varepsilon)$. Pick a rational $\varepsilon > 0$. Let φ be a simple function approximating f such that $\|f - \varphi\|_{L^1} \leq \varepsilon/2$. By the definition of simple function, there is some k' such that φ is constant on all cylinder sets $[\sigma]$ where $\sigma \in 2^{k'}$ ($k \geq k'$). In particular, $\varphi^{(k)} = \varphi$ ($k \geq k'$). Let $k(\varepsilon) = k'$.

Then for $k \geq k(\varepsilon)$,

$$\begin{aligned} \|f - f^{(k)}\|_{L^1} &\leq \|f - \varphi\|_{L^1} + \|\varphi^{(k)} - f^{(k)}\|_{L^1} \\ &= \|f - \varphi\|_{L^1} + \sum_{\sigma \in 2^k} \left| \frac{\int_{[\sigma]} \varphi - f d\lambda}{\lambda([\sigma])} \right| \\ &\leq \|f - \varphi\|_{L^1} + \sum_{\sigma \in 2^k} \frac{\int_{[\sigma]} |\varphi - f| d\lambda}{\lambda([\sigma])} \\ &= 2 \|f - \varphi\|_{L^1} \leq \varepsilon. \end{aligned}$$

□

Recall the following dyadic version of Kolmogorov's inequality.

Fact 4.4 (Dyadic Kolmogorov's inequality, see [12]). *Let $M: 2^{<\omega} \rightarrow [0, \infty)$ be a nonnegative DYADIC MARTINGALE on the $(2^{\mathbb{N}}, \lambda)$, that is $\frac{1}{2}M(\sigma 0) + \frac{1}{2}M(\sigma 1) = M(\sigma)$ for all $\sigma \in 2^{<\omega}$. Then for all $\varepsilon > 0$*

$$\lambda \left(\left\{ x \in 2^{\mathbb{N}} \mid \sup_{k \geq 0} M(x \upharpoonright k) \geq \varepsilon \right\} \right) \leq \frac{M(\emptyset_{string})}{\varepsilon}.$$

As a special case we have the following.

Lemma 4.5. *Given nonnegative $f \in L^1(2^{\mathbb{N}}, \lambda)$,*

$$\lambda \left(\left\{ x \in 2^{\mathbb{N}} \mid \sup_{k \geq 0} f^{(k)}(x) \geq \varepsilon \right\} \right) \leq \frac{\|f\|_{L^1}}{\varepsilon}.$$

Proof. Let $M(\sigma) = \int_{[\sigma]} f d\lambda / \lambda([\sigma])$. This is a nonnegative dyadic martingale since f is nonnegative. Apply Kolmogorov's inequality noting that $f^{(k)}(x) = M(x \upharpoonright k)$ and $\|f\|_{L^1} = \int f d\lambda = M(\emptyset_{string})$.

Now we have the effective version of Proposition 4.1. □

Proposition 4.6 (Effective dyadic Lebesgue differentiation theorem). *Given $f \in L^1_{comp}(2^{\mathbb{N}}, \lambda)$, let*

$$g_k(x) := |f - f(x)|^{(k)}(x) = \frac{\int_{[x \upharpoonright k]} |f(y) - f(x)| d\lambda(y)}{\lambda([x \upharpoonright k])}.$$

Then $g_k \rightarrow 0$ a.e. as $k \rightarrow \infty$ with an effective rate $k(\delta, \varepsilon)$ of a.e. convergence. Hence $f^{(k)} \rightarrow f$ effectively a.e. as $k \rightarrow \infty$.

Further,

$$\frac{\int_{[x \upharpoonright k]} |f(y) - \tilde{f}(x)| d\lambda(y)}{\lambda([x \upharpoonright k])} \xrightarrow[k \rightarrow \infty]{} 0 \quad (\text{on Schnorr random } x).$$

Hence, $f^{(k)}(x) \rightarrow \tilde{f}(x)$ on Schnorr randoms x as $k \rightarrow \infty$.

Proof. Pick $\delta > 0$ and $\varepsilon > 0$. By Proposition 4.3, from f we can effectively find some $k' \in \mathbb{N}$ such that $\|f - f^{(k')}\|_{L^1} \leq \frac{\delta \varepsilon}{4}$. Let $k(\delta, \varepsilon) = k'$. Then for any $k \geq k'$ and all $x \in 2^{\mathbb{N}}$ we have

$$\begin{aligned} (4.1) \quad 0 \leq g_k(x) &= \frac{\int_{[x \upharpoonright k]} |f(y) - f(x)| d\lambda(y)}{\lambda([x \upharpoonright k])} \\ &\leq \frac{\int_{[x \upharpoonright k]} |f(y) - f^{(k')}(y)| d\lambda(y)}{\lambda([x \upharpoonright k])} + \frac{\int_{[x \upharpoonright k]} |f^{(k')}(y) - f^{(k')}(x)| d\lambda(y)}{\lambda([x \upharpoonright k])} \\ &\quad + \frac{\int_{[x \upharpoonright k]} |f^{(k')}(x) - f(x)| d\lambda(y)}{\lambda([x \upharpoonright k])} \\ &= |f - f^{(k')}|^{(k)}(x) + 0 + |f^{(k')}(x) - f(x)|. \end{aligned}$$

To bound the last line, use Lemma 4.5 for the first term,

$$\lambda \left(\left\{ x \mid \sup_{k \geq k'} |f - f^{(k')}|^{(k)}(x) \geq \frac{\varepsilon}{2} \right\} \right) \leq \frac{2 \|f - f^{(k')}\|_{L^1}}{\varepsilon},$$

and use Markov's inequality (Fact A.2) for the last term,

$$\lambda \left(\left\{ x \mid \sup_{k \geq k'} |f^{(k')}(x) - f(x)| \geq \frac{\varepsilon}{2} \right\} \right) \leq \frac{2 \|f - f^{(k')}\|_{L^1}}{\varepsilon}.$$

Putting them together (see Fact A.1), we have

$$\lambda \left(\left\{ x \mid \sup_{k \geq k'} g_k(x) \geq \varepsilon \right\} \right) \leq 4 \frac{\|f - f^{(k')}\|_{L^1}}{\varepsilon} \leq \delta.$$

Since $g_k \geq 0$, this shows that $g_k(x) \rightarrow 0$ effectively a.e.

To show convergence on Schnorr randoms, we substitute $f(x)$ with $\tilde{f}(x)$ in inequality (4.1). Then we have for all Schnorr randoms x ,

$$0 \leq |f - \tilde{f}(x)|^{(k)}(x) \leq |f - f^{(k')}|^{(k)}(x) + |f^{(k')}(x) - \tilde{f}(x)|.$$

(The other instances of f act as equivalence classes and do not require the tilde). Let $h_k = |f - f^{(k')}|^{(k)}(x) + |f^{(k')}(x) - \tilde{f}(x)|$. Since we showed $h_k \rightarrow 0$ effectively a.e., we have that $\tilde{h}_k(x) \rightarrow 0$ on Schnorr randoms x by Lemma 3.19. By the results in Section 3.4 we have that $\tilde{h}_k(x) = |f - f^{(k')}|^{(k)}(x) + |f^{(k')}(x) - \tilde{f}(x)|$ for Schnorr randoms. (For example, for any f and k we have $\tilde{f}^{(k)} = f^{(k)}$ since $f^{(k)}$ is a computable function.) Hence both $|f - \tilde{f}(x)|^{(k)}(x) \rightarrow 0$ and $f^{(k)}(x) \rightarrow \tilde{f}(x)$ on Schnorr randoms x as $k \rightarrow \infty$. \square

4.2. The Lebesgue differentiation theorem. Now I wish to prove an effective version of the Lebesgue differentiation theorem. To simplify the geometry I will use the unit torus \mathbb{T}^d (identified with $[0, 1]^d$) and the Lebesgue measure λ . The argument for $[0, 1]^d$ is similar. First, recall the Lebesgue differentiation theorem. Here $A_r f(x)$ is the average of f over the ball $B(x, r)$,

$$A_r f(x) = \frac{\int_{B(x, r)} f(y) dy}{\lambda(B(x, r))}.$$

Fact 4.7 (Lebesgue differentiation theorem, see [49]). *Given an integrable function f on (\mathbb{T}^d, λ) ,*

$$(4.2) \quad A_r |f - f(x)| (x) = \frac{\int_{B(x, r)} |f(y) - f(x)| dy}{\lambda(B(x, r))} \xrightarrow[r \rightarrow 0]{} 0 \quad (\lambda\text{-a.e. } x \in \mathbb{T}^d).$$

In particular,

$$A_r f(x) \xrightarrow[r \rightarrow 0]{} f(x) \quad (\lambda\text{-a.e. } x \in \mathbb{T}^d).$$


The points x for which the limit (4.2) holds are the LEBESGUE POINTS of f .

If, instead of averaging over balls, we averaged over dyadic sets, the Lebesgue differentiation theorem would be the dyadic Lebesgue differentiation theorem of Fact 4.1. However, the full Lebesgue differentiation theorem is a geometric theorem. The theorem concerns the *simultaneous* convergence of overlapping balls (or cubes). Moreover, if the balls or cubes were replaced by, say, ellipses or rectangles of arbitrary aspect ratio, the theorem would not hold. The main idea behind any proof of the Lebesgue differentiation theorem is to restrict one's attention to a disjoint set of cubes (or balls). The classical proof does this through Vitali's covering lemma (see [49]). Here I use an alternate method of Morayne and Solecki [35], which uses martingale theory and a useful geometric lemma.

If $t = (t_1, \dots, t_d) \in \mathbb{T}^d$ and $Q \subseteq \mathbb{T}^d$, define $t + Q = \{t + x \mid x \in Q\}$, i.e. Q rotated in each i th coordinate by t_i . Let \mathcal{B}_k denote the set of dyadic cubes of measure $(2^{-k})^d$. Define $\mathcal{B}_k^t = \{t + Q \mid Q \in \mathcal{B}_k\}$, i.e. translate the dyadic cubes by the vector $t \in \mathbb{T}^d$. Let $I_k^t(x)$ be the unique element of \mathcal{B}_k^t that contains x . The next fact and lemma show that it is enough to consider convergence along dyadic cubes and finitely many shifts.

Fact 4.8 (Morayne and Solecki [35, Lemma 2]). *Let $x \in \mathbb{T}^d$. Consider a cube $Q = x + (-\delta, \delta)^d$ such that $0 < \delta < 2^{-k}/3$. Then $Q \subseteq \bigcup_{t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d} I_k^t(x)$.*

Proof sketch. The main idea is that any interval of length 2δ where $\delta < 2^{-k}/3$ must either be contained in a dyadic interval of length 2^{-k} , or in a dyadic interval shifted by $2^{-k}/3$ in either direction as this picture shows.



Then notice a dyadic interval of length 2^{-k} shifted by $2^{-k}/3$ is also a (different) dyadic interval of length 2^{-k} shifted by $1/3$. \square

Lemma 4.9. *Let $x \in \mathbb{T}^d$ and $f \in L^1(\mathbb{T}^d, \lambda)$ (such that f is pointwise defined at x). Then the following are equivalent.*

- (1) $A_r |f - f(x)| (x) \xrightarrow{r \rightarrow 0} 0$ (i.e., x is a Lebesgue point of f).
- (2) $\frac{1}{\lambda(Q_\delta(x))} \int_{Q_\delta(x)} |f(y) - f(x)| dy \xrightarrow{\delta \rightarrow 0} 0$ for $Q_\delta(x) = x + (-\delta, \delta)^d$.
- (3) $\frac{1}{\lambda(Q_i)} \int_{Q_i} |f(y) - f(x)| dy \xrightarrow{i \rightarrow \infty} 0$ for any sequence of cubes $Q_0 \supseteq Q_1 \supseteq \dots$ where $\bigcap_i Q_i = \{x\}$ (the sequence need not be computable).
- (4) $\frac{1}{\lambda(I_k^t(x))} \int_{I_k^t(x)} |f(y) - f(x)| dy \xrightarrow{k \rightarrow \infty} 0$ for all $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$.

(1) through (4) also hold when \mathbb{T}^d is replaced by $[0, 1]^d$.

Proof. We will show (4) implies (2) implies (1). The other equivalences are standard results that follow similarly. Their proofs are left to the reader.

For (2) implies (1), pick $r > 0$ and let $\delta = r$. Then $\lambda(B(x, r)) = \lambda(Q_\delta)/C$ for some constant C depending only on the dimension d , and

$$(4.3) \quad A_r |f - f(x)| (x) = \frac{\int_{B(x, r)} |f(y) - f(x)| dy}{\lambda(B(x, r))} \leq C \cdot \frac{\int_{Q_\delta} |f(y) - f(x)| dy}{\lambda(Q_\delta)}.$$

For (2) implies (1), pick $\delta > 0$ and let k be such that $2^{-k}/3 > \delta \geq 2^{-k-1}/3$. Then $\lambda(Q_\delta(x)) \geq \lambda(I_k^t(x))/3^d$ for all $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$. Therefore, by Lemma 4.8,

$$(4.4) \quad \begin{aligned} \frac{\int_{Q_\delta(x)} |f(y) - f(x)| dy}{\lambda(Q_\delta(x))} &\leq \frac{\sum_{t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d} \int_{I_k^t(x)} |f(y) - f(x)| dy}{\lambda(Q_\delta(x))} \\ &\leq \sum_{t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d} \frac{\int_{I_k^t(x)} |f(y) - f(x)| dy}{\lambda(I_k^t(x))/3^d} = 3^d \cdot \sum_{t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d} \frac{\int_{I_k^t(x)} |f(y) - f(x)| dy}{\lambda(I_k^t(x))}. \quad \square \end{aligned}$$

Theorem 4.10 (Effective Lebesgue differentiation theorem). *Given $f \in L^1_{comp}(\mathbb{T}^d, \lambda)$,*

$$A_r |f - f(x)| (x) = \frac{\int_{B(x, r)} |f(y) - f(x)| dy}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} 0 \quad (\lambda\text{-a.e. } x \in \mathbb{T}^d)$$

with an effective rate of a.e. convergence $r(\delta, \varepsilon)$. Hence $A_r f \xrightarrow{r \rightarrow 0} f$ effectively a.e.

Further,

$$A_r |f - \tilde{f}(x)| (x) \xrightarrow{r \rightarrow 0} 0 \quad (\text{on Schnorr random } x).$$

Hence, all Schnorr randoms are Lebesgue points of \tilde{f} and $A_r f(x) \xrightarrow{r \rightarrow 0} \tilde{f}(x)$ on Schnorr randoms x . These statements also hold when \mathbb{T}^d is replaced by $[0, 1]^d$.

Proof. Combining inequalities (4.3) and (4.4) in the proof of Lemma 4.9 we have for $2^{-k}/3 > r \geq 2^{-(k+1)}/3$ that

$$0 \leq A_r |f - f(x)|(x) \leq C \cdot \sum_{t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d} \frac{\int_{I_k^t(x)} |f(y) - f(x)| dy}{\lambda(I_k^t(x))}$$

for some constant C depending only on the dimension of d . Using Proposition 4.6, with $f(y-t)$ in place of $f(y)$, we have that

$$\frac{\int_{I_k^t(x)} |f(y) - f(x)| dy}{\lambda(I_k^t(x))} \xrightarrow[k \rightarrow \infty]{} 0$$

with an effective rate of a.e. convergence for each $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$. Hence, by the squeeze theorem (Proposition 3.16 (3))— $A_r |f - f(x)|(x) \xrightarrow[r \rightarrow 0]{} 0$ with an effective rate of a.e. convergence. The result for Schnorr randomness follows by a similar argument. (Note that if $h(y) := f(y-t)$ for a computable t , then $\tilde{h}(y) = \tilde{f}(y-t)$ by Proposition 3.28)

For $[0, 1]^d$, just use the same argument (as for \mathbb{T}^d), but also adjust for the error near the boundary (which is straightforward, although somewhat tedious). \square

Remark 4.11. Setting aside computational concerns, this proof of the Lebesgue differentiation theorem is very similar to the standard proof. The key differences are that this proof uses Lemma 4.9 to handle the geometric concerns while the standard proof uses the Vitali covering lemma, and we use Kolmogorov's inequality to show convergence, while the standard proof uses the Hardy-Littlewood maximal lemma. The effective proof of Pathak et al. [40] indeed follows the usual proof. For another method to handle the geometry see Brattka et al. [8].

4.3. Corollaries to the Lebesgue differentiation theorem. From the effective Lebesgue differentiation theorem (Theorem 4.10), we have the following corollaries. Note that all of these have “dyadic” versions on $2^{\mathbb{N}}$ as well.

Let A be a measurable set on and $x \in [0, 1]^d$. We say x is a POINT OF DENSITY of A if

$$\frac{\lambda(A \cap B(x, r))}{\lambda(B(x, r))} \xrightarrow[r \rightarrow 0]{} 1.$$

Then we have the following well-known corollary to the Lebesgue differentiation theorem.

Fact 4.12 (Lebesgue density theorem). *Let A be a measurable set. Almost every $x \in A$ is a point of density.*

Corollary 4.13 (Effective Lebesgue density theorem). *Let A be an effectively measurable set in $[0, 1]^d$. Every Schnorr random in \tilde{A} is a point of density.*

Proof. Assume x is in \tilde{A} and is Schnorr random. By Definition 3.17, $\tilde{\mathbf{1}}_A(x) = 1$. The rest follows from the Lebesgue differentiation theorem (Theorem 4.10) applied to $\mathbf{1}_A$. \square

For the next application of the Lebesgue density theorem, if A, B are subsets of \mathbb{R} then denote $A + B := \{x + y \mid x \in A, y \in B\}$, and similarly for $A - B$.

Fact 4.14 (Steinhaus, see [48]). *Let A and B be measurable subsets of \mathbb{R} with positive Lebesgue measure and let x and y be points of density of A and B , respectively. Then $A + B$ contains an open neighborhood around $x + y$. Therefore, if A has positive Lebesgue measure, then $A - A$ contains an open neighborhood around 0.*

Corollary 4.15. *Let $A, B \subseteq [0, 1]^d$ be effectively measurable with positive measure. If $x \in \tilde{A}$ and $y \in \tilde{B}$ are Schnorr randoms, then there is an open neighborhood in $\tilde{A} + \tilde{B}$ around $x + y$.*

Proof. By the effective Lebesgue density theorem (Corollary 4.15), x and y are points of density. Apply Steinhaus' theorem (Fact 4.14). \square

A function $h: [0, 1] \rightarrow \mathbb{R}$ is said to be ABSOLUTELY CONTINUOUS if it is of the form $F(x) = \int_0^x f(y) dy + F(0)$ for some integrable function f . It is clear that absolute continuity implies continuity. We have yet another corollary to the Lebesgue differentiation theorem.

Fact 4.16 (Lebesgue, see [49]). *An absolutely continuous function F is differentiable a.e. with derivative $\frac{d}{dx} F = f$ a.e.*

We say F is EFFECTIVELY ABSOLUTELY CONTINUOUS if the derivative f is L^1 -computable. (This is equivalent to being a computable point in the Banach space $(AC[0, 1], \|\cdot\|_{AC})$ where $\|F\|_{AC} = |f(0)| + \|f\|_{BV}$. See [18].) If F is effectively absolutely continuous, then it is computable (by the computability of integration). However, not every computable and absolutely continuous function is effectively absolutely continuous. This follows from this next corollary combined with the example of Brattka, Miller, and Nies [8] of a computable absolutely continuous function which is only differentiable on Martin-Löf randoms (which are a proper subset of the Schnorr randoms).

Corollary 4.17. *Assume $z \in [0, 1]$ is Schnorr random and F is effectively absolutely continuous, hence $F(x) = \int_0^x f(y) dy + F(0)$ for all x for some L^1 -computable f . Then F is differentiable at z with derivative $\frac{d}{dx} F|_{x=z} = \tilde{f}(z)$.*

Proof. It suffices to show

$$\frac{F(z + t_i) - F(z)}{t_i} = \frac{\int_z^{z+t_i} f(y) dy}{t_i} \xrightarrow{i \rightarrow \infty} \tilde{f}(z)$$

for any decreasing sequence $t_i \rightarrow 0^+$ (and the same for any increasing sequence $t_i \rightarrow 0^-$). Letting $Q_i = [z, t_i]$, this becomes

$$\frac{\int_{Q_i} f(y) dy}{\lambda(Q_i)} \xrightarrow{i \rightarrow \infty} \tilde{f}(z),$$

which follows from the stronger result

$$\frac{\int_{Q_i} |f(y) - \tilde{f}(z)| dy}{\lambda(Q_i)} \xrightarrow{i \rightarrow \infty} 0.$$

By item (3) in Lemma 4.9, this is equivalent to z being a Lebesgue point of \tilde{f} —which z is by the effective Lebesgue differentiation theorem (Theorem 4.10). \square

Variations of Corollary 4.17 are given in Corollary 6.9, Theorem 9.19, and Corollary 9.20. Further, in Section 12, I will give an example showing that Corollary 4.17 characterizes Schnorr randomness.

Related to absolutely continuous functions is the following theorem about Radon-Nikodym derivatives.

Fact 4.18 (Radon-Nikodym, see [49]). *Let μ be a probability measure on $[0, 1]^d$. If μ is absolutely continuous with respect to λ (i.e. $\lambda(A) = 0$ implies $\mu(A) = 0$ for all Borel-measurable A), then there is a λ -a.e. unique integrable function $\frac{d\mu}{d\lambda}$, called the RADON-NIKODYM DERIVATIVE or DENSITY, such that for all Borel-measurable sets A ,*

$$\mu(A) = \int_A \frac{d\mu}{d\lambda}(x) dx.$$

Fact 4.19 (See [49]). *Let μ be a probability measure on $[0, 1]^d$ that is absolutely continuous with respect to λ . Then*

$$\frac{\mu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \frac{d\mu}{d\lambda}(x) \quad (\lambda\text{-a.e. } x).$$

Given a computable measure μ , absolutely continuous with respect to λ , say that μ is COMPUTABLY NORMABLE relative to λ if and only if $\frac{d\mu}{d\lambda} \in L^1_{comp}(\lambda)$. (See [28, 27] for an equivalent characterization of computably normable using norms.)

Corollary 4.20. *Let μ be a computable probability measure on $[0, 1]^d$ that is absolutely continuous with respect to λ , and computably normable relative to λ . Then*

$$\frac{\mu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \widetilde{\frac{d\mu}{d\lambda}}(x) \quad (\text{on } \lambda\text{-Schnorr random } x).$$

Proof. Since μ is computably normable relative to λ , we have $\frac{d\mu}{d\lambda} \in L^1_{comp}(\lambda)$. So then

$$\frac{\mu(B(x, r))}{\lambda(B(x, r))} = \frac{\int_{B(x, r)} \frac{d\mu}{d\lambda}(x) dx}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \widetilde{\frac{d\mu}{d\lambda}}(x)$$

on Schnorr randoms x by the effective Lebesgue differentiation theorem (Theorem 4.10). \square

An extension of Corollary 4.20 to signed measures is given in Theorem 9.12. In Section 12, I will give an example showing that Corollary 4.20 characterizes Schnorr randomness.

I end this section with an application to effective harmonic analysis. Rescale \mathbb{T} to be $[0, 2\pi)$ and here i will denote $\sqrt{-1}$. Let $f \in L^1(\mathbb{T} \rightarrow \mathbb{C})$ be a complex-valued integrable function on \mathbb{T} . Let $\{\hat{f}(j)\}_{j \in \mathbb{Z}}$ be the complex-valued FOURIER COEFFICIENTS of f , that is

$$\hat{f}(j) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) \cdot e^{-ijt} dt.$$

Then f can be approximated by the following complex-valued trigonometric polynomials $\sigma_n(f)$ (arising from the Fejér kernel)

$$\sigma_n(f)(x) = \frac{1}{n+1} \sum_{k=0}^n \sum_{j=-k}^k \hat{f}(j) e^{ijx}.$$

We have the following theorem of Lebesgue. (Note the definition of Lebesgue point naturally extends to complex-valued functions.)

Fact 4.21 (Lebesgue, see [30]). *If $x \in \mathbb{T}$ is a Lebesgue point of f , then $\sigma_n(f)(x) \rightarrow f(x)$ as $n \rightarrow \infty$.*

To given an effective version, note that f is computable in $L^1(\mathbb{T} \rightarrow \mathbb{C})$ (with a suitable choice of test functions) if and only if both its real and imaginary parts are computable in $L^1(\mathbb{T} \rightarrow \mathbb{R})$. It is worth noting that $\hat{f}(j)$ is computable in \mathbb{C} uniformly from f and j (use the facts in Proposition 3.20 and that e^{-ijt} is bounded and computable), and that $\sigma_n(f)$ is a computable complex-valued function uniformly in f and n .

Corollary 4.22. *If $f \in L^1_{\text{comp}}(\mathbb{T} \rightarrow \mathbb{C})$ and x is Schnorr random, then $\sigma_n(f)(x) \rightarrow \tilde{f}(x)$ as $n \rightarrow \infty$.*

Proof. By the effective Lebesgue differentiation theorem (Theorem 4.10) x is a Lebesgue point of both the real and imaginary parts of f . Therefore x is also a Lebesgue point of f . The rest of the corollary follows from Fact 4.21. \square

5. MARTINGALES IN COMPUTABLE ANALYSIS

The remainder of this paper is devoted to the effective convergence properties of martingales and applications thereof. This section develops the theory of martingales in computable analysis.

So far, we have only used dyadic martingales on $2^{\mathbb{N}}$, i.e. functions $M: 2^{<\omega} \rightarrow \mathbb{R}$ that satisfy $\frac{1}{2}M(\sigma 0) + \frac{1}{2}M(\sigma 1) = M(\sigma)$. As motivation, one may represent a dyadic martingale as a sequence of functions, $M_n(x) = M(x \upharpoonright n)$ for $x \in 2^{\mathbb{N}}$. This alternate notation is the common one used in probability theory and it allows for a much more general class of martingales. We will define what it means for a martingale in this more general sense to be computable.

Throughout this section, fix an arbitrary computable probability space (\mathbb{X}, μ) .

5.1. Conditional expectation. An important concept in probability theory is that of conditional expectation. Recall that a σ -algebra is a collection of sets closed under complement, countable intersection and countable union. The collection \mathcal{B} of Borel sets is a σ -algebra. We will only consider sub- σ -algebras of \mathcal{B} , and we will only consider them up to μ -a.e. equivalence. (Two σ -algebras \mathcal{F}, \mathcal{G} are μ -A.E. EQUIVALENT if every $A \in \mathcal{F}$ is μ -a.e. equivalent to some $B \in \mathcal{G}$, and vice versa. For example, a σ -algebra with only measure 0 and measure 1 sets is equivalent to the trivial σ -algebra $\{\emptyset, \mathbb{X}\}$.) Hence every σ -algebra should be understood as a collection of equivalence classes of measurable sets.

An important type of σ -algebra is one generated by a finite partition $\mathcal{P} = \{Q_0, \dots, Q_{k-1}\}$ of \mathbb{X} (i.e. $\bigcup_{i=0}^{k-1} Q_i = \mathbb{X}$ μ -a.e.). Given such a finite partition \mathcal{P} , and given $f \in L^1(\mathbb{X}, \mu)$, the CONDITIONAL EXPECTATION $\mathbb{E}[f \mid \mathcal{P}] \in L^1(\mathbb{X}, \mu)$ is defined by

$$\mathbb{E}[f \mid \mathcal{P}] := \sum_{i=0}^{k-1} \frac{\int_{Q_i} f d\mu}{\mu(Q_i)} \cdot \mathbf{1}_{Q_i}.$$

We may leave $\mathbb{E}[f \mid \mathcal{P}](x)$ undefined when $x \in Q_i$ and $\mu(Q_i) = 0$, as we only wish to define $\mathbb{E}[f \mid \mathcal{P}]$ as an a.e. equivalence class. Notice that $\mathbb{E}[f \mid \mathcal{P}]$ is a step function constant on each Q_i , and so I will sometimes abuse notation and write $\mathbb{E}[f \mid \mathcal{P}](Q_i) := \frac{1}{\mu(Q_i)} \int_{Q_i} f d\mu$ where convenient. Below, and throughout the paper, $\tilde{\mathbb{E}}[f \mid \mathcal{P}](x)$ will mean \tilde{g} where $g = \mathbb{E}[f \mid \mathcal{P}]$.

Proposition 5.1. *Let $\mathcal{P} = \{Q_0, \dots, Q_{k-1}\}$ be a finite partition of \mathbb{X} into effectively measurable sets, and let f be an L^1 -computable function. Then the following hold.*

- (1) $\mathbb{E}[f \mid \mathcal{P}]$ is L^1 -computable uniformly from (the names for) f and \mathcal{P} .
- (2) The value $\mathbb{E}[f \mid \mathcal{P}](Q_i)$ is computable from f and Q_i .
- (3) $\widetilde{\mathbb{E}}[f \mid \mathcal{P}](x) = \mathbb{E}[f \mid \mathcal{P}](Q_i)$ assuming $x \in \widetilde{Q}_i$ and x is Schnorr random.

Proof. Items (1) and (2) are straightforward. For (3), assume x is Schnorr random and $x \in \widetilde{Q}_i$. Then, $\mu(Q_i) > 0$. By Definition 3.17, $\widetilde{\mathbf{1}}_{Q_i}(x) = 1$. Moreover, $\widetilde{\mathbf{1}}_{Q_j}(x) = 0$ for $j \neq i$ (since, by Proposition 3.28, $\widetilde{Q}_i \cap \widetilde{Q}_j = \widetilde{Q_i \cap Q_j} = \widetilde{\emptyset} = \emptyset$). Then, by Proposition 3.28, we have

$$\widetilde{\mathbb{E}}[f \mid \mathcal{P}](x) = \sum_{j=0}^{k-1} \frac{\int_{Q_j} f \, d\mu}{\mu(Q_j)} \cdot \widetilde{\mathbf{1}}_{Q_j}(x) = \frac{\int_{Q_i} f \, d\mu}{\mu(Q_i)} = \mathbb{E}[f \mid \mathcal{P}](Q_i). \quad \square$$

The definition of conditional expectation can be extended to any σ -algebra. The CONDITION EXPECTATION $\mathbb{E}[f \mid \mathcal{F}]$ is the a.e. unique function $\mathbb{E}[f \mid \mathcal{F}] \in L^1(\mathbb{X}, \mu)$ such that $\int_A \mathbb{E}[f \mid \mathcal{F}](x) \, d\mu(x) = \int_A f \, d\mu$ for all measurable $A \in \mathcal{F}$. (Alternately, $\mathbb{E}[f \mid \mathcal{F}]$ can be defined directly using the Radon-Nikodym derivative.) If \mathcal{F} is the σ -algebra generated by a partition \mathcal{P} , then $\mathbb{E}[f \mid \mathcal{F}] = \mathbb{E}[f \mid \mathcal{P}]$ μ -a.e. The following facts about conditional expectation will be used quite often (sometimes without reference).

Fact 5.2 (See [14, 54]). *Assume $f, g, f_n \in L^1(\mathbb{X}, \mu)$, and $\mathcal{F}, \mathcal{F}_1, \mathcal{F}_2$ are σ -algebras.*

- (1) $\mathbb{E}[f \mid \mathcal{F}]$ is \mathcal{F} -measurable.
- (2) $\int \mathbb{E}[f \mid \mathcal{F}](x) \, d\mu(x) = \int f(x) \, d\mu(x)$.
- (3) If f is \mathcal{F} -measurable, then $\mathbb{E}[f \mid \mathcal{F}] = f$ a.e.
- (4) (Tower property) If $\mathcal{F}_1 \subseteq \mathcal{F}_2$ (as σ -algebras), then $\mathbb{E}[\mathbb{E}[f \mid \mathcal{F}_2] \mid \mathcal{F}_1] = \mathbb{E}[f \mid \mathcal{F}_1]$ a.e.
- (5) If $\int |g(x)f(x)| \, d\mu < \infty$ and g is \mathcal{F} -measurable, then $\mathbb{E}[gf \mid \mathcal{F}] = g \cdot \mathbb{E}[f \mid \mathcal{F}]$ a.e.
- (6) (Linearity) $\mathbb{E}[af + g \mid \mathcal{F}] = a\mathbb{E}[f \mid \mathcal{F}] + \mathbb{E}[g \mid \mathcal{F}]$ a.e.
- (7) If $f \leq g$ a.e., then $\mathbb{E}[f \mid \mathcal{F}] \leq \mathbb{E}[g \mid \mathcal{F}]$ a.e.
- (8) (Conditional Jensen's inequality) $|\mathbb{E}[f \mid \mathcal{F}]| \leq \mathbb{E}[|f| \mid \mathcal{F}]$ a.e. (or replace $|\cdot|$ with any convex function).
- (9) If $\mathcal{F}_1 \subseteq \mathcal{F}_2$ (as σ -algebras), then $\|\mathbb{E}[f \mid \mathcal{F}_1]\|_{L^1} \leq \|\mathbb{E}[f \mid \mathcal{F}_2]\|_{L^1} \leq \|f\|_{L^1}$ (also for the L^2 -norm).
- (10) (Conditional Fatou's lemma) $\mathbb{E}[\limsup_{n \rightarrow \infty} f_n \mid \mathcal{F}] \geq \limsup_{n \rightarrow \infty} \mathbb{E}[f_n \mid \mathcal{F}]$ if there is some $g \in L^1$ such that $f_n \geq g$ for all n .

5.2. L^1 -computable martingales. A FILTRATION (\mathcal{F}_k) is a chain of σ -algebras $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \dots$. We say a filtration (\mathcal{F}_k) CONVERGES to the σ -algebra \mathcal{F}_∞ , written $\mathcal{F}_k \uparrow \mathcal{F}_\infty$, when $\mathcal{F}_\infty = \sigma(\bigcup_k \mathcal{F}_k)$. One example of a filtration is a chain of increasingly fine partitions. The only filtration we will use by name is the filtration generated by the chain of partitions (\mathcal{B}_k) where, on $2^\mathbb{N}$, $\mathcal{B}_k = \{[\tau] \mid |\tau| = k\}$, and on \mathbb{T}^d or $[0, 1]^d$, \mathcal{B}_k is the set of dyadic cubes with side length 2^{-k} . It is clear that $\mathcal{B}_k \uparrow \mathcal{B}$, where \mathcal{B} is the Borel σ -algebra.

A MARTINGALE adapted to a filtration (\mathcal{F}_k) is a sequence of integrable functions (M_k) such that M_k is \mathcal{F}_k -measurable and

$$(5.1) \quad \mathbb{E}[M_{k+1} \mid \mathcal{F}_k] = M_k \text{ a.e.}$$

Assuming the filtration (\mathcal{F}_k) is given by a sequence of partitions (\mathcal{P}_k) , then M_k is constant on all $Q \in \mathcal{P}_k$. We then may write $M(Q)$ for $M_k(x)$ where $x \in Q$.

Example 5.3. Every dyadic martingale $M: 2^\omega \rightarrow \mathbb{R}$ is equivalent to a martingale (M_k) on $(2^\mathbb{N}, \lambda)$ with respect to the filtration (\mathcal{B}_k) , and vice versa, under the translation $M_k(x) = M(x \upharpoonright k)$. It is easy to see condition (5.1) is equivalent to

$$M(\sigma 0)\mu(\sigma 0) + M(\sigma 1)\mu(\sigma 1) = M(\sigma)\mu(\sigma).$$

In algorithmic randomness, it is customary to assume the martingales are non-negative. We do not make that assumption here.

Martingales are useful for their well-behaved convergence properties. Also, they have a natural interpretation in terms of gambling. In general, \mathcal{F}_k is the information known to the gambler at time k , and M_k is the capital of the gambler at time k following a betting strategy given by M .

It is not necessary to refer to a specific filtration when talking about martingales. Any martingale (M_k) is also a martingale with respect to the filtration (\mathcal{F}_k) where

$$\mathcal{F}_k = \sigma(M_0, \dots, M_k) = \sigma\left(\bigcup_{i=0}^k \{M_i^{-1}(A) \mid A \in \mathcal{B}\}\right)$$

i.e. the minimal σ -algebra with respect to which M_0, \dots, M_k are measurable. (In the definition of $\sigma(M_0, \dots, M_k)$, it is sufficient to replace \mathcal{B} with any countable generator of \mathcal{B} .) Hence (M_k) is a martingale (with respect to some filtration) if and only if $\mathbb{E}[M_{k+1} \mid M_0, \dots, M_k] = M_k$ (where $\mathbb{E}[M_{k+1} \mid M_0, \dots, M_k]$ is defined as $\mathbb{E}[M_{k+1} \mid \sigma(M_0, \dots, M_k)]$).

We say a martingale (M_k) is an L^1 -COMPUTABLE MARTINGALE if (M_k) is a computable sequence of L^1 -computable functions.

Last, we mention the general form of Kolmogorov's inequality (compared with Fact 4.4) which extends Markov's inequality (Fact A.2). We will use it quite often.

Fact 5.4 (Kolmogorov's inequality, see [54]). *For a martingale (M_k) , and $n, m \in \mathbb{N}$,*

$$\mu\left(\left\{x \in \mathbb{X} \mid \max_{k \in [n, m]} |M_k(x)| \geq \varepsilon\right\}\right) \leq \frac{\|M_m\|_{L^1}}{\varepsilon}.$$

6. THE LÉVY 0-1 LAW AND UNIFORMLY INTEGRABLE MARTINGALES

6.1. Some martingale convergence theorems. Assume in this section that (\mathbb{X}, μ) is a computable probability space. Consider the following class of martingales.

Example 6.1. If $f \in L^1(\mathbb{X}, \mu)$ and (\mathcal{F}_k) is a filtration, then $\mathbb{E}[f \mid \mathcal{F}_k]$ is a martingale on (\mathcal{F}_k) by Fact 5.2 (4). In the case that $\mathbb{X} = 2^{\mathbb{N}}, \mathbb{T}^d, [0, 1]^d$, then the sequence $f^{(k)}$ from the Section 4.1 is equal to $\mathbb{E}[f \mid \mathcal{B}_k]$.

Fact 6.2 (Lévy 0-1 law, see [14, 54]). *Given a filtration (\mathcal{F}_k) such that $\mathcal{F}_k \uparrow \mathcal{F}_\infty$ and $f \in L^1$, then*

$$\mathbb{E}[f \mid \mathcal{F}_k] \xrightarrow[k \rightarrow \infty]{} \mathbb{E}[f \mid \mathcal{F}_\infty] \quad (L^1 \text{ and a.e.}).$$

Therefore, if f is \mathcal{F}_∞ -measurable, then $\mathbb{E}[f \mid \mathcal{F}_\infty] = f$ a.e. and

$$\mathbb{E}[f \mid \mathcal{F}_k] \xrightarrow[k \rightarrow \infty]{} f \quad (L^1 \text{ and a.e.}).$$

In this section I give an effective version of the Lévy 0-1 law.

Theorem 6.3 (Effective Lévy 0-1 law). *Let (\mathcal{F}_k) be any filtration with limit \mathcal{F}_∞ . Assume $f \in L^1_{\text{comp}}$, $\mathbb{E}[f \mid \mathcal{F}_k]$ is L^1 -computable uniformly in k , and $\mathbb{E}[f \mid \mathcal{F}_\infty] \in L^1_{\text{comp}}$. Then*

$$\mathbb{E}[f \mid \mathcal{F}_k] \xrightarrow[k \rightarrow \infty]{} \mathbb{E}[f \mid \mathcal{F}_\infty] \quad (\text{effectively } L^1 \text{ and effectively a.e.}).$$

Hence, by Lemma 3.19,

$$\tilde{\mathbb{E}}[f \mid \mathcal{F}_k](z) \xrightarrow[k \rightarrow \infty]{} \tilde{\mathbb{E}}[f \mid \mathcal{F}_\infty](z) \quad (\text{on Schnorr random } z).$$

To prove this theorem, we will rely on the following characterization of martingales which converge in the L^1 -norm. A martingale (M_k) is called UNIFORMLY-INTEGRABLE if it satisfies either of the following equivalent conditions.

Fact 6.4 (see [14]). *If (M_k) is a martingale on the filtration (\mathcal{F}_k) the following are equivalent.*

- (1) (M_k) converges in the L^1 -norm.
- (2) There exists $f \in L^1$ such that $M_k = \mathbb{E}[f \mid \mathcal{F}_k]$ a.e. for all k .
- (3) The sequence of functions (M_k) is uniformly integrable, i.e.

$$\sup_k \int_{\{x \in \mathbb{X} \mid |M_k(x)| > C\}} |M_k| d\mu \xrightarrow[C \rightarrow \infty]{} 0.$$

(Condition 3 is will not be used in this paper.) By the Lévy 0-1 law, every uniformly-integrable martingale has a limit. By Fact 6.4, the effective Lévy 0-1 law (Theorem 6.3) follows from the next lemma.

Lemma 6.5. *Assume (M_k) is a uniformly-integrable, L^1 -computable martingale with limit $M_\infty \in L^1_{\text{comp}}$. Then*

$$M_k \xrightarrow[k \rightarrow \infty]{} M_\infty \quad (\text{effectively } L^1 \text{ and effectively a.e.}).$$

Hence, by Lemma 3.19, $\widetilde{M}_k(z) \xrightarrow[k \rightarrow \infty]{} \widetilde{M}_\infty(z)$ for Schnorr randoms z .

Proof. Since we know that, $M_k \xrightarrow[k \rightarrow \infty]{L^1} M_\infty$ and since M_∞, M_k are uniformly L^1 -computable, we can find a subsequence (M_{k_j}) such that for all $j \geq i$ we have $\|M_{k_j} - M_{k_i}\|_{L^1} \leq 2^{-i}$. The subsequence converges effectively in L^1 and a.e. (Proposition 3.15).

First, we show convergence in the L^1 -norm. Fix i . Notice that $N_k := (M_k - M_{k_i})$ is a martingale for $k \geq k_i$. (This is easy to verify using conditional expectation facts (Facts 5.2) and the fact that M_{k_i} is \mathcal{F}_{k_i} -measurable.) The L^1 -norm of the martingale (N_k) is nondecreasing (Facts 5.2) and hence for any $j \geq i$,

$$(6.1) \quad \max_{k \in [k_i, k_j]} \|M_k - M_{k_i}\|_{L^1} \leq \|M_{k_j} - M_{k_i}\|_{L^1} \leq 2^{-i}.$$

Since i and j are arbitrary, this shows (M_n) is effectively Cauchy in the L^1 -norm.

To show effective a.e. convergence, again fix i and use Kolmogorov's inequality (Fact 5.4) on the martingale $N_k := (M_k - M_{k_i})$ to get

$$(6.2) \quad \mu \left(\left\{ x \mid \max_{k \in [k_j, k_i]} |M_k(x) - M_{k_i}(x)| \geq 2^{-i/2} \right\} \right) \leq \frac{\|M_{k_j} - M_{k_i}\|_{L^1}}{2^{-i/2}} \leq 2^{-i/2}.$$

Since i and j are arbitrary, this shows (M_n) is effectively a.e. Cauchy. \square

Remark 6.6. Notice in the case that $\mathbb{X} = 2^{\mathbb{N}}, \mathbb{T}^d, [0, 1]^d$ and $M_k = f^{(k)}$ (as in Section 4.1), then Lemma 6.5 follows from the effective Lebesgue approximation theorem (Proposition 4.3) (L^1 convergence) and the effective dyadic Lebesgue differentiation theorem (Proposition 4.6) (a.e. convergence).

If the martingale is L^2 -computable and L^2 -BOUNDED, i.e. $\sup_k \|M_k\|_{L^2} < \infty$, then it is sufficient to know the L^2 -bound instead of the limit. (This is not true of the L^1 case.)

Fact 6.7 (See [14]). *Assume (M_k) is an L^2 -bounded martingale. Then (M_k) is uniformly-integrable, has a limit M_∞ in the L^2 -norm (and L^1 -norm), and $\sup_k \|M_k\|_{L^2} = \|M_\infty\|_{L^2}$.*

Corollary 6.8. *Assume (M_k) is an L^2 -computable martingale with limit M_∞ and with computable L^2 -bound $b = \sup_k \|M_k\|_{L^2} = \|M_\infty\|_{L^2}$. Then*

$$M_k \xrightarrow[k \rightarrow \infty]{} M_\infty \quad (\text{eff. } L^2, \text{eff. } L^1, \text{eff. a.e.}).$$

Therefore, M_∞ is L^2 - and L^1 -computable (uniformly from (M_k) and b), and $\widetilde{M}_k(z) \xrightarrow[k \rightarrow \infty]{} \widetilde{M}_\infty(z)$ for Schnorr randoms z .

Proof. The space of L^2 -functions is a Hilbert space and the conditional expectation $f \mapsto \mathbb{E}[f \mid \mathcal{F}]$ is a projection onto the space of \mathcal{F} -measurable functions [14]. Therefore, by the Pythagorean theorem, for $k \geq j$,

$$\|M_k - M_j\|_{L^2}^2 = \|M_k\|_{L^2}^2 - \|M_j\|_{L^2}^2 \leq b^2 - \|M_j\|_{L^2}^2.$$

Since the L^2 -bound b is finite and computable, this implies effective convergence in the L^2 -norm and hence in the L^1 -norm as well. Hence the limit is L^1 - and L^2 -computable (uniformly from (M_k) and b). Since (M_k) converges in L^1 , the martingale is uniformly-integrable (Fact 6.4). The rest follows from Lemma 6.5. \square

This gives the following variation of Corollary 4.17.

Corollary 6.9. *Let $F: [0, 1] \rightarrow \mathbb{R}$ be a computable function which is also absolutely-continuous with derivative $f = \frac{d}{dx}F$. Assume that $\|f\|_{L^2}$ is computable. Then f is L^2 -computable (uniformly from F and $\|f\|_{L^2}$), F is effectively absolutely continuous, and F is differentiable on Schnorr randoms.*

Proof. For any non-dyadic real $x \in [0, 1]$, let $x \upharpoonright n$ denote the binary expansion of x truncated at the n th bit and let $0.x \upharpoonright n$ denote the corresponding dyadic rational. Then

$$\frac{d}{dx} F(x) = \lim_{n \rightarrow \infty} \frac{F(2^{-n} + 0.x \upharpoonright n) - F(0.x \upharpoonright n)}{2^{-n}}.$$

The term under the limit is an L^2 -computable martingale as follows. If f is the derivative of F , then

$$\frac{F(2^{-n} + 0.(x \upharpoonright n)) - F(0.(x \upharpoonright n))}{2^{-n}} = \frac{\int_{[x \upharpoonright n]} f d\lambda}{2^{-n}} = f^{(n)}(x)$$

where $f^{(n)}(x)$ is the martingale defined in Section 4.1 (see Example 6.1). Each $f^{(n)}$ is L^2 -computable from F and n since it is a test function. We know $f^{(n)} \xrightarrow[n \rightarrow \infty]{L^1} f$ (Fact 4.2). Since $\|f\|_{L^2}$ is computable, by Corollary 6.8, the derivative f is L^2 -computable and F is effectively absolutely continuous. The rest follows by Corollary 4.17. \square

In Section 12, I will give examples showing that the theorems of this section characterize Schnorr randomness.

7. MORE MARTINGALE CONVERGENCE RESULTS

7.1. Martingale convergence results. A martingale (M_k) is said to be L^1 -BOUNDED if $\sup_k \|M_k\|_{L^1} < \infty$.

The Lévy 0-1 Law above is a special case of the following theorem.

Fact 7.1 (Doob's martingale convergence theorem, see [14, 54]). *If (M_k) is an L^1 -bounded martingale, then M_k converges pointwise a.e. and in measure to an integrable function.*

Example 7.2. If a martingale is uniformly-integrable or nonnegative then it is L^1 -bounded. Indeed, given a uniformly-integrable martingale (M_k) , there is some $f \in L^1$ such that $M_k = \mathbb{E}[f \mid \mathcal{F}_k]$ (Fact 6.4) and $\|\mathbb{E}[f \mid \mathcal{F}_k]\|_{L^1} \leq \|f\|_{L^1}$ (Facts 5.2). For a nonnegative martingale (M_k) , we have (using Facts 5.2) that

$$\|M_k\|_{L^1} = \int M_k d\mu = \int \mathbb{E}[M_k \mid \mathcal{F}_0] d\mu = \int M_0 d\mu = \|M_0\|_{L^1}.$$

While martingale convergence in general is not effective, it can be under certain circumstances. We have already seen the case when the martingale is uniformly-integrable.

Unlike uniform integrability, being merely L^1 -bounded only implies pointwise convergence, not convergence in the L^1 -norm.

Example 7.3. Consider a doubling strategy, whereby the gambler bets all his capital on at each stage until he loses. The limit of his capital is almost-surely zero, but the martingale is nonnegative, so the L^1 -norm stays constant and does not converge in the L^1 -norm.

Now I consider the case when (M_k) is a nonnegative singular supermartingale. A SUPERMARTINGALE (M_k) is an ADAPTED PROCESS, i.e. M_k is \mathcal{F}_k -measurable such that $\mathbb{E}[M_{k+1} \mid \mathcal{F}_k] \leq M_k$ for all k . (A SUBMARTINGALE (M_k) is the same except $\mathbb{E}[M_{k+1} \mid \mathcal{F}_k] \geq M_k$.) Notice, every martingale is a supermartingale (and submartingale). A supermartingale (M_k) is SINGULAR if $M_k(x) \xrightarrow[k \rightarrow \infty]{} 0$ a.e.

Lemma 7.4. *Let M be a nonnegative L^1 -computable singular supermartingale. Then $M_k \xrightarrow[k \rightarrow \infty]{} 0$ effectively a.e., and hence (by Lemma 3.19) $\widetilde{M}_k(z) \xrightarrow[k \rightarrow \infty]{} 0$ for all Schnorr randoms z .*

Proof. By Fact 7.1, $M_k \xrightarrow[k \rightarrow \infty]{} 0$ in measure. Hence we can effectively find a subsequence (k_i) such that (M_{k_i}) converges rapidly the metric d_{meas} (Fact 3.11), namely

$$d_{meas}(M_{k_i}, 0) = \|\min\{|M_{k_i}|, 1\}\|_{L^1} < 2^{-(i+1)}.$$

Fix i . Since M_{k_i} is nonnegative, it follows by Markov's inequality (Fact A.2) that

$$(7.1) \quad \mu\left(\underbrace{\{x \mid 0 \leq M_{k_i}(x) < 1\}}_{=: C_i}\right) \leq 1 - 2^{-(i+1)}.$$

The set C_i in $\sigma(M_{k_i})$, and hence $C_i \in \mathcal{F}_{k_i}$ for any filtration (\mathcal{F}_k) to which (M_k) is adapted. For $k > k_i$ let $N_k := \mathbf{1}_{C_i} M_k$. The following calculation shows that $(N_k)_{k \geq k_i}$ is still a supermartingale adapted to (\mathcal{F}_k) :

$$\mathbb{E}[\mathbf{1}_{C_i} M_{k+1} \mid \mathcal{F}_k] = \mathbf{1}_{C_j} \mathbb{E}[M_{k+1} \mid \mathcal{F}_k] \leq \mathbf{1}_{C_j} M_k \text{ a.e.}$$

(Intuitively what makes N_k a supermartingale is that on C_i , the process (N_k) behaves as the supermartingale (M_k) , and on the complement of C_i , the process (N_k) is the constant zero supermartingale.) The L^1 -norms of nonnegative supermartingales decrease, and therefore for all $k \geq k_j$,

$$\|\mathbf{1}_{C_i} M_k\|_{L^1} \leq \|\mathbf{1}_{C_i} M_{k_i}\|_{L^1} \leq \|\min(M_{k_i}, 1)\|_{L^1} \leq 2^{-(i+1)}.$$

Kolmogorov's inequality (Fact 5.4) also holds for nonnegative supermartingales, and therefore for $j > i$

$$\mu\left(\left\{x \mid \max_{k \in [k_i, k_j]} \mathbf{1}_{C_i}(x) M_k(x) \geq 2^{-(i+1)/2}\right\}\right) \leq \frac{2^{-(i+1)}}{2^{-(i+1)/2}} \leq 2^{-(i+1)/2}.$$

Call this set A_i . Then

$$\mu\left(\left\{x \mid \max_{k \in [k_i, k_j]} M_k(x) \geq 2^{-(i+1)/2}\right\}\right) \leq \mu(A_i) + (1 - \mu(C_i)) \leq 2^{-i/2}.$$

As i and j are arbitrary, $M_k \rightarrow 0$ effectively a.e. \square

Our goal, however, is to show any martingale converges effectively a.e. if the L^1 -bound and the limit are known. To prove this, I will use two complimentary martingale decompositions. In this next decomposition, M_k^+ denotes the nonnegative part of the martingale decomposition, whereas $[M_k]^+$ will mean $\max(M_k, 0)$ —and similarly for M_k^- and $[M_k]^-$. (Whereas (M_k^+) is a martingale, $([M_k]^+)$ is only a submartingale.) Also, for a martingale $N = (N_k)$, denote $\|N\|_{M^1} = \sup_k \|N_k\|_{L^1}$.

Fact 7.5 (Krickeberg Decomposition, see [9, Chapter V, Section 4]). *Let (M_k) be an L^1 -bounded martingale with respect to the filtration (\mathcal{F}_k) . Then there are two nonnegative martingales (M_k^+) and (M_k^-) such that such that $M_k = M_k^+ - M_k^-$ a.e. for all k , and $\|M\|_{M^1} = \|M^+\|_{M^1} + \|M^-\|_{M^1} = \|M_k^+\|_{L^1} + \|M_k^-\|_{L^1}$ for all k . Further, this decomposition is a.e. unique; $M_k^+ = \sup_n \mathbb{E}[[M_n]^+ \mid \mathcal{F}_k]$ a.e.; $M_k^- = \sup_n \mathbb{E}[[M_n]^- \mid \mathcal{F}_k]$ a.e.; $\lim_{k \rightarrow \infty} M_k^+ = [\lim_k M_k]^+$ a.e.; and $\lim_{k \rightarrow \infty} M_k^- = [\lim_k M_k]^-$ a.e.*

Fact 7.6 (Uniformly Integrable/Singular Decomposition, see [9, Chapter V, Section 4]). *Let (M_k) be an L^1 -bounded martingale with respect to the filtration (\mathcal{F}_k) and let $M_\infty = \lim_n M_n$. Then there is a uniformly integrable martingale (M_k^{ui}) and a singular martingale (M_k^s) such that $M_k = M_k^s + M_k^{ui}$ a.e. for all k . Further, this decomposition is a.e. unique; $M_k^{ui} = \mathbb{E}[M_\infty \mid \mathcal{F}_k]$ a.e.; $M_k^s = \mathbb{E}[M_k - M_\infty \mid \mathcal{F}_k]$ a.e.; and $\|M_k\|_{M^1} = \|M_k^s\|_{M^1} + \|M_k^{ui}\|_{M^1}$.*

Remark 7.7. To make the decompositions computable, we need the filtration to be computable. The filtration (\mathcal{F}_k) can be represented by the sequence of operators $f \mapsto \mathbb{E}[f \mid \mathcal{F}_k]$ from L^1 to L^1 . Say that (\mathcal{F}_k) is COMPUTABLE if $f \mapsto \mathbb{E}[f \mid \mathcal{F}_k]$ is a computable operator from L^1 to L^1 uniformly in k . If (\mathcal{P}_k) is a computable chain of computable partitions, where \mathcal{P}_{k+1} is a refinement of \mathcal{P}_k , then the corresponding filtration is computable. Assuming the filtration (\mathcal{F}_k) is computable, the above decompositions are computable using the L^1 -bound $\|M\|_{M^1}$ and the limit M_∞ , respectively, as follows.

Proposition 7.8 (Effective Krickeberg decomposition). *Let (M_k) be an L^1 -computable martingale with respect to a computable filtration (\mathcal{F}_k) . Then the Krickeberg decomposition (M_k^+) , (M_k^-) is computable from (M_k) , (\mathcal{F}_k) , and the L^1 -bound $\|M\|_{M^1}$. Further, the limits $\lim_{k \rightarrow \infty} M_k^+ = [M_\infty]^+$ and $\lim_{k \rightarrow \infty} M_k^- = [M_\infty]^-$ are L^1 -computable from the limit M_∞ .*

Proof. We wish to compute $M_k^+ = \sup_n \mathbb{E}[[M_n]^+ \mid \mathcal{F}_k]$ and $M_k^- = \sup_n \mathbb{E}[[M_n]^- \mid \mathcal{F}_k]$. Note that $\mathbb{E}[[M_n]^+ \mid \mathcal{F}_k]$ is L^1 -computable from n , k , and (\mathcal{F}_k) , since the filtration is computable. To show each supremum is L^1 -computable, fix $\varepsilon > 0$ and k . Then choose $n > k$ such that

$$\|\mathbb{E}[[M_n]^+ \mid \mathcal{F}_k]\|_{L^1} + \|\mathbb{E}[[M_n]^- \mid \mathcal{F}_k]\|_{L^1} > \|M\|_{M^1} - \varepsilon$$

Since $M_k^+ \geq \mathbb{E}[[M_n]^+ | \mathcal{F}_k]$ and $M_k^- \geq \mathbb{E}[[M_n]^- | \mathcal{F}_k]$ for all n , we have

$$\begin{aligned} & \|M_k^+ - \mathbb{E}[[M_n]^+ | \mathcal{F}_k]\|_{L^1} + \|M_k^- - \mathbb{E}[[M_n]^- | \mathcal{F}_k]\|_{L^1} \\ &= \|M_k^+\|_{L^1} + \|M_k^-\|_{L^1} - (\|\mathbb{E}[[M_n]^+ | \mathcal{F}_k]\|_{L^1} + \|\mathbb{E}[[M_n]^- | \mathcal{F}_k]\|_{L^1}) \\ &\leq \|M\|_{M^1} - \|\mathbb{E}[[M_n]^+ | \mathcal{F}_k] + \mathbb{E}[[M_n]^- | \mathcal{F}_k]\|_{L^1} \leq \varepsilon. \end{aligned}$$

Hence M_k^+ and M_k^- are L^1 -computable uniformly in k .

To compute the limits, just use that fact that $[M_\infty]^+$ and $[M_\infty]^-$ are L^1 -computable from M_∞ . \square

Proposition 7.9 (Effective Uniformly Integrable/Singular Decomposition). *Let (M_k) be an L^1 -computable martingale with respect to a computable filtration (\mathcal{F}_k) . Then the decomposition $(M_k^{ui}), (M_k^s)$ is computable from (M_k) , (\mathcal{F}_k) , and the limit M_∞ . Further, the L^1 -bound $\|M^s\|_{M^1} = \|M\|_{M^1} - \|M^{ui}\|_{M^1} = \|M\|_{M^1} - \|M_\infty\|_{L^1}$ is computable from $\|M\|_{M^1}$.*

Proof. Since the filtration is computable, $M_k^{ui} = \mathbb{E}[M_\infty | \mathcal{F}_k]$ is computable in the L^1 -norm uniformly from M_∞ , k , and (\mathcal{F}_k) . Then $M_k^s = M_k - M_k^{ui}$ is computable in the L^1 -norm. To compute $\|M^s\|_{M^1}$ just use that $\|M^{ui}\|_{M^1} = \|M_\infty\|_{L^1}$ is computable. \square

In the martingale convergence results so far, there have been no computability requirements on the filtration (\mathcal{F}_k) . We can continue to work without specifying the computability of the filtration. The trick is to approximate M by a different martingale whose filtration is given by a chain of partitions.

Proposition 7.10. *Let M be an L^1 -computable martingale (resp. supermartingale, submartingale). There is a computable martingale (resp. supermartingale, submartingale) N adapted to a computable chain of computable partitions (\mathcal{P}_k) such that for all k , $\mathcal{P}_k \subseteq \sigma(M_0, \dots, M_k)$ and $\|N_k - M_k\|_{L^1} \leq 2^{-k}$. If M is nonnegative, then so is N . Further, if M is a martingale or nonnegative submartingale, then $\sup_n \|M_n\|_{L^1} = \sup_n \|N_n\|_{L^1}$.*

Proof. The main idea is to take each σ -algebra in the canonical filtration $\mathcal{F}_k = \sigma(M_0, \dots, M_k)$ and approximate it with a finite sub- σ -algebra, i.e. a partition $\mathcal{P}_k \subseteq \mathcal{F}_k$.

For each k , let $T_k: (\mathbb{X}, \mu) \rightarrow \mathbb{R}^{k+1}$ be the map $T_k = (M_0, \dots, M_k)$. Recall that $\sigma(M_0, \dots, M_k) = \sigma(T_k) = \sigma(\{T_k^{-1}(B) \mid B \in \mathcal{C}\})$ where $\sigma(\mathcal{C})$ is generated by the Borel sigma algebra on the push forward measure space $(\mathbb{R}^{k+1}, \mu_* T_k)$. Recall that $\mu_* T_k$ is computable (Proposition 3.25) and therefore we can take $\mathcal{C} = \text{Basis}(\mathbb{R}^{k+1}, \mu_* T_k)$ as in Lemma 3.5. Let $\{B_i^k\}_i$ be a computable enumeration of $\text{Basis}(\mathbb{R}^{k+1}, \mu_* T_k)$. Then by Proposition 3.20, $\{T_k^{-1}(B_i^k)\}_{i,k}$ is a computable double sequence of effectively measurable sets which generates $\sigma(M_0, \dots, M_k)$. That is, if $\mathcal{Q}_i^k = \{T_k^{-1}(B_0^k), \dots, T_k^{-1}(B_{i-1}^k)\}$, then $\sigma(\mathcal{Q}_i^k) \uparrow_{i \rightarrow \infty} \sigma(M_0, \dots, M_k)$.

By the Lévy 0-1 law (Fact 6.2), $\mathbb{E}[M_k | \mathcal{Q}_i^k] \xrightarrow{i \rightarrow \infty} M_k$. Since each $\mathbb{E}[M_k | \mathcal{Q}_i^k]$ is L^1 -computable from i and k , find some i_k such that $\|\mathbb{E}[M_k | \mathcal{Q}_{i_k}^k] - M_k\|_{L^1} \leq 2^{-k}$. Define $\mathcal{P}_k = \mathcal{Q}_{i_k}^k$ and $N_k = \mathbb{E}[M_k | \mathcal{P}_k]$.

If M is a supermartingale, then N is as well. Indeed, by two applications of the tower property (Facts 5.2),

$$\begin{aligned} \mathbb{E}[N_{k+1} | \mathcal{P}_k] &= \mathbb{E}[\mathbb{E}[M_{k+1} | \mathcal{P}_{k+1}] | \mathcal{P}_k] && \text{(definition of } N_{k+1}) \\ &= \mathbb{E}[M_{k+1} | \mathcal{P}_k] && \text{(tower property)} \\ &= \mathbb{E}[\mathbb{E}[M_{k+1} | \sigma(M_0, \dots, M_k)] | \mathcal{P}_k] && \text{(tower property)} \\ &\leq \mathbb{E}[M_k | \mathcal{P}_k] && (M \text{ is a supermartingale)} \\ &= N_k && \text{(definition of } N_k). \end{aligned}$$

If M is a martingale, or submartingale, the same argument works.

In general, $\|N_k\|_{L^1} = \|\mathbb{E}[M_k | \mathcal{P}_k]\|_{L^1} \leq \|M_k\|_{L^1}$ which is just a property of conditional expectation (Facts 5.2). Moreover, $\|\|M_k\|_{L^1} - \|N_k\|_{L^1}\| \leq \|M_k - N_k\|_{L^1}$. If M is a martingale or nonnegative submartingale, then $\|M_k\|_k$ is increasing and hence $\sup_n \|N_k\|_k = \sup_n \|M_k\|_k$. \square

Theorem 7.11. *Let M be an L^1 -computable martingale with computable L^1 -bound $\|M\|_{M^1}$ and L^1 -computable limit M_∞ . Then $M_k \xrightarrow{k \rightarrow \infty} M_\infty$ effectively a.e., and hence, by Lemma 3.19, $\widetilde{M}_k(z) \xrightarrow{k \rightarrow \infty} \widetilde{M}_\infty(z)$ for all Schnorr randoms z .*

Proof. Let N be as in Proposition 7.10. Since $\|N_k - M_k\|_{L^1} \leq 2^{-k}$ for all k , $(N_k - M_k) \xrightarrow[k \rightarrow \infty]{} 0$ effectively a.e. It follows that $M_k \xrightarrow[k \rightarrow \infty]{} M_\infty$ effectively a.e. if and only if $N_k \xrightarrow[k \rightarrow \infty]{} M_\infty$ effectively a.e.

Since N is a martingale with respect to a computable sequence of partitions, N is effectively decomposable (Proposition 7.9) into a uniformly integrable part N^{ui} and a singular part N^s . We know $N_k^{ui} \xrightarrow[k \rightarrow \infty]{} M_\infty$ converges effectively a.e. by the effective Lévy 0-1 law (Theorem 6.3).

Since $\|M\|_{M^1}$ is computable, then so is $\|N^s\|_{M^1}$. Therefore, N^s can be effectively decomposed (Proposition 7.8) into two nonnegative L^1 -computable singular martingales N^{s+} and N^{s-} . By Lemma 7.4, $N_k^{s+} \xrightarrow[k \rightarrow \infty]{} 0$ and $N_k^{s-} \xrightarrow[k \rightarrow \infty]{} 0$ effectively a.e.

Putting this all together we have that $N_k = N_k^{ui} + N_k^{s+} - N_k^{s-} \xrightarrow[k \rightarrow \infty]{} M_\infty$ effectively a.e. \square

In Section 12, I show that Lemma 9.6 (and hence Theorem 7.11) characterizes Schnorr randomness.

8. SUBMARTINGALES AND SUPERMARTINGALES

Recall from the previous section, a sequence (X_k) of integrable functions is a submartingale (resp. supermartingale) adapted to a filtration (\mathcal{F}_n) if X_k is \mathcal{F}_k -measurable for all k , and $\mathbb{E}[X_{k+1} \mid \mathcal{F}_k] \geq X_k$ (resp. $\mathbb{E}[X_{k+1} \mid \mathcal{F}_k] \leq X_k$) for all k .

It can be shown that L^1 -computable, nonnegative submartingales and supermartingales converge effectively a.e. when their L^1 -bounds and limits are known. The proofs are different for each.

Theorem 8.1. *Let (X_n) be a nonnegative L^1 -computable supermartingale whose limit X_∞ is L^1 -computable. Then $X_n \xrightarrow[n \rightarrow \infty]{} X_\infty$ effectively a.e. and, by Lemma 3.19, $\tilde{X}_n(x) \xrightarrow[n \rightarrow \infty]{} \tilde{X}_\infty(x)$ on Schnorr randoms x . (Instead assuming X_n is nonnegative, we may assume that $X_n \geq Z$ for some integrable function Z .)*

Proof. As in the proof of Theorem 7.11, we may use Proposition 7.10 to assume, without loss of generality, that X_n is adapted to a computable filtration. By the fact that (X_n) is a supermartingale, the fact that X_n is nonnegative (or bounded from below by an integrable function Z), and the conditional Fatou's theorem (Facts 5.2), we have

$$X_n \geq \liminf_k \mathbb{E}[X_k \mid \mathcal{F}_n] \geq \mathbb{E}[X_\infty \mid \mathcal{F}_n].$$

Then we have a nonnegative, L^1 -computable supermartingale $Y_n = X_n - \mathbb{E}[X_\infty \mid \mathcal{F}_n]$ which converges to 0 a.e. But Y_n converges to 0 effectively a.e. by Lemma 7.4. Also $\mathbb{E}[X_\infty \mid \mathcal{F}_n]$ converges effectively a.e. by the effective Lévy 0-1 law. Putting them together completes the proof. \square

For the submartingale case, I first use an effective version of the monotone convergence theorem.

Proposition 8.2 (Effective monotone convergence theorem). *Assume f_n is an nondecreasing sequence of L^1 -computable functions. Also assume $\sup_n \|f_n\|_{L^1}$ is finite and computable. Then $f_n \rightarrow \sup_n f_n$ effectively in the L^1 -norm and effectively a.e. By Lemma 3.19, $\tilde{f}_n \rightarrow \widetilde{\sup_n f_n}$ (or equivalently $\sup_n \tilde{f}_n = \widetilde{\sup_n f_n}$) on Schnorr randoms.*

Proof. Find a subsequence (n_k) such that $(\sup_n \|f_n\|_{L^1}) - \|f_{n_k}\|_{L^1} \leq 2^{-k}$. Fix k . By monotonicity, $\|f_n - f_{n_k}\|_{L^1} \leq 2^{-k}$ for all $n \geq n_k$. Also, by monotonicity, Markov's inequality, and the monotone convergence theorem,

$$\begin{aligned} \mu \left(\left\{ \sup_n |f_n - f_{n_k}| > 2^{-k/2} \right\} \right) &= \mu \left(\left\{ \left(\sup_n f_n \right) - f_{n_k} > 2^{-k/2} \right\} \right) \\ &\leq \frac{\|\sup_n f_n - f_{n_k}\|_{L^1}}{2^{-k/2}} \\ &= \frac{\sup \|f_n\|_{L^1} - \|f_{n_k}\|_{L^1}}{2^{-k/2}} \leq 2^{-k/2}. \end{aligned}$$

Since k is arbitrary, this gives effective convergence in L^1 and effective a.e. convergence. \square

I also use an effective version of Doob's decomposition theorem.

Fact 8.3 (Doob decomposition, see [54]). *Let (X_n) be a submartingale with respect to (\mathcal{F}_n) . Then there is a martingale (M_n) with respect to (\mathcal{F}_n) and a PREDICTABLE PROCESS A_n (i.e. A_{n+1} is \mathcal{F}_n measurable) such that $A_0 = 0$ and $X_n = M_n + A_n$. Moreover, this decomposition is a.e. unique; $A_{n+1} - A_n = \mathbb{E}[X_{n+1} - X_n \mid \mathcal{F}_n]$; and A_n is nondecreasing.*

Proposition 8.4 (Effective Doob Decomposition). *If (X_n) is an L^1 -computable submartingale and (\mathcal{F}_n) is a computable filtration, then the Doob decomposition is effective.*

Proof. It is enough that $\mathbb{E}[X_{n+1} - X_n \mid \mathcal{F}_n]$ is L^1 -computable from the parameters. \square

Theorem 8.5. *Let (X_n) be a nonnegative, L^1 -computable submartingale such that the L^1 -bound $\sup_n \|X_n\|_{L^1}$ is computable and the limit X_∞ is L^1 -computable. Then $X_n \xrightarrow[n \rightarrow \infty]{} X_\infty$ effectively a.e. and $\tilde{X}_n(x) \xrightarrow[n \rightarrow \infty]{} \tilde{X}_\infty(x)$ on Schnorr randoms x .*

Proof. With out loss of generality, the filtration (\mathcal{F}_n) is one of partitions (Proposition 7.10; the same argument holds for submartingales). Then decompose $X_n = M_n + A_n$ as in the effective Doob decomposition (Proposition 8.4). Notice that $0 \leq A_n \leq X_n$ using induction on the formula for A_n , hence both (M_n) and (A_n) are nonnegative. Recall, also that $\|M_n\|_{L^1}$ is nondecreasing in n since (M_n) is a martingale, and $\|A_n\|_{L^1}$ is nondecreasing since (A_n) is nondecreasing. Hence

$$\sup_n \|X_n\|_{L^1} = \sup_n (\|M_n\|_{L^1} + \|A_n\|_{L^1}) = \left(\sup_n \|M_n\|_{L^1} \right) + \left(\sup_n \|A_n\|_{L^1} \right)$$

Since each term is lower semicomputable and $\sup_n \|X_n\|_{L^1}$ is computable, both $\sup_n \|M_n\|_{L^1}$ and $\sup_n \|A_n\|_{L^1}$ are computable.

Moreover, let $X_\infty, M_\infty, A_\infty$ be the limits of $(X_n), (M_n), (A_n)$, respectively. Clearly, $X_\infty = M_\infty + A_\infty$. Notice X_∞ is L^1 -computable by assumption, and A_∞ is L^1 -computable by the effective monotone convergence theorem (Proposition 8.2). Hence M_∞ is L^1 -computable. Therefore, the convergence of (M_n) and (A_n) is effective a.e. using the effective convergence theorem for martingales (Theorem 7.11) and the effective monotone convergence theorem (Proposition 8.2). Convergence on Schnorr randoms follows similarly. \square

In Section 12, I will show these theorems characterize Schnorr randomness.

These theorems both require a lower bound and are not as general as they could be. We leave the following open problem.

Problem 8.6. Let (X_n) be a nonnegative, L^1 -computable submartingale (or supermartingale) such that the L^1 -bounds $\sup_n \|X_n\|_{L^1}$ and $\inf_n \|X_n\|_{L^1}$ are computable and the limit X_∞ is L^1 -computable. Does (X_n) converge to X_∞ effectively a.e.? What if $\|X_n\|_{L^1}$ is computable? What if the rate of convergence of $\|X_n\|_{L^1}$ is computable?

9. MORE DIFFERENTIABILITY RESULTS

In this section we will explore some more differentiability-type results. The results follow from Sections 6 and 7. In some cases, we only sketch the details.

9.1. Signed measures and Radon-Nikodym derivatives. Signed measures are (informally) measures that may assign positive or negative mass to sets. A signed measure ν has a total variation norm $\|\nu\|_{TV}$ that represents the sum of both the positive and negative mass. If μ is a positive measure on $[0, 1]^d$ (i.e. a measure that gives nonnegative mass to every set), then $\|\mu\|_{TV} = \mu([0, 1]^d)$. We will only consider finite signed measures, i.e. where $\|\mu\|_{TV} < \infty$. The (finite) signed measures can be characterized by the Riesz representation theorem as follows. We will use this as our definition of SIGNED MEASURE.

Fact 9.1 (Riesz representation theorem, see [49]). *There is a one-to-one correspondence between (finite) signed measures ν on $[0, 1]^d$ and bounded linear functionals $T: \mathcal{C}([0, 1]^d) \rightarrow \mathbb{R}$, namely each T is the integration map $f \mapsto \int f d\nu$ of a signed measure ν . Further, $\|\nu\|_{TV}$ is equal to the operator norm $\|T\| := \sup_{f \in \mathcal{C}([0, 1]^d)} |T(f)| / \|f\|_\infty$.*

Definition 9.2. A signed measure ν is said to be COMPUTABLE if the corresponding functional T_ν is computable (i.e. $\int f d\nu$ is computable uniformly from f).¹³

Remark 9.3. If T_ν is positive (i.e. $T_\nu(f) \geq 0$ when $f \geq 0$), then ν is a POSITIVE MEASURE and $\|\nu\|_{TV} = T_\nu(\mathbf{1}_{[0,1]^d})$, which is computable from T_ν . A little thought reveals that the positive, computable signed measures are precisely the computable measures of Definition 2.4. Similarly, the positive, computable signed measures with norm one are precisely the computable probability measures.

Recall that λ denotes the Lebesgue measure. In this next fact, which extends Fact 4.19, ν -a.e. means outside a measurable set C such that $\nu(B) = 0$ for all measurable $B \subseteq C$.

Fact 9.4 (Radon-Nikodym theorem and decomposition, see [49]). *Given a signed measure ν on $[0, 1]^d$, there is a λ -a.e. unique, λ -integrable function f and a ν -a.e. unique, λ -null set D such that for all measurable sets A ,*

$$\nu(A) = \int_A f d\lambda + \nu(A \cap D).$$

The function f is the RADON-NIKODYM DERIVATIVE $d\nu/d\lambda$.

Fact 9.5 (See [49]). *Let ν be a signed measure on $[0, 1]^d$. Then*

$$\frac{\nu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \frac{d\nu}{d\lambda}(x) \quad (\lambda\text{-a.e. } x).$$

When ν is a nonnegative absolutely continuous measure, Fact 9.5 is equivalent to Fact 4.19, which is a version of the Lebesgue differentiation theorem (Fact 4.7). An effective version of Fact 9.5 will be given in Theorem 9.12, but first consider the “singular” case where $d\nu/d\lambda = 0$.

Lemma 9.6. *If μ is a positive measure on $[0, 1]^d$ such that $d\mu/d\lambda = 0$ then*

$$\frac{\mu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} 0$$

effectively a.e. and for all λ -Schnorr randoms x .

Proof sketch. Without loss of generality we may work on (\mathbb{T}^d, λ) . By modifying the argument in Lemma 4.9, it is enough to show on λ -Schnorr randoms x that

$$\frac{\mu(I_k^t(x))}{\lambda(I_k^t(x))} \xrightarrow{k \rightarrow \infty} 0$$

for all $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$. However, $\mu(I_k^t(x))$ may not be computable, which happens when the boundary of the cube $I_k^t(x)$ has positive mass. To handle this, replace $I_k^t(x)$ by $I_k^{t+s}(x)$ (that is the dyadic cube shifted by $t+s$ that contains x) for some computable vector $s \in [0, 1]^d$, such that $\mu(I_k^{t+s}(x))$ is computable for all $k \in \mathbb{N}$ and all $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$. One can show, by a diagonalization argument, that there is such an s .

Fix such an s . It is enough to show that

$$\frac{\mu(I_k^{t+s}(x))}{\lambda(I_k^{t+s}(x))} \xrightarrow{k \rightarrow \infty} 0.$$

We have $M_k^t(x) := \mu(I_k^{t+s}(x))/\lambda(I_k^{t+s}(x))$ is a nonnegative, singular, L^1 -computable martingale for each $t \in \{-\frac{1}{3}, 0, \frac{1}{3}\}^d$. The statement of the lemma follows from Lemma 7.4. \square

To effectivize Fact 9.5 in its full generality, I will use two decompositions, which are analogies to the martingale decompositions in Section 7.

¹³In general, the norm $\|\nu\|_{TV}$ is only lower semicomputable, so the space of signed measures is not a computable Banach space. The representation I am using implicitly uses the weak-* topology (or topology of pointwise convergence) on the space of bounded linear functionals of $\mathcal{C}([0, 1])$. That is the minimal topology for which each T_ν is continuous. The unit ball in this topology is metrizable and one could alternately use this fact to classify the computable signed measures as the computable points in the corresponding computable metric space.

Fact 9.7 (Lebesgue decomposition, see [13]). *Given a signed measure ν on $[0, 1]^d$, there is a unique decomposition of ν into two signed measures ν_{ac} and ν_s (the absolutely continuous part and the singular part, respectively) such that $\nu = \nu_{ac} + \nu_s$; $\nu_{ac}(A) = \int_A (d\nu_{ac}/d\lambda) d\lambda$; and $d\nu_s/d\lambda = 0$. Further, $\|\nu\|_{TV} = \|\nu_{ac}\|_{TV} + \|\nu_s\|_{TV}$; if f and D are as in the Radon-Nikodym theorem (Fact 9.4), then $d\nu_{ac}/d\lambda = f$ and for all measurable A ,*

$$\nu_{as}(A) = \int_A f d\lambda \quad \text{and} \quad \nu_s(A) = \nu(A \cap D).$$

Recall, the notation $[f]^+ = \max\{f, 0\}$ and $[f]^- = \max\{-f, 0\}$.

Fact 9.8 (Jordan decomposition, see [13]). *Given a signed measure ν on $[0, 1]^d$, there is a unique decomposition of ν into two signed measures ν^+ and ν^- such that $\nu = \nu^+ - \nu^-$, $\|\nu\|_{TV} = \|\nu^+\|_{TV} + \|\nu^-\|_{TV}$. Further $d\nu^+/d\lambda = [d\nu/d\lambda]^+$, $d\nu^-/d\lambda = [d\nu/d\lambda]^-$.*

Denote $|\nu| = \nu^+ + \nu^-$. The Jordan decomposition is related to the Hahn Decomposition.

Fact 9.9 (Hahn decomposition, see [13]). *Given a signed measure ν on $[0, 1]^d$, there is a unique partition of $[0, 1]^d$ into measurable sets N, P such that $\nu^+(A) = \nu(A \cap P)$ and $\nu^- = \nu(A \cap N)$.*

Here are effective versions of the Lebesgue and Jordan decompositions.

Proposition 9.10 (Effective Lebesgue decomposition). *Let ν be a computable signed measure on $[0, 1]^d$ such that $d\nu/d\lambda$ is L^1 -computable. Then the Lebesgue decomposition ν_{ac} , ν_s is computable. Further, $\|\nu_{ac}\|_{TV}$ and $\|\nu_s\|_{TV}$ are computable from $\|\nu\|_{TV}$.*

Proof. It is easy to see that ν_{ac} , defined by $\nu_{ac}(A) = \int_A f d\lambda$, is a computable signed measure where f is the L^1 -computable Radon-Nikodym derivative. Then define $\nu_s := \nu - \nu_{ac}$.

Notice $\|\nu_{ac}\|_{TV} = \|f\|_{L^1}$, so $\|\nu_s\|_{TV} = \|\nu\|_{TV} - \|\nu_{ac}\|_{TV}$ is computable when $\|\nu\|_{TV}$ is computable. \square

Let ν be a computable signed measure on $[0, 1]^d$ such that $\|\nu\|_{TV}$ is computable. Then the Lebesgue decomposition ν^+ , ν^- is computable. Further, if $d\nu/d\lambda$ is L^1 -computable, then the Radon-Nikodym derivatives $d\nu^+/d\lambda = [d\nu/d\lambda]^+$ and $d\nu^-/d\lambda = [d\nu/d\lambda]^-$ are L^1 -computable. (Further, P and N are effectively measurable in the probability measure $|\nu|/\|\nu\|_{TV}$.)

Proof. The proof is very similar to Proposition 7.8. Using the total variation of ν , the Riesz representation, and the fact that computable functions are dense in $\mathcal{C}([0, 1]^d)$, we can effectively find a computable function $f: [0, 1]^d \rightarrow [-1, 1]$ such that $\|\nu\|_{TV} - \int f d\nu \leq \varepsilon$ for any ε . This function approximates the Hahn decomposition $\mathbf{1}_P - \mathbf{1}_N$. Notice for any computable $\varphi: [0, 1]^d \rightarrow [0, 1]$, we have by nonnegativity,

$$\int \varphi d\nu^+ \geq \int \varphi \cdot [f]^+ d\nu^+ \geq \int \varphi \cdot [f]^+ d\nu^+ - \int \varphi \cdot [f]^+ d\nu^- = \int \varphi \cdot [f]^+ d\nu$$

and similarly $\int \varphi d\nu^- \geq \int -\varphi \cdot [f]^- d\nu$. Then we have

$$\begin{aligned} & \left| \int \varphi d\nu^+ - \int \varphi \cdot [f]^+ d\nu \right| + \left| \int \varphi d\nu^- - \int -\varphi \cdot [f]^- d\nu \right| \\ &= \int \varphi \cdot (1 - f) d\nu^+ - \int \varphi \cdot (1 + f) d\nu^+ \\ &\leq \int (1 - f) d\nu^+ - \int (1 + f) d\nu^+ \quad (-1 \leq f \leq 1) \\ &= \int d|\nu| - \int f d\nu \leq \varepsilon. \end{aligned}$$

Hence ν^+ and ν^- are computable from $\|\nu\|_{TV}$ and ν . Moreover, this shows that $\mathbf{1}_P - \mathbf{1}_N$ is L^1 -computable in $|\nu|/\|\nu\|_{TV}$ and therefore P and N are effectively measurable.

If $d\nu/d\lambda$ is L^1 -computable, then so are $[d\nu/d\lambda]^+$ and $[d\nu/d\lambda]^-$ (Proposition 3.1). \square

Theorem 9.12. *If ν is a computable signed measure such that $\|\nu\|_{TV}$ is computable and $d\nu/d\lambda$ is L^1 -computable, then*

$$\frac{\nu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \frac{d\nu}{d\lambda} \quad (\text{effectively a.e.})$$

and

$$\frac{\nu(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \widetilde{\frac{d\nu}{d\lambda}}(x) \quad (\text{on Schnorr random } x).$$

Proof. By the effective decompositions (Propositions 9.10 and 9.11) decompose ν into $\nu = \nu_{ac}^+ + \nu_{ac}^- + \nu_s^+ + \nu_s^-$ (the order of the decompositions does not matter). Then

$$\frac{\nu_s^+(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} 0 \quad \text{and} \quad \frac{\nu_s^-(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} 0 \quad (\lambda\text{-a.e.})$$

and

$$\frac{\nu_{ac}^+(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \left[\frac{d\nu}{d\lambda} \right]^+ \quad \text{and} \quad \frac{\nu_{ac}^-(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} \left[\frac{d\nu}{d\lambda} \right]^- \quad (\lambda\text{-a.e.}).$$

Apply Lemma 9.6 and Corollary 4.20 respectively. \square

Remark 9.13. An alternate proof would be to prove the following stronger version of Fact 9.5. Since signed measures form a vector space, denote $a \cdot \nu$ for the signed measure given by scaling ν by $a \in \mathbb{R}$. Also by $|\nu|$ we mean the positive measure $\nu^+ + \nu^-$. One can show that

$$\frac{|\nu - \frac{d\nu}{d\lambda}(x) \cdot \lambda|(B(x, r))}{\lambda(B(x, r))} \xrightarrow{r \rightarrow 0} 0 \quad (\lambda\text{-a.e.}).$$

We could decompose this effectively into $|\nu - \frac{d\nu}{d\lambda}(x) \cdot \lambda| = |\nu_{ac} - \frac{d\nu}{d\lambda}(x) \cdot \lambda| + \nu_s^+ + \nu_s^-$. The first term can be handled by the same proof as the effective Lebesgue differentiation theorem (Theorem 4.10), and the last terms can be handled using Lemma 9.6.

In Section 12, I give some examples which show the theorems of this section characterize Schnorr randomness.

9.2. Functions of bounded variation. A function $f: [0, 1] \rightarrow \mathbb{R}$ is of BOUNDED VARIATION if there is some bound b such that for all finite sequences $0 = a_0 \leq a_1 \leq \dots \leq a_k = 1$ we have

$$\sum_{i < k} |f(a_{i+1}) - f(a_i)| \leq b.$$

The smallest such b is the TOTAL VARIATION (NORM) of f and is written $V(f)$. We have the following fact.

Fact 9.14 (See [13]). *Every function on $[0, 1]$ of bounded variation is differentiable almost-everywhere, and the derivative is integrable.*

Since every absolutely continuous function is of bounded variation, Fact 9.14 implies Fact 4.16.

There are a number of approaches to represent functions of bounded variation and their differentiability using computable analysis. The simplest approach is to only consider computable functions of bounded variation [8]. However, not all bounded variation functions are continuous.

The most general approach is to consider functions defined on a computably enumerable, countable, dense subset of $[0, 1]$. Then instead of differentiability we will consider pseudo-differentiability. This approach has been used in both constructive mathematics [5, 10] and computable analysis [32, 8, 29].

Definition 9.15. Let $\{a_n\}_{n \in \mathbb{N}}$ be a uniformly computable dense sequence of distinct reals in $[0, 1]$ with $a_0 = 0$ and $a_1 = 1$. Let $f: \{a_n\}_{n \in \mathbb{N}} \rightarrow \mathbb{R}$ be a function. We say f is COMPUTABLE if $f(a_n)$ is uniformly computable from n . Define the TOTAL VARIATION of f as follows where the supremum is over finite sequences $a_{n_0} < \dots < a_{n_k}$ in $\{a_n\}_{n \in \mathbb{N}}$.

$$V(f) = \sup_{a_{n_0} < \dots < a_{n_k}} \sum_{i < k} |f(a_{i+1}) - f(a_i)|$$

Let $x \in (0, 1)$. Then define the PSEUDO-DERIVATIVE of f at x as

$$(9.1) \quad \check{D}f(x) = \lim_{|b-a| \rightarrow 0} \frac{f(b) - f(a)}{b - a}$$

where the limit is over all $a, b \in \{a_n\}_{n \in \mathbb{N}}$ such that $a < x < b$. Say f is PSEUDO-DIFFERENTIABLE at x if the limit converges.

Proposition 9.16. *All functions f as in Definition 9.15 such that $V(f) < \infty$ are pseudo-differentiable for a.e. $x \in (0, 1)$, and the derivative is an integrable function.*

Proof. Just extend f to a total bounded variation function g by setting

$$g(x) = \begin{cases} f(x) & x \in \{a_n\}_{n \in \mathbb{N}} \\ \lim_{a \rightarrow x^-} f(x) & (a \in \{a_n\}_{n \in \mathbb{N}}) \text{ otherwise} \end{cases}.$$

(This limit exists since $V(f) < \infty$.) Then apply Fact 9.14. \square

Consider these examples of functions of bounded variation.

Example 9.17. Assume $g: [0, 1] \rightarrow \mathbb{R}$ is a computable (and hence continuous) function of bounded variation. Assume $\{a_n\}_{n \in \mathbb{N}}$ is as in Definition 9.15. Let $f = g \upharpoonright \{a_n\}_{n \in \mathbb{N}}$ (i.e. the restriction of g to $\{a_n\}_{n \in \mathbb{N}}$). Then $f: \{a_n\}_{n \in \mathbb{N}} \rightarrow \mathbb{R}$ is computable (as in Definition 9.15) and of bounded variation. Moreover, $V(f) = V(g)$ and the derivative $\frac{d}{dx}g$ is equal to $\check{D}f$ for all $x \in (0, 1)$.

Conversely, assume $f: \{a_n\}_{n \in \mathbb{N}} \rightarrow \mathbb{R}$ is a computable function of bounded variation with a continuous extension g and that $V(f)$ is computable. Assume that f can be extended to a continuous function $g: [0, 1] \rightarrow \mathbb{R}$ (i.e. $f = g \upharpoonright \{a_n\}_{n \in \mathbb{N}}$). Then g is a computable function (uniformly computable from $V(f)$ and f). (The modulus of continuity is computable from the variation. See Lu and Weihrauch [32].)

Remark 9.18. One could also consider L^1 -computable functions of bounded variation, as well as functions of the form $f(x) = \nu([0, x])$ for some computable signed measure ν . However, it requires some care to work with these types of functions and I will not do so here.

Theorem 9.19. *Let $f: \{a_n\}_{n \in \mathbb{N}} \rightarrow \mathbb{R}$ be computable (as in Definition 9.15). Assume $V(f)$ is computable (and hence finite) and the derivative $F := \check{D}f$ is L^1 -computable. Then f is pseudo-differentiable on all Schnorr randoms. Further*

$$\frac{f(b) - f(a)}{b - a} \xrightarrow{(b-a) \rightarrow 0} F \quad (a, b \in \{a_n\}_{n \in \mathbb{N}}, a < x < b).$$

converges effectively a.e., and $\check{D}f(x) = \tilde{F}(x)$ on Schnorr randoms x .

Proof sketch. Follow the arguments of Section 9.1. Replace the norm $\|\nu\|_{TV}$ with the total variation norm $V(f)$; positive measures with increasing functions; the Radon-Nikodym derivative with the pseudo-derivative; absolutely continuous measures with absolutely continuous functions; singular measures with functions of derivative zero; and the Lebesgue/Jordan decompositions with their corresponding versions for functions of bounded variation. See [29] for an effective version of the Jordan decomposition for functions of bounded variation. \square

Corollary 9.20. *Let $g: [0, 1] \rightarrow \mathbb{R}$ be a computable function of bounded variation. Assume $V(g)$ is computable and the derivative $G := \frac{d}{dx}g$ is L^1 -computable. Then g is differentiable on all Schnorr randoms. Further the derivative converges effectively a.e. to G , and $\frac{d}{dx}g|_{x=z} = \tilde{G}(z)$ on Schnorr randoms z .*

Proof. Use Theorem 9.19 and Example 9.17. \square

In Section 12, I give some examples which show the theorems of this section characterize Schnorr randomness.

10. THE ERGODIC THEOREM

There has been a great deal of interest in the effectivity of the ergodic theorems, both in terms of rates of convergence and randomness. In this section, I briefly summarize the results for Schnorr randomness.

Fact 10.1 (See [51]). *Let (\mathbb{X}, μ, T) be a measure-preserving system. Define $A_n f = \frac{1}{n} \sum_{i < n} f \circ T^i$. Let $\mathcal{I}nv(T)$ be the σ -algebra of T -invariant sets. Then $A_n f \rightarrow f^* := \mathbb{E}[f \mid \mathcal{I}nv(T)]$ a.e. and in the L^1 -norm. If f is L^2 , then convergence is in the L^2 -norm as well. If $\mathcal{I}nv(T)$ is TRIVIAL ($\mathbb{E}[f \mid \mathcal{I}nv(T)] = \int f d\mu$ for all f), then the system is said to be ERGODIC and $A_n f \rightarrow f^* = \int f d\mu$.*

This next theorem is a combination of results from a number of authors. I use techniques from this paper to fill in a few gaps not explicitly in the literature.

Theorem 10.2. *Let (\mathbb{X}, μ, T) be a measure preserving system where (\mathbb{X}, μ) is a computable probability space and $T: (\mathbb{X}, \mu) \rightarrow (\mathbb{X}, \mu)$ is an effectively measurable measure-preserving map.*

- (1) *If f is L^1 -computable and the limit f^* is L^1 -computable, then $A_n f \rightarrow f^*$ both effectively in L^1 and effectively a.e. Hence $\widetilde{A_n f}(z) \rightarrow \widetilde{f^*}(z)$ on Schnorr randoms z . In particular, the system is ergodic or if $\mathbb{E}[\cdot | \text{Inv}(T)]$ is a computable operator on $L^1 \rightarrow L^1$, then f^* is L^1 -computable and the results in the preceding sentence hold.*
- (2) *If f is L^2 -computable and $\|f^*\|_{L^2}$ is computable, then f^* is L^2 -computable and $A_n f \rightarrow f^*$ effectively in the L^2 -norm, the L^1 -norm, and effectively a.e. Hence $\widetilde{A_n f}(z) \rightarrow \widetilde{f^*}(z)$ on Schnorr randoms z . In particular, if the system is ergodic or if $\mathbb{E}[\cdot | \text{Inv}(T)]$ is a computable operator on $L^2 \rightarrow L^2$, then $\|f^*\|_{L^2}$ is computable and the results in the preceding sentence hold.*

(Hoyrup [23] mentions that $\mathbb{E}[\cdot | \text{Inv}(T)]$ is a computable operator on $L^1 \rightarrow L^1$ if and only if the ergodic decomposition $x \mapsto \mu_x$ is effectively measurable (layerwise computable). The same is true of L^2 .)

Proof. The first sentence of (1) follows from Avigad, Gerhardy, Towsner [2] and Galatolo, Hoyrup, Rojas [22] in the case that the system is ergodic.¹⁵ The Galatolo et al. proof also holds in the non-ergodic case by replacing $\int f d\mu$ with the L^1 -computable limit f^* [personal communication with Hoyrup and Rojas]. The first sentence of (2) follows from Avigad, Gerhardy, Towsner [2].

The part about Schnorr randomness follows from Lemma 3.19 (see also [21, 40]). \square

A Martin-Löf random version of this next corollary can be found in Bienvenu, Day, Hoyrup, Mezhirov, and Shen [3]. The proof is the same. It is a generalization of Kučera's theorem.

Corollary 10.3. *Assume T is an effectively measurable, ergodic, measure preserving action on (\mathbb{X}, μ) and A is an effectively measurable set. Then for all Schnorr randoms x , there are infinitely-many k such that $\widetilde{T}^k(x) \in A$.*

Proof. By Theorem 10.2, $\frac{1}{n} \sum_{k < n} \mathbf{1}_A(\widetilde{T}^k(x)) \rightarrow \mu(A) > 0$. Hence, there are infinitely many k such that $\widetilde{T}^k(x) \in A$. \square

Corollary 10.4 (Kučera's theorem for Schnorr randomness). *If $C \subseteq 2^{\mathbb{N}}$ is a closed set of positive measure and $x \in 2^{\mathbb{N}}$ is Schnorr random, then some tail of x is in C .*

Proof. In the previous result, let T be the left shift map ($T(0x) = T(1x) = x$) and let $A = C$. \square

11. BACKWARDS MARTINGALES AND THEIR APPLICATIONS

In this section, I discuss backwards martingales. Unlike “forward martingales” and ergodic averages, backwards martingales have not before been used before in algorithmic randomness. However, like forward martingales and ergodic averages, they are a powerful tool.

The definition of martingale can be extended to any linearly ordered (or partially ordered) index set I . Namely, $(\mathcal{F}_i)_{i \in I}$ is a FILTRATION if $\mathcal{F}_i \subseteq \mathcal{F}_j$ for any $i \leq j$, and $(M_i)_{i \in I}$ is a MARTINGALE adapted to $(\mathcal{F}_i)_{i \in I}$ if each M_i is \mathcal{F}_i -measurable and $\mathbb{E}[M_j | \mathcal{F}_i] = M_i$ for any $i \leq j$. If the index set I is the nonpositive integers, then we say M is a BACKWARDS (or REVERSE) MARTINGALE, often written (M_{-k}) to denote that the martingale is backwards. As opposed to “forward martingales”, backwards martingales always converge a.e. and in the L^1 -norm.

Fact 11.1 (See [14]). *Let (M_{-k}) be a backwards martingale adapted to the filtration (\mathcal{F}_{-k}) and let $\mathcal{F}_{-\infty} = \bigcap_k \mathcal{F}_{-k}$. Then $M_{-k} \rightarrow M_{-\infty} = \mathbb{E}[M_{-0} | \mathcal{F}_{-\infty}]$ both in L^1 and a.e.*

We have the following analog of Theorem 6.5 and Corollary 6.8.

Theorem 11.2. *Fix a computable probability space (\mathbb{X}, μ) .*

- (1) *If (M_{-k}) is an L^1 -computable backwards martingale, and the limit $M_{-\infty}$ is L^1 -computable, then $M_{-k} \rightarrow M_{-\infty}$ converges effectively in the L^1 -norm and effectively a.e. Hence, $\widetilde{M_{-k}}(z) \rightarrow \widetilde{M_{-\infty}}(z)$ on Schnorr randoms z .*

¹⁵For Avigad et al. the measure preserving map T is “computable” if the corresponding operator $f \mapsto f \circ T$ is a computable from L^2 to L^2 . By Proposition 3.30, this is the same as effectively measurable.

While the Galatolo et al. result is for a.e. computable T , the proof works for effectively measurable T by the fact that if f is L^1 - or L^2 -computable, then so is $f \circ T$ (uniformly from f and T) (Proposition 3.29).

- (2) If (M_{-k}) is an L^2 -computable backwards martingale, and $\|M_{-\infty}\|_{L^2} = \inf_k \|M_{-k}\|_{L^2}$ is computable, then $M_{-k} \rightarrow M_{-\infty}$ converges effectively in the L^2 -norm and effectively a.e. Hence, $M_{-\infty}$ is L^2 -computable, and $\widetilde{M}_{-k}(z) \rightarrow \widetilde{M}_{-\infty}(z)$ on Schnorr randoms z .

Proof. In the L^1 case, the proof is basically the same as that of Lemma 6.5. Since the limit is known, there is an effectively convergent subsequence. Further, since the inequalities (6.1) and (6.2) only apply to finite intervals of indices, they remain true for backwards martingales.

In the L^2 -case the argument resembles Corollary 6.8. For any $k \in \mathbb{N}$, $M_{-\infty} = \mathbb{E}[M_{-k} \mid \mathcal{F}_{-\infty}]$. By the Pythagorean theorem,

$$\|M_{-k} - M_{-\infty}\|_{L^2}^2 = \|M_{-k}\|_{L^2}^2 - \|M_{-\infty}\|_{L^2}^2.$$

So $M_{-k} \rightarrow M_{-\infty}$ effectively in L^2 . The rest follows from the L^1 -case. \square

Remark 11.3. Theorem 11.2 is analogous to both the effective ergodic theorem (Theorem 10.2) and the effective Lévy 0-1 law (Theorem 6.3), as seen in Table 3. Note that all three theorems can be viewed as taking place on structured probability spaces. The ergodic theorems takes place in a measure preserving system (\mathbb{X}, μ, T) with T effectively measurable, and the martingale theorems take place in a FILTERED PROBABILITY SPACE $(\mathbb{X}, \mu, (\mathcal{F}_n))$, where (\mathcal{F}_n) is a computable filtration (Remark 7.7). In such a computable system, a.e. convergence is computable when $f \mapsto \mathbb{E}[f \mid \mathcal{G}]$ is a computable operator for the limit σ -algebra \mathcal{G} .

	Ergodic averages	Backwards martingales	Lévy 0-1 law
Space	(\mathbb{X}, μ, T)	$(\mathbb{X}, \mu, (\mathcal{F}_{-n}))$	$(\mathbb{X}, \mu, (\mathcal{F}_n))$
Averages	$\frac{1}{n} \sum_{k < n} f \circ T^k$	$\mathbb{E}[f \mid \mathcal{F}_{-n}]$	$\mathbb{E}[f \mid \mathcal{F}_n]$
Limit	$\mathbb{E}[f \mid \text{Inv}(T)]$	$\mathbb{E}[f \mid \mathcal{F}_{-\infty}]$	$\mathbb{E}[f \mid \mathcal{F}_{\infty}]$
Limit σ -algebra	$\text{Inv}(T)$	$\mathcal{F}_{-\infty}$	\mathcal{F}_{∞}
“Nicest” system	ergodic system	$\mathcal{F}_{-\infty}$ is trivial	\mathcal{F}_{∞} is Borel σ -alg.

TABLE 3. Comparison of three convergence theorems.

Backwards martingales are quite useful. I will give three applications. The first application is a variation of Kučera’s theorem for Schnorr randomness (Corollary 10.4). However, this version does not follow directly from the ergodic theoretic Corollary 10.3.

Corollary 11.4. *On $(2^{\mathbb{N}}, \lambda)$, assume A is effectively measurable and $\lambda(A) > 0$. Then for all Schnorr random $x \in 2^{\mathbb{N}}$, there is some Schnorr random $y \in \widetilde{A}$ such that y is a permutation of finitely-many bits of x . In particular, if A is an effectively closed set of computable positive measure, then $y \in A$.*

Proof. Let \mathcal{F}_{-n} be the sigma-algebra of sets invariant under permutations of the first n bits. Notice that $\mathcal{F}_{-0} \supseteq \mathcal{F}_{-1} \supseteq \dots$ and that $\mathbb{E}[\mathbf{1}_A \mid \mathcal{F}_{-n}] = \frac{1}{n!} \sum_T \mathbf{1}_A \circ T$ where $T: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ ranges over permutations of bits which permute only the first n bits. Hence $\mathbb{E}[\mathbf{1}_A \mid \mathcal{F}_{-n}]$ is an L^1 -computable backwards martingale. Further, $\mathcal{F}_{-\infty} = \bigcap_n \mathcal{F}_{-n}$ is trivial (i.e. all sets in \mathcal{F}_{-n} are measure one or measure zero). Let x be Schnorr random. By Theorem 11.2,

$$\frac{1}{n!} \sum_T \mathbf{1}_{\widetilde{A}}(T(x)) = \mathbb{E}[\mathbf{1}_A \mid \mathcal{F}_{-n}](x) \xrightarrow{n \rightarrow \infty} \mathbb{E}[\mathbf{1}_A \mid \mathcal{F}_{-\infty}](x) = \lambda(A) > 0.$$

Therefore, $\frac{1}{n!} \sum_T \mathbf{1}_{\widetilde{A}}(T(x)) > 0$ for some n , and, moreover, $T(x) \in \widetilde{A}$ for some T which permutes the first n bits. \square

Before giving the next two examples, recall the probabilistic mindset. For the remainder of this section we fix a computable probability space (Ω, \mathbb{P}) as our sample space. We are not concerned with what this space is. A measurable function $X: (\Omega, \mathbb{P}) \rightarrow \mathbb{R}$ is called a RANDOM VARIABLE. Recall its DISTRIBUTION (or PUSH-FORWARD PROBABILITY MEASURE) \mathbb{P}_X is a probability measure on \mathbb{R} defined by

$$(11.1) \quad \int \varphi d\mathbb{P}_X = \mathbb{E}[\varphi(X)]$$

for any bounded continuous $\varphi: \mathbb{R} \rightarrow \mathbb{R}$. (This equation then extends to all $\varphi \in L^1(\mathbb{R}, \mathbb{P}_X)$.) Given a sequence of random variables $X = (X_i)_{i \in \mathbb{N}}$, the joint distribution of X is the probability measure \mathbb{P}_X on $\mathbb{R}^{\mathbb{N}}$ given by the equation

$$\int \varphi d\mathbb{P}_X = \mathbb{E}[\varphi(X_0, \dots, X_{d-1})]$$

for any bounded continuous $\varphi: \mathbb{R}^{\mathbb{N}} \rightarrow \mathbb{R}$ depending only on the first d coordinates.. *In other words, one may just think of (Ω, \mathbb{P}) as $(\mathbb{R}^{\mathbb{N}}, \mathbb{P}_X)$. Then X_i just becomes the i th coordinate of $\mathbb{R}^{\mathbb{N}}$.*¹⁶ A sequence $X = (X_i)$ is INDEPENDENT AND IDENTICALLY DISTRIBUTED (I.I.D.) if the joint distribution \mathbb{P}_X is the product measure $\mu^{\mathbb{N}} := \mu \otimes \mu \otimes \dots$ where $\mu = \mathbb{P}_{X_0}$. Equivalently, for all bounded continuous functions $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$,

$$(11.2) \quad \mathbb{E}[\varphi(X_0, \dots, X_{d-1})] = \int \varphi d\mu^d.$$

Fact 11.5 (Strong law of large numbers, see [14]). *Let (X_i) be a sequence of i.i.d. integrable random variables with partial sums $S_k = \sum_{i=0}^{k-1} X_i$. Then $S_k/k \rightarrow \mathbb{E}[X_0]$ a.e. (and in the L^1 -norm).*

Corollary 11.6 (Effective strong law of large numbers). *Let (X_i) be sequence of i.i.d. L^1 -computable random variables with partial sums $S_k = \sum_{i=0}^{k-1} X_i$. Then $S_k/k \rightarrow \mathbb{E}[X_0]$ effectively a.e. and effectively in the L^1 -norm. Hence, $\tilde{S}_k(\omega)/k \rightarrow \mathbb{E}[X_0]$ on Schnorr randoms ω .*

Proof. It is known that $M_{-k} := S_k/k$ is a backwards martingale adapted to the filtration $\mathcal{F}_{-k} = \sigma(S_k, S_{k+1}, \dots) = \sigma(S_k, X_{k+1}, X_{k+2}, \dots)$ [14, Example 5.61]. (\mathcal{F}_{-k} is the σ -algebra of sets invariant under permuting X_0, \dots, X_{k-1} .) Clearly (M_{-k}) is L^1 -computable. By the strong law of large numbers, we know (M_{-k}) converges to $\mathbb{E}[X_0]$, and the expectation is a computable real number. Hence by Theorem 11.2 $S_k/k \rightarrow \mathbb{E}[X_0]$ effectively in the L^1 -norm and effectively a.e. Hence, by Lemma 3.19, $\tilde{S}_k/k \rightarrow \mathbb{E}[X_0]$ on Schnorr randoms. \square

Remark 11.7. Taking $(\Omega, P) = (2^{\mathbb{N}}, \lambda)$ and $X_i(x) = x(i)$, the previous corollary implies that all Schnorr randoms z have an equal density of 1s and 0s—a fact which is well known. In Section 12, I use an extension of this fact to show that the strong law of large numbers characterizes Schnorr randomness. Corollary 11.6 could also be proved using the effective ergodic theorem (Theorem 10.2). Indeed, this is another similarity between backwards martingales and ergodic averages.

Now, I consider de Finetti's theorem. A sequence of random variables $X = (X_i)$ is EXCHANGEABLE if the joint distribution of (X_0, \dots, X_{d-1}) is the same as that of $(X_{\sigma(0)}, \dots, X_{\sigma(d-1)})$ for any permutation σ . In other words, the joint distribution \mathbb{P}_X is unchanged by permuting coordinates. De Finetti's theorem says that every exchangeable sequence is a convex combination of i.i.d. sequences.

Fact 11.8 (de Finetti's theorem, see [14, 19]). *Every exchangeable sequence of random variables $X = (X_i)$ is i.i.d. conditioned on some random measure μ . That is there is a (Ω, \mathbb{P}) -measurable random map $\mu: \omega \mapsto \mu_\omega$ where μ_ω is a probability measure on \mathbb{R} , such that for any bounded continuous function $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$,*

$$(11.3) \quad \mathbb{E}[\varphi(X_0, \dots, X_{d-1}) \mid \mu](\omega) = \int \varphi d\mu_\omega^d \quad (\mathbb{P}\text{-a.e. } \omega)$$

where $\mathbb{E}[\cdot \mid \mu]$ is conditioning on the least σ -algebra for which the map $\omega \mapsto \mu_\omega$ is measurable. This random measure $\omega \mapsto \mu_\omega$, called the DIRECTING MEASURE, is \mathbb{P} -a.s. unique.

Moreover, the following a.e. convergence theorems hold. For every $f \in L^1(\mathbb{R}^d, \mathbb{P}_{X_0})$,

$$(11.4) \quad \frac{1}{k} \sum_{i=0}^{k-1} f(X_i(\omega)) \rightarrow \mathbb{E}[f(X_0) \mid \mu](\omega) \quad (\mathbb{P}\text{-a.e. } \omega)$$

This can be extended to all $f \in L^1(\mathbb{R}^d, \mathbb{P}_{X_0, \dots, X_{d-1}})$ as follows.

$$(11.5) \quad A_k(f) = \frac{1}{k!/(k-d)!} \sum_{\sigma} f(X_{\sigma(0)}, \dots, X_{\sigma(d-1)}) \rightarrow \mathbb{E}[f(X_0, \dots, X_{d-1}) \mid \mu](\omega) \quad (\mathbb{P}\text{-a.e. } \omega)$$

where the average is over all $\frac{k!}{(k-d)!}$ many injections $\sigma: \{0, \dots, d-1\} \rightarrow \{0, \dots, k-1\}$.

¹⁶This intuition also holds in computable probability. A probability measure μ on $\mathbb{R}^{\mathbb{N}}$ is computable if and only if there is a sequence $X = (X_i)$ of uniformly effectively measurable (even a.e. computable) random variables on $(2^{\mathbb{N}}, \lambda)$ such that $\mu = \mathbb{P}_X$ [45, 26].

First, note the connection with the strong law of large numbers. If $X = (X_i)$ is i.i.d., then $\omega \mapsto \mu_\omega$ is constant. Therefore, the strong law of large numbers follows from equation (11.4) using $f(x) = x$. Second, note the similarity between equations (11.5) and the ergodic theorem.

Theorem 11.9 (Computable de Finetti's theorem (Freer, Roy [19])). *If $X = (X_i)$ is a sequence of exchangeable random variables with computable distribution \mathbb{P}_X , then the distribution \mathbb{P}_μ of the directing measure μ is computable from \mathbb{P}_X and vice versa.*

We now can show this effective a.e. convergence theorem.

Corollary 11.10. *Let $X = (X_i)$ be a uniformly computable sequence of effectively measurable, exchangeable random variables with directing measure μ . Then for all $f \in L^1_{\text{comp}}(\mathbb{R}^d, \mathbb{P}_{X_0, \dots, X_{d-1}})$,*

$$A_k(f) \rightarrow \mathbb{E}[f(X_0, \dots, X_{d-1}) \mid \mu]$$

both effectively a.e. and effectively in L^1 . Hence, for all Schnorr random ω ,

$$\widetilde{A}_k(\widetilde{f})(\omega) \rightarrow \widetilde{\mathbb{E}}[f(X_0, \dots, X_{d-1}) \mid \mu](\omega)$$

where

$$\widetilde{A}_k(f) := \frac{1}{k!/(k-d)!} \sum_{\sigma} f(\widetilde{X}_{\sigma(0)}, \dots, \widetilde{X}_{\sigma(d-1)}).$$

Proof. Since $X = (X_i)$ is uniformly effectively measurable, the distribution \mathbb{P}_X is computable (Propositions 3.25 and 3.27). Then by Theorem 11.9, the distribution \mathbb{P}_μ is also computable. Also, for any $f \in L^1(\mathbb{R}^d, \mathbb{P}_{X_0, \dots, X_{d-1}})$, we have $M_{-k} = A_k(f)$ is a backwards martingale [14, Chapter 5].

Let $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$ be a bounded computable function. Then $M_{-k} = A_k(\varphi)$ is an L^2 -computable backwards martingale. By Theorem 11.2, it is enough to compute the (square of the) L^2 -norm of the limit

$$\begin{aligned} \|\mathbb{E}[\varphi(X_0, \dots, X_{d-1}) \mid \mu]\|_{L^2(\Omega, P)}^2 &\stackrel{\text{eq. (11.3)}}{=} \int \left(\int \varphi d\mu_\omega \right)^2 dP(\omega) \\ &\stackrel{\text{eq. (11.1)}}{=} \int \left(\int \varphi d\nu \right)^2 dP_\mu(\nu). \end{aligned}$$

This last integral is computable since $\nu \mapsto \left(\int \varphi d\nu \right)^2$ is a computable map.

Hence we have proved the result for bounded computable $\varphi: \mathbb{R}^d \rightarrow \mathbb{R}$. For $f \in L^1_{\text{comp}}(\mathbb{R}^d, P_{X_0, \dots, X_{d-1}})$, take some φ which approximates f in the L^1 -norm, then

$$\begin{aligned} &\|\mathbb{E}[\varphi(X_0, \dots, X_{d-1}) \mid \mu] - \mathbb{E}[f(X_0, \dots, X_{d-1}) \mid \mu]\|_{L^1(\Omega, P)} \\ &= \|\mathbb{E}[(\varphi - f)(X_0, \dots, X_{d-1}) \mid \mu]\|_{L^1(\Omega, P)} \\ &\leq \|(\varphi - f)(X_0, \dots, X_{d-1})\|_{L^1(\Omega, P)} \\ &= \|\varphi - f\|_{L^1(\mathbb{R}^d, P_{X_0, \dots, X_{d-1}})}. \end{aligned}$$

Since the last term is uniformly computable, we can compute the limit $\mathbb{E}[f(X_0, \dots, X_{d-1}) \mid \mu]$ in the L^1 -norm. By Theorem 11.2, this completes the proof. \square

Example 11.11 (Pólya's urn). Consider an urn with one black ball and one red ball. At each stage k we take a ball from the urn, then return that ball to the urn along with another ball of the same color. Let X_k be the color of the k th ball drawn (0 for red, 1 for black). It turns out the sequence of random variables (X_k) is exchangeable. Let $S_k = \sum_{i=0}^{k-1} X_i$. By de Finetti's theorem the average S_k/k converges a.s., meaning that the ratio of red balls to black balls approaches a limit a.s. Now suppose, Pólya's Urn is modeled on a computer such that the random variables (X_k) are a.e. computable with respect to a uniformly distributed random real $x \in [0, 1]$. Then if x is Schnorr random, the simulation of Pólya's urn is guaranteed to converge to a fixed ratio of red and black balls.

Remark 11.12. There are other computable aspects of the ergodic theorem that could be explored for de Finetti's theorem. For one, the map $\omega \mapsto \mu_\omega$ is a form of ergodic decomposition. Hoyrup [23] has a number of results about the computability of the ergodic decomposition. In particular, I suspect that the map $x \mapsto \mu_x$ is effectively measurable. I also suspect Schnorr random points ω satisfy the following "typicalness"

property (similar to [21]) for de Finetti's theorem: for all bounded continuous (not necessarily computable) functions $\varphi: \mathbb{R}^d \rightarrow [0, 1]$, we have

$$\lim_{k \rightarrow \infty} \widetilde{A}_k(\varphi)(\omega) = \int \varphi d\tilde{\mu}_\omega^d.$$

Pursuing this, however, would take me too far afield.

12. CHARACTERIZING SCHNORR RANDOMNESS

In this section, I show that most of the effective a.e. convergence theorems in this paper are optimal in that Schnorr randomness cannot be strengthened to another form of randomness. In other words, combined with the effective a.e. convergence theorems in this paper, these examples characterize Schnorr randomness. See Table 1 in the introduction for how to match these examples to the corresponding a.e. convergence theorem(s).

12.1. Monotone convergence, the Lebesgue differentiation theorem, absolutely continuous functions and measures, and uniformly integrable martingales.

Example 12.1. Fix (\mathbb{X}, μ) and let (U_n) be a Schnorr test. Consider the following function f . By Remark 2.10, we may assume (U_n) is decreasing, and also assume $\mu(U_n) \leq 2^{-2n}$ by taking a subsequence. Let $f = \sum_n \mathbf{1}_{U_n}$. The following calculation shows that $f \in L_{comp}^2$.

$$\begin{aligned} \left\| f - \sum_{n=0}^{m-1} \mathbf{1}_{U_n} \right\|_{L^2} &= \left\| \sum_{n=m}^{\infty} \mathbf{1}_{U_n} \right\|_{L^2} \leq \sum_{n=m}^{\infty} \|\mathbf{1}_{U_n}\|_{L^2} = \sum_{n=m}^{\infty} \mu(U_n)^{1/2} \\ &\leq \sum_{n=m}^{\infty} 2^{-n} = 2^{-m+1} \end{aligned}$$

Clearly, $f(x) = \infty$ if x is covered by (U_n) .

This example is similar to the Schnorr integral tests of Miyabe [34]. This example will allow me to characterize Schnorr randomness using the monotone convergence theorem, the Lebesgue differentiation theorem, differentiation of absolutely continuous functions, differentiation of absolutely continuous measures, and convergence of uniformly integrable martingales.

Theorem 12.2 (Example of monotone convergence). *Let (U_n) be a Schnorr test on (\mathbb{X}, μ) . There is an increasing sequence of bounded computable functions (f_n) such that $\sup_n \|f_n\|_{L^2} = \infty$ and $\sup_n f_n(x) = \infty$ for all x covered by (U_n) .*

Proof. Let $f = \sum_n \mathbf{1}_{U_n}$ be as in Example 12.1. Define $g_n = \sum_{k < n} \mathbf{1}_{U_k}$. We can find a computable $f_n \leq g_n$ such that $\|g_n - f_n\|_{L^2} \leq 2^{-n}$ and $\sup_n f_n = \sup_n g_n = f$. Namely, by effective inner regularity (Proposition 3.22) find a closed set $C_n \subseteq U_n$ of computable measure such that $\mu(U_n - C_n) \leq 2^{-(k+1)}$. Then, using the effective Tietze extension theorem [53] we can find a computable function $h_k \leq \mathbf{1}_{U_k}$ such that $h_k = 0$ on U_k^c and $h_k = 1$ on C_k . Then $f_n = \sum_{k < n} h_k$ is as desired. \square

Theorem 12.3 (Example of Lebesgue differentiation theorem). *For any Schnorr test (U_i) on $([0, 1]^d, \lambda)$, there is an $f \in L_{comp}^2([0, 1]^d, \lambda)$ such that $\frac{1}{\lambda(B(x, r))} \int_{B(x, r)} f d\lambda \rightarrow \infty$ for all x covered by (U_i) . (This holds as well for \mathbb{T}^d and for the dyadic version on $2^{\mathbb{N}}$.)*

Proof. Take the L^2 -computable f from Example 12.1. Let x be covered by (U_i) . Then for each k , there is some r_k such that $B(x, r_k) \subseteq U_k$. Since (U_k) is decreasing, $f(y) \geq k$ for all $y \in B(x, r_k)$. Hence, $\frac{1}{\lambda(B(x, r))} \int_{B(x, r)} f d\lambda \geq k$. Hence $\limsup_{r \rightarrow 0} \frac{1}{\lambda(B(x, r))} \int_{B(x, r)} f d\lambda = \infty$. \square

Theorem 12.4 (Example of absolutely continuous measure). *Let (U_n) be a Schnorr test on $([0, 1]^d, \lambda)$. There is an absolutely continuous, positive measure μ with L^2 -computable derivative $\frac{d\mu}{d\lambda}$ such that $\frac{\mu(B(z, r))}{\lambda(B(z, r))} \xrightarrow{r \rightarrow 0} \infty$ for all z covered by (U_n) .*

Proof. Take the L^2 -computable f from Example 12.1. let μ be defined by $\mu(A) = \int_A f d\lambda$. The rest of the proof is the same as the previous one. \square

Theorem 12.5 (Example of absolutely continuous function). *Let (U_n) be a Schnorr test on $([0, 1], \lambda)$. There is an increasing, absolutely continuous, computable function F with L^2 -computable derivative such that $\frac{d}{dx}F|_{x=z} = \infty$ for all z covered by (U_n) .*

Proof. Take the L^2 -computable f from Example 12.1. Let $F(x) = \int_0^x f(t) dt$. Then F is computable, increasing, and absolutely continuous. By the same argument as in Theorem 12.3, $\frac{d}{dx}F|_{x=z} = \infty$ for all z covered by (U_n) . \square

Theorem 12.6 (Example of a dyadic uniformly integrable martingale). *Let (U_n) be a Schnorr test on $(2^{\mathbb{N}}, \lambda)$. There is a nonnegative, computable, dyadic, uniformly integrable, martingale (M_k) with limit $M_\infty \in L^2_{\text{comp}}$ (and hence in L^1_{comp}) such that $M_k(x) \rightarrow \infty$ on all x covered by (U_n) .*

Proof. Take the L^2 -computable f from Example 12.1. Then let $M_k = f^{(k)} = \mathbb{E}[f \mid \mathcal{B}_k]$ as in Example 6.1. This is a computable, dyadic martingale with limit $M_\infty = f$. If x is covered by (U_n) then $M_k(x) \rightarrow \infty$ by the same argument as Theorem 12.3. \square

In this next theorem, $x \in (\mathbb{X}, \mu)$ is KURTZ RANDOM if it is not in any Σ_2^0 null set. Every Kurtz random is Schnorr random. All a.e. computable functions f are defined on Kurtz randoms, since the domain of f is a measure one Π_2^0 set. Further, no Kurtz randoms are on the boundary of a ball in $\text{Basis}(\mathbb{X}, \mu)$, since the set of boundaries is a Σ_2^0 null set. Therefore for each decomposition of \mathbb{X} into finitely many cells, a Kurtz random x is in the interior of one of the cells. See Rute [42], for more discussion.

Theorem 12.7 (Example of a uniformly integrable martingale). *Fix (\mathbb{X}, μ) . Let (U_n) be a Schnorr test. From (U_n) we can construct an a.e. computable, uniformly integrable, L^2 -computable (and hence L^1 -computable) martingale (M_k) with limit $M_\infty \in L^2_{\text{comp}}$ (and hence in L^1_{comp}) such that $M_k(x)$ diverges for all Kurtz random x covered by U_n . (Since M_k is a.e. computable, it is well-defined on Kurtz randoms.)*

Proof. The idea is the same as the previous proof, except that one needs a “canonical filtration” for the space (\mathbb{X}, μ) . Recall the collection $\text{Basis}(\mathbb{X}, \mu)$ from Lemma 3.5 which has an enumeration $\{B_i\}$. Let \mathcal{P}_k be the partition generated by $\{B_0, \dots, B_k\}$. This generates a filtration $\sigma(\mathcal{P}_k)$ of \mathbb{X} such that $\sigma(\mathcal{P}_k) \uparrow \mathcal{B}(\mathbb{X})$ (the Borel σ -algebra).

Now let f be as in Example 12.1. Let $M_k = \mathbb{E}[f \mid \mathcal{P}_k]$ although we will define it in an a.e. computable manner as follows. To compute $M_k(x)$, just find the atom $Q \in \mathcal{P}_k$ that x is in, and then computing $\frac{1}{\mu(Q)} \int_Q f d\mu$. This can be done for almost every x , namely all x in the interior of some $Q \in \mathcal{P}_k$ with positive measure (all Kurtz randoms x have this property).

If x is Kurtz random and covered by (U_n) , then take the intersection of the first N many sets U_n that contain x . There is a ball $B(x, r)$ in the intersection (since we are assuming the U_n are decreasing). Since $\text{Basis}(\mathbb{X}, \mu)$ is an effective basis, there is a computable sequence of sets $\{Q_i\}$ from $\bigcup_k \mathcal{P}_k$ such that $B(x, r) = \bigcup_i Q_i$ μ -a.e. If x is Kurtz random, then $x \in Q$ for some $Q = Q_i \in \mathcal{P}_k$ for some k . Then we have for all $\ell \geq k$,

$$M_\ell(x) = \mathbb{E}[f \mid \mathcal{P}_\ell](x) \geq \mathbb{E}[N \cdot \mathbf{1}_Q \mid \mathcal{P}_\ell](x) = N \cdot \mathbf{1}_Q(x) = N.$$

Hence $M_k(x) \xrightarrow[k \rightarrow \infty]{} \infty$. \square

12.2. Singular martingales, functions of bounded variation, and measures. Consider these two examples of nonnegative, dyadic, singular martingales (the limit is zero) corresponding to a Schnorr test (U_n) . The main idea is to bet when it looks like x is in another U_n , and then to “bet away” the money back down to zero. One puts all its mass (bets all its money) on a countable set of points. The other puts its mass on a measure-zero set, without atoms.

Example 12.8 (Singular “atomic” martingale). Let (U_n) be a Schnorr test on $(2^{\mathbb{N}}, \lambda)$. Assume (U_n) is decreasing, and assume $\mu(U_n) \leq 2^{-n}$. Effectively partition $U_n = \bigcup_m [\sigma_m^n]$ (that is, a prefix-free representation of U_n). Let $(\sigma_i)_i$ be a reordering of $(\sigma_m^n)_{n,m}$. If x is covered by (U_n) then $x \in [\sigma_i]$ for infinitely-many i .

For each i , create a martingale as follows. For each i , let a_i be the “midpoint” of $[\sigma_i]$ (that is $a_i = \sigma_i 100\dots$). Let $b_i = \lambda(\sigma_i) / \sqrt{\lambda(U_n)}$ for the n such that $\sigma_i \subseteq U_n$. Then define a computable dyadic martingale $M^{(i)}$ as the one that puts all its money on the point a_i and has starting capital b_i . That is, for each $\tau \in 2^{<\omega}$, define

$$M^{(i)}(\tau) = \begin{cases} b_i / \lambda(\tau) & \text{if } a_i \in [\tau] \\ 0 & \text{otherwise} \end{cases}.$$

It is easy to verify each $M^{(i)}$ is a computable, nonnegative, singular, dyadic martingale. Define $M = \sum_i M^{(i)}$. This is also a nonnegative, singular, dyadic martingale, and M is finite and computable since $\sum_i b_i = \sum_n \lambda(U_n)/\sqrt{\lambda(U_n)} \leq \sum_n 2^{-n/2}$ and $\sum_i b_i$ is computable. If x is covered by (U_n) then for every n we have some i such that $x \in \sigma_i \subseteq U_n$ and $M(\sigma_i) \geq M^{(i)}(\sigma_i) = b_i/\lambda(\sigma_i) = 1/\sqrt{\lambda(U_n)} \geq 2^{n/2}$. Hence $\limsup_k M_k(x) = \infty$.

This first example allows us to characterize Schnorr randomness by singular martingales, atomic measures, and bounded variation functions consisting only of jumps.

Theorem 12.9 (Example of singular martingale). *Let (U_n) be a Schnorr test. There is a nonnegative, computable, singular, dyadic martingale (M_k) such that $\limsup_k M_k(x) = \infty$ for all x covered by U_n .*

Proof. Use the martingale in Example 12.8 (or in Example 12.13 below). \square

On $([0, 1]^d, \lambda)$ redefine $I_k(x)$ to be the OPEN DYADIC SET containing x (in the absolutely continuous case, it did not much matter if $I_k(x)$ was open or half-open). Define $\overline{I}_k(x)$ as the corresponding closed set.

Lemma 12.10. *Let μ be a computable positive measure on $2^{\mathbb{N}}$. There is a corresponding computable positive measure ν on $[0, 1]^d$ such that $\nu(I_k(x)) \leq \mu(x \upharpoonright dk) \leq \nu(\overline{I}_k(x))$ for all vectors x with no dyadic rational coordinates.*

Proof. Let $T: 2^{\mathbb{N}} \rightarrow [0, 1]^d$ be the (usual) computable map $T(x) = (y_0, \dots, y_{d-1})$ where $y_i = 0.x(i)x(d+i)x(2d+1)\dots$ in particular, $T^{-1}(I_k(x)) \subsetneq [x \upharpoonright dk] \subsetneq T^{-1}(\overline{I}_k(x))$. That is, the first d bits of x correspond to the first bit of each coordinate in (x_1, \dots, x_d) . Define ν as the push-forward measure of μ along T , hence $\nu(\overline{I}_k(x)) \leq \mu(x \upharpoonright dk) \leq \nu(I_k(x))$. By Proposition 3.25, ν is computable. \square

Theorem 12.11 (Example of atomic, singular measure). *Let (U_n) be a Schnorr test on $([0, 1]^d, \lambda)$. There is an atomic, singular positive measure ν such that $\limsup_r \frac{\nu(B(z, r))}{\lambda(B(z, r))} = \infty$ for all z covered by (U_n) .*

Proof. Let (V_n) be a test on $(2^{\mathbb{N}}, \lambda)$ which covers the points in $2^{\mathbb{N}}$ corresponding to the points that (U_n) covers in $[0, 1]^d$. (Partition each U_n into closed dyadic sets and replace each with the corresponding basic open set $[\sigma]$.)

Let μ be the computable positive measure on $2^{\mathbb{N}}$ associated with the martingale M in Example 12.8. That is, $\mu(\sigma) = M(\sigma)\lambda(\sigma)$. Notice that μ is atomic. Let ν be the computable positive measure on $[0, 1]^d$ as in Lemma 12.10; ν is still atomic. Without loss of generality, we assume in Example 12.8 that $\limsup_k M_{dk}(x) = \infty$ for the x covered by U_n . (Just require the σ_i to be of length dk for some k .) Then by Lemma 12.10, for all x covered by (U_n) . We have $\limsup_k \nu(\overline{I}_k(x))/\lambda(\overline{I}_k(x)) \geq \limsup_k M_k(x) = \infty$. By an geometric argument similar to the proof of Lemma 4.9, we have $\limsup_{r \rightarrow 0} \frac{\nu(B(z, r))}{\lambda(B(z, r))} = \infty$. \square

Theorem 12.12 (Example of bounded variation function with jumps). *Let (U_n) be a Schnorr test on $([0, 1], \lambda)$. There is a nondecreasing function F and a computable sequence of pairs of reals (a_i, b_i) such that $F(x) = \sum_{a_i \leq x} b_i$ (F only consists of jumps), $V(F) = \sum_i b_i$ is computable, and $\frac{d}{dx}F|_{x=z}$ does not exist for all z covered by (U_n) .*

Proof. Let a_i, b_i be from Example 12.8 (except a_i is now the corresponding real in $[0, 1]$). Let ν be the measure from the previous example. Notice that each a_i is an atom of ν with weight b_i . Hence, $F(x) = \sum_{a_i \leq x} b_i = \nu([0, x])$. By the previous proof, the derivative of F does not exist at z covered by (U_n) . \square

Now for the second example martingale.

Example 12.13 (Singular “continuous” martingale). Define σ_i and b_i the same as in Example 12.8. But now, we want to put the mass on a set of points in $[\sigma]$. Define $N^{(i)}$ as follows. If $|\tau| \leq |\sigma|$ then bet all the money on σ .

$$N^{(i)}(\tau) = \begin{cases} b_i/\lambda(\tau) & \text{if } \tau \preceq \sigma \\ 0 & \text{otherwise} \end{cases}.$$

If τ is incomparable with σ then $N^{(i)}(\tau) = 0$. If $\tau \succeq \sigma$, then bet that the even bits are all 1s, ignoring the odd bits. That is,

$$\begin{aligned} N^{(i)}(\tau 0) &= \begin{cases} 0 & |\tau| \text{ is even} \\ N^{(i)}(\tau) & |\tau| \text{ is odd} \end{cases} \\ N^{(i)}(\tau 1) &= \begin{cases} 2 \cdot N^{(i)}(\tau) & |\tau| \text{ is even} \\ N^{(i)}(\tau) & |\tau| \text{ is odd} \end{cases}. \end{aligned}$$

This nonnegative dyadic martingale will almost surely converge to 0 and is therefore singular. Define $M = \sum_i M^{(i)}$. As before, M is computable. If x is covered by (U_n) , by the same argument as in Example 12.8, we have $\limsup_k M_k(x) = \infty$.

Theorem 12.14. *Let (U_n) be a Schnorr test on $([0, 1]^d, \lambda)$. There is a continuous, singular, positive measure μ such that $\limsup_{r \rightarrow 0} \frac{\mu(B(z, r))}{\lambda(B(z, r))} = \infty$ for all z covered by (U_n) .*

Proof. Follow the proof of Theorem 12.11, except use Example 12.13 to get a continuous measure. \square

Theorem 12.15. *Let (U_n) be a Schnorr test on $([0, 1], \lambda)$. There is a continuous, nondecreasing function F with zero derivative almost surely such that $\frac{d}{dx} F|_{x=z}$ does not exist for all z covered by (U_n) .*

Proof. Let $F(x) = \nu([0, x])$ where ν is the measure in Theorem 12.14. Therefore, the derivative of F does not exist for all z covered by (U_n) . \square

12.3. Backwards martingales, the strong law of large numbers, de Finetti's theorem, and the ergodic theorem. Schnorr proved the following fact (which has been extended by Gács, Hoyrup, and Rojas), that shows each non-Schnorr random can fail to satisfy the law of large numbers.

Proposition 12.16 (Schnorr [44, Theorem 12.1]; Gács, Hoyrup, Rojas [21]). *If (U_n) is a Schnorr test, then there is an a.e. computable measure preserving transformation $\varphi: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ such that for all x not covered by (U_n) , if $y = \varphi(x)$, then $\limsup_k \frac{1}{k} \sum_{i < k} y(i) \geq \frac{2}{3}$.*

This fact will allow us to use the strong law of large numbers, de Finetti's theorem, backwards martingale convergence, and the ergodic theorem to characterize Schnorr randomness on $(2^{\mathbb{N}}, \lambda)$.

Corollary 12.17. *If (U_n) is a Schnorr test on $(2^{\mathbb{N}}, \lambda)$, then the following hold.*

- (1) *There is a computable i.i.d. sequence of i.i.d. 0, 1-valued random variables (X_i) such that $\frac{1}{k} \sum_{i < k} X_i(x)$ diverges for all x covered by (U_n) .*
- (2) *There is a computable exchangeable sequence of a.e. computable random variables (X_i) and a bounded computable $\psi: \mathbb{R} \rightarrow \mathbb{R}$, such that $\frac{1}{k} \sum_{i < k} \psi(X_i(x))$ diverges for all x covered by (U_n) .*
- (3) *There is a bounded a.e. computable backwards martingale (M_{-k}) with a constant, computable limit $M_{-\infty}$ such that $M_{-k}(x)$ diverges for all x covered by (U_n) .*
- (4) (Gács, Hoyrup, Rojas [21]) *There is a bounded a.e. computable function $f: 2^{\mathbb{N}} \rightarrow \mathbb{R}$ and an a.e. computable, ergodic, measure preserving $T: 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ such that $\frac{1}{k} \sum_{i < k} f(T^n(x))$ diverges for all x covered by (U_n) .*

Proof. Take φ as in Proposition 12.16. Slightly modify φ so that $\frac{1}{k} \sum_{i < k} \varphi(n)(i)$ diverges on all x covered by (U_n) . (Just swap the 0s and 1s whenever $\frac{1}{k} \sum_{i < k} \varphi(n)(i) > \frac{6}{5}$ or $< \frac{4}{5}$.)

For (1) and (2), let $X_n(x)$ be the n th bit of $\varphi(x)$. This sequence is i.i.d. (and therefore exchangeable). For (2), also let ψ be the identity map.

For (3), let $M_{-k} = \frac{1}{k} \sum_{j < k} X_k$ from (1). Recall from Corollary 11.6, this is a backwards martingale with limit $\frac{1}{2}$.

For (4), Gács, Hoyrup, and Rojas [21] showed that φ can be constructed to have an a.e. computable inverse. Set $T = \varphi \circ \sigma \circ \varphi^{-1}$ where σ is the left shift map, and set f to be the first bit of $\varphi(x)$. Then $\frac{1}{n} \sum_{k < n} f(T^n(x))$ is equal to $\frac{1}{k} \sum_{j < k} X_k$ which diverges. \square

12.4. Convergence of test functions to 0.

Theorem 12.18. *Let (U_n) be a Schnorr test in $(2^{\mathbb{N}}, \lambda)$. There is a computable sequence (φ_n) of dyadic test functions, such that $\|\varphi_n\|_{L^2} < 2^{-n}$ but $\limsup_n \varphi_n(x) = \infty$ on all x covered by (U_n) .*

Proof. By Remark 2.10, we may assume that $\lambda(U_n) \leq 2^{-(2n+2)}$. We may also computably break up (U_n) into a disjoint union of dyadic intervals $U_n = \bigcup_m [\sigma_m^n]$. (For each n , the set $\{\sigma_m^n\}$ be infinite or finite—it is enough to know it is computably enumerable uniformly in n .) Then

$$\sum_{n,m} \left(1/\sqrt{\lambda(U_n)}\right) \cdot \lambda(\sigma_m^n) = \sum_n \left(1/\sqrt{\lambda(U_n)}\right) \cdot \lambda(U_n) \leq 1,$$

and the sum is computable. Renumber $\{\sigma_i\}_i = \{\sigma_{n,m}\}_{n,m}$ using a computable pairing function. Effectively partition the double sequence $(\sigma_i)_i$ into finite sequences $(\sigma_{i(k)}, \sigma_{i(k)+1}, \dots, \sigma_{i(k+1)-1})$ such that $\sum_{j=\sigma(k)}^{i(k)-1} \left(1/\sqrt{\lambda(U_n)}\right) \cdot \lambda(\sigma_i) \leq 2^{-k}$ where i codes the pair (n, m) (break up the $[\sigma_i]$ into smaller intervals if needed).

Let $\varphi_k = \sum_{j=\sigma(k)}^{i(k)-1} (1/\sqrt{\lambda(U_n)}) \cdot \mathbf{1}_{[\sigma_i]}$. By the pigeonhole principle, if x is covered by (U_n) then for each n , $\varphi_k(x) > n$ for infinitely many k . \square

Theorem 12.19. *Let (U_n) be a Schnorr test on (\mathbb{X}, μ) . There is a computable sequence (φ_n) of test functions, such that $\|\varphi_n\|_{L^2} < 2^{-n}$ but $\limsup_n \varphi_n(x) = \infty$ on all Kurtz random x covered by (U_n) .*

Proof. The proof is the same as the previous one. Just replace dyadic intervals $[\sigma]$ with finite Boolean combinations of $\text{Basis}(\mathbb{X}, \mu)$ from Lemma 3.5. (Also, make the sets slightly larger to cover their measure-zero boundaries.) \square

Theorem 12.20. *Let (U_n) be a Schnorr test on (\mathbb{X}, μ) . There is a computable sequence (f_k) of computable functions such that $\|f_k\|_{L^2} < 2^{-k}$ but $\limsup_k f_k(x) = \infty$ on all x covered by (U_n) .*

Proof. Take the test functions (φ_k) from the previous two theorems. Approximate them with computable functions f_k as follows. For each $[\sigma]$ in φ_k (or the corresponding finite Boolean combination B of basis elements), find a computable function h_σ such that on $h_\sigma = 1$ on $[\sigma]$ and $\|h_\sigma - \mathbf{1}_\sigma\|_{L^2}$ is sufficiently small. This can be done by defining $h_\sigma = 1$ on $[\sigma]$ (or in the other case, on the closure \bar{B} which has the same measure), using effective outer regularity (Proposition 3.22) to find an open set $V \supseteq [\sigma]$ of similar measure, defining $h_\sigma = 0$ on V^c and then using the effective Tietze extension theorem [53] to extend this to a computable function. \square

Theorem 12.21. *Let (U_n) be a Schnorr test on $([0, 1]^d, \lambda)$. There is a computable sequence (p_k) of rational polynomials such that $\|p_k\|_{L^2} < 2^{-k}$ but $\limsup_k p_k(x) = \infty$ on all x covered by (U_n) .*

Proof. Take the computable functions (f_n) in the last theorem. Effectively approximate (f_n) by polynomials using the effective Weierstrass approximation theorem [41]. Since they are close in the uniform norm, they are close in the L^2 -norm. \square

APPENDIX A. PROOFS FROM SECTION 3.

A.1. Useful facts. The following set of calculations are straightforward, but useful.

Fact A.1. *If $f \leq g$ (a.e.), then*

$$\mu\{f > \varepsilon\} \leq \mu\{g > \varepsilon\}.$$

Also

$$\mu\{f_1 + f_2 > \varepsilon_1 + \varepsilon_2\} \leq \mu\{f_1 > \varepsilon_1\} + \mu\{f_2 > \varepsilon_2\}.$$

and

$$\mu\left\{\sum_i f_i > \sum_i \varepsilon_i\right\} \leq \sum_i \mu\{f_i > \varepsilon_i\}.$$

Also, recall Markov's inequality and a useful variation for the metric d_{meas} .

Fact A.2 (Markov's inequality, see [48]). Assume f is an integrable function and $\varepsilon > 0$. Then

$$\mu\{x \mid |f| \geq \varepsilon\} \leq \frac{\|f\|_{L^1}}{\varepsilon}.$$

Also given \mathbb{Y} -valued measurable functions f and g and $0 < \varepsilon \leq 1$,

$$\mu\{x \mid d_{\mathbb{Y}}(f, g) \geq \varepsilon\} = \mu\{x \mid \min\{d_{\mathbb{Y}}(f, g), 1\} \geq \varepsilon\} \leq \frac{d_{meas}(f, g)}{\varepsilon}.$$

A.2. Integrable functions, measurable functions, and measurable sets.

Restatement of Proposition 3.7. The measure of each cell of $\text{Basis}(\mathbb{X}, \mu)$ is computable from its code σ .


Proof. Given a cell $C = A_1 \cap \dots \cap A_\ell \cap B_1^c \cap \dots \cap B_k^c$ (where $A_1, \dots, A_\ell, B_1, \dots, B_k \in \text{Basis}(\mathbb{X}, \mu)$, that is balls of null boundary), then C is in between the effectively open and effectively closed sets $A_1 \cap \dots \cap A_\ell \cap \overline{B_1^c} \cap \dots \cap \overline{B_k^c}$ and $\overline{A_1} \cap \dots \cap \overline{A_\ell} \cap B_1^c \cap \dots \cap B_k^c$ which have the same measure. Since the measure of effectively open sets is lower semicomputable and closed sets is upper semicomputable (Proposition 2.5), the measure of C is computable (uniformly from its code σ). \square

Proposition A.3. Let A be a set formed by combining elements of $\text{Basis}(\mathbb{X}, \mu)$ using finitely-many connectives Boolean connectives $\cup, \cap, ^c$ as well as the closure operator. Then $\mu(A)$ is computable from (the code for) A .

Proof. A finite Boolean combination can be decomposed into a finite union of pairwise disjoint cells (basically disjunctive normal form). Since the boundaries of the cells have measure zero, the closure operator does not effect the measure. \square

A.3. Effective modes of convergence.

Restatement of Proposition 3.15 (Modes of effective convergence). On a computable probability space (\mathbb{X}, μ) , the following implications are effective—in that a rate of convergence for the latter is computable from the former. (L^1 and L^2 only apply to real-valued functions.)



(1) The dotted arrow represents that if $f_i \rightarrow f$ with a geometric rate of convergence in the metric d_{meas} , e.g. $\forall j \geq i \ d_{meas}(f_j, f) \leq 2^{-i}$, then $f_i \rightarrow f$ effectively almost uniformly.

(2) For the arrow going to “Schnorr”, see Lemma 3.19.

Proof. ($L^2 \rightarrow L^1 \rightarrow d_{meas}$): Use that $d_{meas}(f_i, f) \leq \|f_i - f\|_{L^1} \leq \|f_i - f\|_{L^2}$.

($d_{meas} \rightarrow \text{measure}$): Assume $n(\varepsilon)$ is a rate of convergence in the metric d_{meas} . I claim $m(\varepsilon_1, \varepsilon_2) = n(\varepsilon_1 \varepsilon_2)$ is a rate of convergence in measure (assuming $0 < \varepsilon < 1$). Indeed, for $i \geq n(\varepsilon_1 \varepsilon_2)$, $d_{meas}(f_i, f) \leq \varepsilon_1 \varepsilon_2$ and by Markov's inequality (Fact A.2),

$$\mu\{d_{\mathbb{Y}}(f_i, f) > \varepsilon_1\} \leq \frac{d_{meas}(f_i, f)}{\varepsilon_1} \leq \frac{\varepsilon_1 \varepsilon_2}{\varepsilon_1} = \varepsilon_2.$$

(Measure $\rightarrow d_{meas}$): Let $m(\varepsilon_1, \varepsilon_2)$ be a rate of convergence in measure. I claim that $n(\varepsilon) = m(\varepsilon/2, \varepsilon/2)$ is a rate of convergence in the metric d_{meas} . Indeed, for $i \geq m(\varepsilon/2, \varepsilon/2)$ we have that

$$d_{meas}(f_i, f) = \int \min\{d_{\mathbb{Y}}(f_i, f), 1\} d\mu \leq \mu\{d_{\mathbb{Y}}(f_i, f) > \varepsilon/2\} + \varepsilon/2 \leq \varepsilon/2 + \varepsilon/2.$$

(Almost uniform $\rightarrow \text{measure}$): A rate of effective almost uniform convergence $n(\varepsilon_1, \varepsilon_2)$ is also a rate of convergence in measure since if $i \geq n(\varepsilon_1, \varepsilon_2)$,

$$\mu\{d_{\mathbb{Y}}(f_i, f) > \varepsilon_1\} \leq \mu\left\{\sup_{i \geq n(\varepsilon_1, \varepsilon_2)} d_{\mathbb{Y}}(f_i, f) > \varepsilon_1\right\}.$$

(1): Assume $\forall j \geq i$ $d_{\text{meas}}(f_j, f) \leq 2^{-i}$. Based on this rate of convergence in the metric d_{meas} , the rate of convergence in measure is $n(\varepsilon_1, \varepsilon_2) \geq -\log_2(\varepsilon_1 \varepsilon_2)$. I claim $n(\varepsilon_1, \varepsilon_2) = -\log_2\left(\frac{\varepsilon_1 \varepsilon_2}{(2+\sqrt{2})^2}\right)$ is a rate of almost uniform converge. Indeed, if $n = -\log_2\left(\frac{\varepsilon_1 \varepsilon_2}{(2+\sqrt{2})^2}\right)$, by Facts A.1 and A.2 and the fact that $\sum_{i \geq 0} 2^{-i/2} = 2 + \sqrt{2}$ we have

$$\begin{aligned} \mu \left\{ \sup_{i \geq n} d_{\mathbb{Y}}(f_i, f) > \varepsilon_1 \right\} &\leq \mu \left\{ \sum_{i \geq n} d_{\mathbb{Y}}(f_i, f) > \varepsilon_1 \right\} \\ &\leq \sum_{i \geq 0} \mu \left\{ d_{\mathbb{Y}}(f_{i+n}, f) > \frac{2^{-i/2}}{2+\sqrt{2}} \cdot \varepsilon_1 \right\} \\ &\leq \sum_{i \geq 0} \frac{d_{\text{meas}}(f_{i+n}, f)}{\frac{2^{-i/2}}{2+\sqrt{2}} \cdot \varepsilon_1} \leq \sum_{i \geq 0} \frac{2^{-(i+n)}}{\frac{2^{-i/2}}{2+\sqrt{2}} \cdot \varepsilon_1} \\ &= \frac{(2+\sqrt{2})}{2^n \varepsilon_1} \sum_{i \geq 0} 2^{-i/2} = \frac{(2+\sqrt{2})^2}{2^n \varepsilon_1} = \varepsilon_2. \quad \square \end{aligned}$$

Restatement of Proposition 3.16. *Let (f_n) and f be uniformly effectively measurable real-valued functions.*

- (1) *If $f_n \rightarrow f$ effectively a.e.. and $g_n \rightarrow g$ effectively a.e., then $f_n + g_n \rightarrow f + g$ effectively a.e..*
- (2) *If $f_n^j \rightarrow f^j$ effectively a.e.. ($j \in \{0, \dots, k-1\}$), and g is computable with a uniform modulus of continuity, then $g(f_n^0, \dots, f_n^{k-1}) \rightarrow g(f^0, \dots, f^{k-1})$ effectively a.e..*
- (3) *(Squeeze theorem) Assume $f_n \leq g_n \leq h_n$ a.e. and that $f_n \rightarrow g$ effectively a.e.. and $h_n \rightarrow g$ effectively a.e., then $g_n \rightarrow g$ effectively a.e..*

Further, in all cases the rates of convergence for the latter are computable from the former (in (2) use the modulus of continuity for g). Indeed, we do not need to assume the functions are effectively measurable, just that the rates of convergence are computable. The same results hold for continuous convergence, e.g. $f_r \rightarrow f$ as $r \rightarrow 0$.

Proof. (1): Assume $f_i \rightarrow f$ and $g_i \rightarrow g$ with rates $n(\varepsilon_1, \varepsilon_2)$ and $n'(\varepsilon_1, \varepsilon_2)$, respectively, of a.e.. convergence. I claim $m(\varepsilon_1, \varepsilon_2) = \max\left\{n\left(\frac{\varepsilon_1}{2}, \frac{\varepsilon_2}{2}\right), n'\left(\frac{\varepsilon_1}{2}, \frac{\varepsilon_2}{2}\right)\right\}$ is a rate of almost uniform convergence for $f_i + g_i \rightarrow f + g$. Indeed, if $m = m(\varepsilon_1, \varepsilon_2)$ then

$$\begin{aligned} \mu \left\{ \sup_{i \geq m} |(f_i + g_i) - (f + g)| > \varepsilon_1 \right\} &\leq \mu \left\{ \left(\sup_{i \geq m} |f_i - f| \right) + \left(\sup_{i \geq m} |g_i - g| \right) > \varepsilon_1 \right\} \\ &\leq \mu \left\{ \sup_{i \geq m} |f_i - f| > \frac{\varepsilon_1}{2} \right\} + \mu \left\{ \sup_{i \geq m} |g_i - g| > \frac{\varepsilon_1}{2} \right\} \leq \frac{\varepsilon_2}{2} + \frac{\varepsilon_2}{2} = \varepsilon_2. \end{aligned}$$

(2): Assume $f_i^j \rightarrow f^j$ with a rate of a.e. convergence $n_j(\varepsilon_1, \varepsilon_2)$. Also assume $g: \mathbb{R}^k \rightarrow \mathbb{R}$ is a continuous function with a computable modulus of continuity $\delta(\varepsilon)$, that is for all $x_0, \dots, x_{k-1}, y_0, \dots, y_{k-1} \in \mathbb{R}$.

$$\sum_{j=0}^{k-1} |x_j - y_j| \leq \delta(\varepsilon) \quad \rightarrow \quad |g(x_1, \dots, x_j) - g(y_1, \dots, y_k)| \leq \varepsilon.$$

Fix $\varepsilon_1, \varepsilon_2 > 0$. Let $m = \max_{j < k} n_j(\frac{\delta(\varepsilon_1)}{k}, \frac{\varepsilon_2}{k})$. Then

$$\begin{aligned} & \mu \left\{ \sup_{i \geq m} |g(f_i^0, \dots, f_i^{k-1}) - g(f^0, \dots, f^{k-1})| > \varepsilon_1 \right\} \\ & \leq \mu \left\{ \sum_{j < k} \sup_{i \geq m} |f_i^j - f^j| > \delta(\varepsilon_1) \right\} \\ & \leq \sum_{j < k} \mu \left\{ \sup_{i \geq m} |f_i^j - f^j| > \frac{\delta(\varepsilon_1)}{k} \right\} \\ & \leq \sum_{j < k} \frac{\varepsilon_2}{k} = \varepsilon_2. \end{aligned}$$

(3): Assume $f_n \leq g_n \leq h_n$ a.e. and $f_i \rightarrow g$ and $h_i \rightarrow g$ with computable rates of a.e. convergence. Let a rate of a.e. convergence for $f_i \rightarrow g$ be $n(\varepsilon_1, \varepsilon_2)$. By part (2), a rate of a.e. convergence $n'(\varepsilon_1, \varepsilon_2)$ for $(h_i - f_i) \rightarrow 0$ is computable. We claim that $g_i \rightarrow g$ with a rate of $m(\varepsilon_1, \varepsilon_2) = \max \{n(\frac{\varepsilon_1}{2}, \frac{\varepsilon_2}{2}), n'(\frac{\varepsilon_1}{2}, \frac{\varepsilon_2}{2})\}$. Indeed, for $n = n(\varepsilon_1, \varepsilon_2)$, we have

$$\begin{aligned} & \mu \left\{ \sup_{i \geq m} |g_i - g| > \varepsilon_1 \right\} \leq \mu \left\{ \sup_{i \geq m} (|f_i - g| + (h_i - f_i)) > \varepsilon_1 \right\} \\ & \leq \mu \left\{ \sup_{i \geq m} |f_i - g| > \frac{\varepsilon_1}{2} \right\} + \mu \left\{ \sup_{i \geq m} (h_i - f_i) > \frac{\varepsilon_1}{2} \right\} \leq \frac{\varepsilon_2}{2} + \frac{\varepsilon_2}{2} = \varepsilon_2. \end{aligned}$$

As for continuous convergence, the proofs are the same. \square

A.4. Convergence on Schnorr randomness.

Remark A.4. A SOLOVAY TEST FOR SCHNORR RANDOMNESS (U_n) is a computable sequence of effectively open sets U_n such that the sum $\sum_n \mu(U_n)$ is finite and computable. (This follows when $\mu(U_n)$ is computable uniformly from n and $\mu(U_n) \leq 2^{-n}$ or any other sequence with a finite sum.) If $x \in U_n$ for infinitely-many n , then say n is SOLOVAY COVERED by (U_n) . Then x is Schnorr random if and only if it is not Solovay-covered by any Solovay test for Schnorr randomness [11, 21]. (This is an effective version of the Borel-Cantelli lemma.)

Lemma A.5. *Suppose (φ_n) is a computable sequence of test functions which converge effectively a.e. to $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$.*

- (1) *(Existence) The limit $\lim_{n \rightarrow \infty} \varphi_n(x)$ exists on all Schnorr randoms x .*
- (2) *(Uniqueness) Given another sequence of test functions (ψ_n) converging effectively a.e. to f ,*

$$\lim_{n \rightarrow \infty} \varphi_n(x) = \lim_{n \rightarrow \infty} \psi_n(x) \quad (\text{on Schnorr random } x).$$

Proof. First we show existence by showing that $\varphi_n(x)$ is Cauchy for Schnorr randoms x . The main idea is to break up the indices into finite intervals. Since the rate of effectively a.e. convergence is computable, effectively choose n_k so that

$$\mu \left\{ \sup_{n \geq n_k} d_{\mathbb{Y}}(\varphi_n, \varphi_{n_k}) > 2^{-(k+1)} \right\} \leq 2^{-(k+1)}.$$

Our Solovay test for Schnorr randomness is

$$U_k = \left\{ x \mid \max_{n \in [n_k, n_{k+1}]} d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_k}(x)) > 2^{-(k+1)} \right\}.$$

Each set is effectively open uniformly in k . (As a technicality, let x only range over the interiors of the cells in φ_n . This guarantees that U_k is effectively open. It is also sufficient for our purposes since the boundary of each cell is a measure zero effectively closed set and therefore cannot contain Schnorr randoms.) This is a Solovay test since $\mu(U_k)$ is computable and by our choice of n_k ,

$$\mu(U_k) \leq \mu \left\{ \sup_{n \geq n_k} d_{\mathbb{Y}}(\varphi_n, \varphi_{n_k}) > 2^{-(k+1)} \right\} \leq 2^{-(k+1)}.$$

Now, let x be Schnorr random (and hence is not on the boundary of any cell). We have that x is in at most finitely many U_k . Hence for some k_0 large enough, for all $k \geq k_0$ and all $n \in [n_k, n_{k+1}]$ we have $d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_k}(x)) \leq 2^{-(k+1)}$. It follows that for all $k \geq k_0$ and for all $n \geq n_k$ that

$$d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_k}(x)) \leq \sum_{j \geq k} 2^{-(j+1)} \leq 2^{-k}.$$

Hence $\varphi_n(x)$ is Cauchy.

For uniqueness, take (φ_n) and (ψ_n) and interleave them, $\varphi_0, \psi_0, \varphi_1, \psi_1, \dots$. It is easy to see this sequence still has an effectively rate of a.e. convergence. Hence it converges on Schnorr randoms and each subsequence must converge to the same value. \square

Restatement of Proposition 3.18. *Suppose $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable with Cauchy-name (φ_n) (in the metric d_{meas} , L^1 -norm, or L^2 -norm).*

- (1) (Existence) *The limit $\lim_{n \rightarrow \infty} \varphi_n(x)$ exists on all Schnorr randoms x .*
- (2) (Uniqueness) *Given another Cauchy-name (ψ_n) for f ,*

$$\lim_{n \rightarrow \infty} \varphi_n(x) = \lim_{n \rightarrow \infty} \psi_n(x) \quad (\text{on Schnorr random } x).$$

Proof. A Cauchy-name has an effective rate of a.e. convergence by Proposition 3.15 and the rest follows from Lemma A.5. \square

Restatement of Lemma 3.19 (Convergence Lemma). *Suppose that (f_k) and f are uniformly effectively measurable. If*

$$f_k \rightarrow f \quad (\text{effectively a.e.})$$

then

$$\tilde{f}_k(x) \rightarrow \tilde{f}(x) \quad (\text{for all Schnorr random } x).$$

Proof. First, I will approximate (f_k) with a sequence of test functions (ψ_k) which converges effectively a.e. to f , and then show that (\tilde{f}_k) is close to (ψ_k) on Schnorr randoms.

For each k , let $(\varphi_n^k)_{n \in \mathbb{N}}$ be a Cauchy-name for f_k . Since a rate of a.e. convergence of $(\varphi_n^k)_{n \in \mathbb{N}}$ is computable from k , effectively choose $(n_{k,i})_{k,i \in \mathbb{N}}$ so that

$$\mu \left\{ \sup_{n \geq n_{k,i}} d_{\mathbb{Y}}(\varphi_n^k, \varphi_{n_{k,i}}^k) > 2^{-(k+i+1)} \right\} \leq 2^{-(k+i+1)}.$$

Consider the sequence $\psi_k = \varphi_{n_{k,0}}^k$. I will show that $\psi_k \xrightarrow[k \rightarrow \infty]{} f$ effectively a.e. as follows. Choose ε and δ . Since $f_k \rightarrow f$ effectively a.e., we can effectively choose k' such that

$$\mu \left\{ \sup_{k \geq k'} d_{\mathbb{Y}}(f_k, f) > \frac{\varepsilon}{2} \right\} \leq \frac{\delta}{2}.$$

Let $k(\varepsilon, \delta) = \max\{-2 \log_2 \varepsilon, -2 \log_2 \delta, k'\}$. Then $\sum_{k \geq k(\varepsilon, \delta)} 2^{-(k+1)} = 2^{-k(\varepsilon, \delta)} \leq \min\{\varepsilon/2, \delta/2\}$

$$\begin{aligned} \mu \left\{ \sup_{k \geq k(\varepsilon, \delta)} d_{\mathbb{Y}}(\varphi_{n_{k,0}}^k, f) > \varepsilon \right\} &\leq \sum_{k \geq k(\varepsilon, \delta)} \mu \left\{ d_{\mathbb{Y}}(\varphi_{n_{k,0}}^k, f_k) > 2^{-(k+1)} \right\} + \mu \left\{ \sup_{k \geq k'} d_{\mathbb{Y}}(f_k, f) > \frac{\varepsilon}{2} \right\} \\ &\leq \sum_{k \geq k(\varepsilon, \delta)} 2^{-(k+1)} + \frac{\delta}{2} \leq \delta. \end{aligned}$$

Hence, $\psi_k(x) \rightarrow \tilde{f}(x)$ on Schnorr randoms.

To show convergence of \tilde{f}_k , consider the Solovay test

$$U_{k,i} = \left\{ x \mid \max_{n \in [n_{k,i}, n_{k,i+1}]} d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_{k,i}}(x)) > 2^{-(k+i+1)} \right\}.$$

(Again, as in Lemma A.5, use the convention that x only ranges over the interiors of the cells.) This is a Solovay test since each $\mu(U_{k,i})$ is computable from k, i and since

$$\sum_k \sum_i \mu(U_{k,i}) \leq \sum_k \sum_i 2^{-(k+i+1)} = 2.$$

Now, let x be Schnorr random (and hence not on the boundary of any cell). We have that x is in at most finitely many $U_{k,i}$. Hence for some k_0 large enough, for all $k \geq k_0$, for all $i \geq 0$, and for all $n \in [n_{k,i}, n_{k,i+1}]$ we have $d_{\mathbb{Y}}(\varphi_n^k(x), \varphi_{n_{k,i}}^k(x)) \leq 2^{-(k+i+1)}$. It follows that for all $k \geq k_0$ and for all $n \geq n_{k,0}$ that

$$d_{\mathbb{Y}}(\varphi_n^k(x), \psi_k(x)) = d_{\mathbb{Y}}(\varphi_n^k(x), \varphi_{n_{k,0}}^k(x)) \leq \sum_{i \geq 0} 2^{-(k+i+1)} \leq 2^{-k}.$$

Hence $d_{\mathbb{Y}}(\tilde{f}_k(x), \psi_k(x)) \leq 2^{-k}$. Therefore, $\lim_k \tilde{f}_k(x) = \lim_k \psi_k(x) = \tilde{f}(x)$. \square

A.5. Properties of effectively measurable functions.

Restatement of Proposition 3.20. *The following implications hold for real-valued functions (and all the computations are uniform).*

- (1) $f \in L_{\text{comp}}^2 \Rightarrow f \in L_{\text{comp}}^1 \Rightarrow f \in L_{\text{comp}}^0$. (The converses do not hold in general.)
- (2) If $0 \leq f \leq 1$, then $f \in L_{\text{comp}}^2 \Leftrightarrow f \in L_{\text{comp}}^1 \Leftrightarrow f \in L_{\text{comp}}^0$.
- (3) $f \in L_{\text{comp}}^1 \Leftrightarrow (f \in L_{\text{comp}}^0 \text{ and } \|f\|_{L^1} \text{ is computable})$.
- (4) $f \in L_{\text{comp}}^2 \Leftrightarrow (f \in L_{\text{comp}}^0 \text{ and } \|f\|_{L^2} \text{ is computable})$.
- (5) If $f \in L_{\text{comp}}^1$ then $\int f d\mu$ is computable.
- (6) If B is effectively measurable, then $\mu(B)$ is computable.
- (7) If $0 \leq g \leq 1$, $g \in L_{\text{comp}}^1$, and $f \in L_{\text{comp}}^1$, then $g \cdot f \in L_{\text{comp}}^1$.

Proof. (1): Use that $\|f - \varphi\|_{L^2} \geq \|f - \varphi\|_{L^1} \geq d_{\text{meas}}(f, \varphi)$.

(2): In this case, $\|f - \varphi\|_{L^2}^2 \leq \|f - \varphi\|_{L^1} = d_{\text{meas}}(f, \varphi) \leq \|f - \varphi\|_{L^2}$.

(3): Given f effectively measurable, break up $\max\{f, 0\} = \sum_{n \in \mathbb{N}} f_n$ where $f_n = \min\{\max\{f, n\}, n+1\} - n$ and similarly for $\min\{-f, 0\}$. By (2), f_n is L^1 -computable from n . Use $\|f\|_{L^1}$ to approximate f in L^1 with finite sums of (f_n) .

(4): Same as (4).

(5): Use $\int f d\mu = \|\max\{f, 0\}\|_{L^1} + \|\min\{f, 0\}\|_{L^1}$ and that L^1 is a computable lattice.

(6): Use $\mu(B) = \mu(B \triangle \emptyset) = d(B, \emptyset)$ and that \emptyset is effectively measurable.

(7): Use that $g \in L_{\text{comp}}^1$ by (2) and

$$\begin{aligned} \|g \cdot f - \psi \cdot \varphi\|_{L^1} &\leq \|g \cdot (f - \varphi)\|_{L^1} + \|(g - \psi) \cdot \varphi\|_{L^1} \\ &\leq \|f - \varphi\|_{L^1} + \|g - \psi\|_{L^1} \cdot \|\varphi\|_{\infty}. \end{aligned}$$

Approximate f with a test function φ and then approximate g with ψ . \square

Restatement of Proposition 3.21 (Effective Lusin's theorem). *Given an effectively measurable $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$, and some rational $\varepsilon \geq 0$, there is an effectively closed set K of computable measure $\mu(K) \geq 1 - \varepsilon$ and a computable function $g: K \rightarrow \mathbb{Y}$ such that $g = \tilde{f} \upharpoonright K$ on Schnorr randoms. (Further, g and K are computable uniformly from ε and any name for f .) Moreover, if $\mathbb{Y} = \mathbb{R}$, then $g: K \rightarrow \mathbb{Y}$ can be extended (uniformly from its name) to a total computable function $g: \mathbb{X} \rightarrow \mathbb{Y}$ such that $g = \tilde{f} \upharpoonright K$ on Schnorr randoms.*

Proof. Let (φ_n) be the Cauchy-name for f in the metric d_{meas} . Let (U_k) be the Solovay test for Schnorr randomness from Lemma A.5, that is

$$U_k = \{x \mid \max_{n \in [n_k, n_{k+1}]} d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_k}(x)) > 2^{-k}\}$$

for some computable sequence (n_k) . Again, we ignore the boundaries of the cells corresponding to φ_n for $n \in [n_k, n_{k+1}]$. Recall, $\mu(U_k)$ is computable from k and $\mu(U_k) \leq 2^{-k}$. To handle the boundaries, we can find an effectively open set V_k of computable measure $\mu(V_k) \leq 2^{-k}$ such that V_k covers the boundaries of the cells corresponding to φ_n for $n \in [n_k, n_{k+1}]$.

Let

$$K = \left(\bigcup_{k \geq 2 - \log_2 \varepsilon} U_k \cup V_k \right)^c.$$

Then

$$1 - \mu(K) \leq \sum_{k \geq 2 - \log_2 \varepsilon} \mu(U_k \cup V_k) \leq \sum_{k \geq 2 - \log_2 \varepsilon} 2 \cdot 2^{-k} \leq \varepsilon$$

and $\mu(K)$ is computable (the measure of every finite union is computable, and the measure of the remaining tail can be made arbitrarily small).

As in the proof of Lemma A.5, it follows that for all $x \in K$ and all $k \geq 2 - \log_2 \varepsilon$, that $x \in U_k$ and is not on the boundaries of the relevant cells. Therefore

$$d_{\mathbb{Y}}(\varphi_n(x), \varphi_{n_k}(x)) \leq \sum_{j \geq k} 2^{-(j+1)} \leq 2^{-k}.$$

Use this to compute the value of $g(x) := \lim_n \varphi_n(x)$ for $x \in K$. If x is Schnorr random this is equal to $\tilde{f}(x)$.

If $\mathbb{Y} = \mathbb{R}$, then by the effective Tietze extension theorem [53], we can extend g to a total computable function. \square

Restatement of Proposition 3.22 (Effective inner/outer regularity). *Given $A \subseteq (\mathbb{X}, \mu)$ effectively measurable, and some rational $\varepsilon > 0$, there is an effectively open set U and an effectively closed set C both of computable measure such that $C \subseteq \tilde{A} \subseteq U$ for Schnorr randoms such that $\mu(U) - \mu(C) \leq \varepsilon$. (The sets U, C and their measures $\mu(U), \mu(C)$ are uniformly computable from ε and any name for A .)*

Proof. From the effective Lusin's theorem (Proposition 3.21), we can choose an effectively closed K of computable measure $\mu(K) \geq 1 - \varepsilon$ and a computable function $g: K \rightarrow \{0, 1\}$ such that $\tilde{1}_A \upharpoonright C = g$ on Schnorr randoms. Then let $C = \{x \in K \mid g(x) = 1\}$ and $U = \mathbb{X} \setminus \{x \in K \mid g(x) = 0\}$. These are effectively closed and open. Then $C \subseteq \tilde{A} \subseteq U$ for Schnorr randoms since $\tilde{A} \upharpoonright C = g$, and $\mu(U) - \mu(C) = 1 - \mu(K) \leq \varepsilon$.

The measures $\mu(C)$ and $\mu(U)$ are computable as follows. From a name for g , we can enumerate a sequence of balls $\{B_i^0\}$ and $\{B_j^1\}$ from $\text{Basis}(\mathbb{X}, \mu)$ such that if $x \in B_i^0$ and $x \in K$ then $f(x) = 0$ and similarly for B_j^1 . Notice $\bigcup_i B_i^0 \cup \bigcup_j B_j^1$ covers K .

Let $V = K^c$ and enumerate a sequence of balls $\{A_i\}$ from $\text{Basis}(\mathbb{X}, \mu)$ such that $V = \bigcup_i A_i$ and hence $\mathbb{X} = \bigcup_i B_i^0 \cup \bigcup_j B_j^1 \cup \bigcup_k A_k$. Find a finite subsequence of these balls such that

$$\mu(B_1^0 \cup \dots \cup B_\ell^0 \cup B_1^1 \cup \dots \cup B_n^1 \cup A_1 \cup \dots \cup A_m) \approx 1.$$

Then $\mu(C) \approx \mu((B_1^1 \cup \dots \cup B_n^1) \setminus (A_1 \cup \dots \cup A_m))$ and $\mu(U) \approx 1 - \mu((B_1^0 \cup \dots \cup B_\ell^0) \setminus (A_1 \cup \dots \cup A_m))$. \square

Restatement of Proposition 3.23 (Schnorr layerwise computability). *Consider a (pointwise-defined) measurable function $f: \mathbb{X} \rightarrow \mathbb{Y}$ that is SCHNORR LAYERWISE COMPUTABLE, that is, there is a computable sequence (C_n) of effectively closed sets of computable measure $\mu(C_n) \leq 2^{-n}$, such that $f \upharpoonright C_n$ is computable on C_n uniformly in n . Then there is an effectively measurable $g: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ such that $\tilde{g} = f$ on Schnorr randoms.*

Proof. Fix $\varepsilon > 0$. Choose C_n such that $\mu(C_n) \geq 1 - \varepsilon$. From a name for $f \upharpoonright C_n$, we can enumerate a sequence of balls $\{B_i\}_i$ from $\text{Basis}(\mathbb{X}, \mu)$ and values c_i for which if $x \in B_i$ and $x \in C_n$ then $d_{\mathbb{Y}}(f(x), c_i) \leq \varepsilon$. Note that $\{B_i\}_i$ covers C_n , so we can compute a subsequence B_0, \dots, B_{k-1} such that $\mu(B_0 \cup \dots \cup B_{k-1}) \geq 1 - 2\varepsilon$.

Let φ be the test function made from all cells of B_0, \dots, B_{k-1} (except the cell $B_0^c \cup \dots \cup B_{k-1}^c$). Use the approximations c_i to determine the value of φ on each cell. Then $d_{\mathbb{Y}}(\varphi(x), f(x)) \leq \varepsilon$ unless $x \notin C_n$ or $x \notin B_0 \cup \dots \cup B_{k-1}$. Therefore,

$$d_{\text{meas}}(\varphi, f) \leq \varepsilon + (1 - \mu(C_n)) + (1 - \mu(B_0 \cup \dots \cup B_{k-1})) \leq 4\varepsilon.$$

Hence, f is almost-everywhere equal to an effectively measurable function g with Cauchy name φ . Moreover, $\varphi_n(x) \rightarrow f(x)$ for all x in all but finitely-many C_n . This is true of all Schnorr randoms x , since (C_n^c) forms a Solovay test for Schnorr randomness. \square

Restatement of Proposition 3.24 (Examples of effectively measurable functions and sets). *All of these functions $f: \mathbb{X} \rightarrow \mathbb{Y}$ and sets $A \subseteq \mathbb{X}$ are effectively measurable, and $\tilde{f} = f$ and $\tilde{A} = A$ on Schnorr randoms.*

- (1) Test functions and test sets as in Propositions 3.1 and 3.3 and in Definition 3.8.
- (2) Computable functions and decidable sets (i.e., computable 0,1-valued functions).
- (3) Almost-everywhere computable functions $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and almost-everywhere decidable sets (i.e., almost everywhere computable 0,1-valued functions).
- (4) Nonnegative lower semicomputable functions $f: \mathbb{X} \rightarrow \mathbb{R}$ with a computable integral, effectively open sets $U \subseteq \mathbb{X}$ of computable measure, and effectively closed sets $C \subseteq \mathbb{X}$ of computable measure.

Proof. (1): This is obvious from the definition of effectively measurable and of \tilde{f} .

(2): See (3).

(3): We will show that almost-everywhere computable functions are Schnorr layerwise computable. From a name for f from n , we can enumerate a sequence of balls $\{B_i^n\}_i$ from $\text{Basis}(\mathbb{X}, \mu)$ and values c_i^n for which if $x \in B_i^n$ then $d_{\mathbb{Y}}(f(x), c_i^n) \leq 2^{-n}$. Moreover, $\mu(\bigcup_i B_i^n) = 1$. Choose ε . For each n , find a subsequence $(B_0^n, \dots, B_{k(n)-1}^n)$ such that $\mu(B_0^n \cup \dots \cup B_{k(n)-1}^n) \geq 1 - \varepsilon/2^n$. Then let $C_\varepsilon = \bigcap_n (\overline{B}_0^n \cup \dots \cup \overline{B}_{k(n)-1}^n)$.

I will show that C_ε is an effectively closed set of computable measure $\mu(C_\varepsilon) \geq 1 - 2\varepsilon$ such that f is computable on C_ε . It is clearly effectively closed. It has computable measure since $\mu(\bigcap_{n \leq m} (\overline{B}_0^n \cup \dots \cup \overline{B}_{k(n)-1}^n))$ is computable and

$$\mu(C_\varepsilon) - \mu\left(\bigcap_{m > n} (\overline{B}_0^m \cup \dots \cup \overline{B}_{k(m)-1}^m)\right) \leq \sum_{m > n} (1 - \mu(\overline{B}_0^m \cup \dots \cup \overline{B}_{k(m)-1}^m)) \leq \sum_{m > n} \varepsilon/2^m = \varepsilon/2^n.$$

Similarly, $1 - \mu(C_\varepsilon) \leq \sum_n (1 - \mu(\overline{B}_0^n \cup \dots \cup \overline{B}_{k(n)-1}^n)) \leq \sum_n \varepsilon/2^n \leq 2\varepsilon$. Finally, f is computable on C_ε since for any n and x in C_ε we can wait until $x \in B_i^n$ for some i , and we know that $f(x)$ is within 2^{-n} of c_i^n .

(4): Let $f = \sup g_n$ where (g_n) is a computable sequence of computable functions. Then $\|f - g_n\|_{L^1} = \int f - g_n d\mu$ and from monotonicity we can compute an effective rate of a.e. convergence of g_n to f . Therefore f is effectively measurable and $\tilde{f} = \lim_n \tilde{g}_n = \lim_n g_n = f$. For effectively open U of computable measure, just use $f = 1_U$ which is lower semi computable. The same for effectively closed C of computable measure. \square

Restatement of Proposition 3.25 (Push-forward measures). *If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable, then the push-forward measure $(\mathbb{Y}, \mu_* f)$ is a computable probability space (uniformly from (\mathbb{X}, μ) , \mathbb{Y} , and f).*

Proof. It is enough to compute $\int \varphi d\mu_* f = \int \varphi \circ f d\mu$ uniformly from a computable function $\varphi: \mathbb{Y} \rightarrow [0, 1]$. By the effective Lusin's theorem (Proposition 3.21) \tilde{f} is Schnorr layerwise computable. Since φ is a computable function, we have that $\varphi \circ \tilde{f}$ is Schnorr layerwise (since from the definition of Schnorr layerwise computable, the composition of a computable function with a Schnorr layerwise computable function is still Schnorr layerwise computable). By Proposition 3.23, $\varphi \circ f$ is effectively measurable. Since $\varphi \circ f$ is effectively measurable and bounded, the integral $\int \varphi \circ f d\mu$ is computable (Proposition 3.20). \square

Restatement of Proposition 3.26 (Preservation of Schnorr randomness). *If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable and x is Schnorr random, then $\tilde{f}(x)$ is Schnorr random on $(\mathbb{Y}, \mu_* f)$.*

Proof. Assume not. Let (U_n) be a $(\mathbb{Y}, \mu_* f)$ -Schnorr test which covers $\tilde{f}(x)$. Let $g = \sum_n 1_{U_n}$. Then g is a lower semicomputable function and hence $g = \sup_n \varphi_n$ for a computable sequence of computable functions. We can also assume that $0 \leq \varphi_n \leq n$. By the same argument as in the previous proof, $\varphi_n \circ f$ is effectively measurable uniformly in n and $\varphi_n \circ f = \varphi_n \circ \tilde{f}$ on (\mathbb{X}, μ) -Schnorr randoms x .

Moreover, we can show that $\varphi_n \circ f \rightarrow g \circ f$ effectively in measure since $d_{\text{meas}}(\varphi_n \circ f, g \circ f) = d_{\text{meas}}(\varphi_n, g) \leq \|g - \varphi_n\|_{L^1}$ which is computable since $\int g d\mu_* f = \sum_n \mu_* f(U_n)$ is computable and $\int \varphi_n d\mu_* f$ is computable since φ_n is computable and bounded. Restricting to a subsequence (n_k) we have that $\varphi_{n_k} \circ f \rightarrow g \circ f$ converges effectively a.e. By Lemma 3.19, $(\varphi_{n_k} \circ \tilde{f})(x)$ must converge (to something in \mathbb{R}) since x is Schnorr random. However, $\lim_k (\varphi_{n_k} \circ \tilde{f})(x) = \infty \notin \mathbb{R}$. \square

Restatement of Proposition 3.27 (Composition and tuples).

- (1) (Composition) Given $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and $g: (\mathbb{Y}, \mu_* f) \rightarrow \mathbb{Z}$ effectively measurable, the composition $g \circ f$ is effectively measurable (uniformly from f and g) and

$$\widetilde{g \circ f} = \tilde{g} \circ \tilde{f} \quad (\text{on Schnorr randoms}).$$

- (2) (Tuples) Given $f_n: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}_n$ effectively measurable (uniformly in n), the tuples

$$(f_0, \dots, f_{k-1}): (\mathbb{X}, \mu) \rightarrow \mathbb{Y}_0 \times \dots \times \mathbb{Y}_{k-1}$$

and

$$(f_n)_{n \in \mathbb{N}}: (\mathbb{X}, \mu) \rightarrow \prod_{n \in \mathbb{N}} \mathbb{Y}_n$$

are effectively measurable (uniformly from (f_n)) and

$$(f_0, \dots, f_{k-1}) = (\widetilde{f_0}, \dots, \widetilde{f_{k-1}}) \quad \text{and} \quad (\widetilde{f_i})_{i \in \mathbb{N}} = (\widetilde{f_i})_{i \in \mathbb{N}} \quad (\text{on Schnorr randoms}).$$

Proof. (1): Consider f and g with Cauchy-names, (φ_n) and (ψ_n) . First we show that $\psi_n \circ f$ is effectively measurable uniformly in f and ψ_n . Fix $\varepsilon > 0$. We can effectively choose some small $\varepsilon' > 0$ such that all but a small μ -measure of x are more than ε' from the boundary of the cells which make up ψ_n . Then choose φ_m such that $\mu\{d_{\mathbb{Y}}(\varphi_m, f) > \varepsilon'\} < \varepsilon$. Outside of this bad set, we have the $\varphi_m(x)$ and $f(x)$ are in the same cell of $\psi_n(x)$, and hence $\psi_n \circ \varphi_m - \psi_n \circ f = 0$. Hence $d_{\text{meas}}(\varphi_m, f) \leq \varepsilon$. Therefore, $\psi_n \circ \varphi_m \xrightarrow{m \rightarrow \infty} \psi_n \circ f$ effectively a.e. and therefore $\psi_n \circ f$ is effectively measurable uniformly from f and ψ_n . (This required that $\psi_n \circ \varphi_m$ is a test function. To ensure this, one may need to slightly modify φ_m to avoid hitting the boundary of the cells in ψ_n .) By Lemma 3.19, $\widetilde{\psi_n \circ f} = \lim_m \widetilde{\psi_n \circ \varphi_m} = \widetilde{\psi_n \circ \widetilde{f}}$ on Schnorr randoms.

Next, we show that $g \circ f$ is effectively measurable uniformly in f and g . This is straightforward since $d_{\text{meas}}(g \circ f, \psi_n \circ f) = d_{\text{meas}}(g, \psi_n)$. Moreover, by Lemma 3.19, $\widetilde{g \circ f} = \lim_n \widetilde{\psi_n \circ f} = \lim_n \widetilde{\psi_n \circ \widetilde{f}} = \widetilde{g \circ \widetilde{f}}$ on Schnorr randoms x (since $\widetilde{f}(x)$ is Schnorr random).

(2): I just do the infinite case. Let $\mathbb{Y} = \prod_{n \in \mathbb{N}} \mathbb{Y}_n$ with metric $d_{\mathbb{Y}} = \sum_{n \in \mathbb{N}} 2^{-(n+1)} \min\{d_{\mathbb{Y}_n}, 1\}$. For each n , let $(\varphi_n^j)_{j \in \mathbb{N}}$ be the Cauchy-name for f_n in the metric d_{meas} . Then approximate $f = (f_n)_{n \in \mathbb{N}}$ with $\psi_k = (\varphi_n^{2^{k+n+2}})_{n \in \mathbb{N}}$. Then

$$d_{\text{meas}}(f, \psi_k) = \int d_{\mathbb{Y}}(f, \psi_k) d\mu = \sum_{n \in \mathbb{N}} 2^{-(n+1)} \int \min\{d_{\mathbb{Y}_n}(f_n, \varphi_n^{2^{k+n+2}}), 1\} d\mu \leq 2^{-k}.$$

Therefore f is effectively measurable and $(\widetilde{f_i})_{i \in \mathbb{N}} = \lim_{j \rightarrow \infty} (\varphi_i^j)_{i \in \mathbb{N}} = (\widetilde{f_i})_{i \in \mathbb{N}}$. \square

Restatement of Proposition 3.28 (Combinations of measurable functions).

- (1) (*Computable pointwise operations*). All computable pointwise operations, including vector, lattice, and Boolean algebra operations preserve effective measurability. Moreover, given $f, g: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ and $A, B \subseteq (\mathbb{X}, \mu)$ effectively measurable, we have

$$\begin{aligned} \widetilde{f + g} &= \widetilde{f} + \widetilde{g}, & \widetilde{af} &= a\widetilde{f}, & \widetilde{f \cdot g} &= \widetilde{f} \cdot \widetilde{g} \\ \widetilde{\min(f, g)} &= \min(\widetilde{f}, \widetilde{g}), & \widetilde{\max(f, g)} &= \max(\widetilde{f}, \widetilde{g}), & \widetilde{|f|} &= |\widetilde{f}| \\ \widetilde{A \cup B} &= \widetilde{A} \cup \widetilde{B}, & \widetilde{A \cap B} &= \widetilde{A} \cap \widetilde{B}, & \widetilde{A^c} &= \widetilde{A}^c, & \widetilde{\mathbb{X}} &= \mathbb{X}, & \widetilde{\emptyset} &= \emptyset \end{aligned}$$

on Schnorr randoms, and

$$f \leq g \text{ a.e.} \iff \widetilde{f} \leq \widetilde{g} \text{ (on Schnorr randoms)}$$

$$A \subseteq B \text{ a.e.} \iff \widetilde{A} \subseteq \widetilde{B} \text{ (on Schnorr randoms)}.$$

- (2) (*Inverse image*) Given $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ and $B \subseteq (\mathbb{Y}, \mu_*)$ effectively measurable then $f^{-1}(B)$ is effectively measurable and $\widetilde{f^{-1}(B)} = \widetilde{f}^{-1}(\widetilde{B})$ on Schnorr randoms.
- (3) (*Rotations*) Given $f: (\mathbb{T}^d, \lambda) \rightarrow \mathbb{R}$ effectively measurable, and a computable vector $t \in \mathbb{T}^d$, then $h(x) := f(x - t)$ is effectively measurable and $\widetilde{h}(x) = \widetilde{f}(x - t)$ on Schnorr randoms.
- (4) (*Indicator functions*) Given $A \subseteq (\mathbb{X}, \mu)$, A is effectively measurable if and only if $\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ is effectively measurable (equivalently, L^1 -computable by Proposition 3.20 (2)) and $x \in \widetilde{A}$ if and only if $\widetilde{\mathbf{1}_A}(x) = 1$ on Schnorr randoms. (Notice the codomain of $\mathbf{1}_A$ is \mathbb{R} here rather than $\{0, 1\}$ as in Definition 3.17.)

Proof. (1): This is a direct application of Propositions 3.24 and 3.27. Also if $f \leq g$ a.e., then $g - f = \max\{g - f, 0\}$ a.e., and $\widetilde{g} - \widetilde{f} = \widetilde{g - f} = \max\{\widetilde{g - f}, 0\} = \max\{\widetilde{g} - \widetilde{f}, 0\} \geq 0$ on Schnorr randoms. Similarly for $A \subseteq B$.

(2): Use that $\mathbf{1}_{f^{-1}(B)} = \mathbf{1}_B \circ f$. The rest follows from Proposition 3.27.

(3): Let $g(x) := x - t$. Then g is computable and measure preserving, that is $\lambda_* g = \lambda$. Hence, $h = f \circ g$, and $\widetilde{h} = \widetilde{f} \circ g$ by Propositions 3.24 and 3.27.

(4): Consider the computable inclusion map $i: \{0, 1\} \rightarrow \mathbb{R}$. We have $(\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R}) = i \circ (\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \{0, 1\})$. By Proposition 3.27, if A is effectively measurable then $\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ is. For the other direction,

if $\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R}$ is effectively measurable, then consider the partial computable map $g: \mathbb{R} \rightarrow \{0, 1\}$ which sends $0 \mapsto 0$ and $1 \mapsto 1$. This map is almost-everywhere computable on $(\mathbb{R}, \mu_* \mathbf{1}_A)$ (which only has mass on 0 and 1). Now, $(\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \{0, 1\}) = g \circ (\mathbf{1}_A: (\mathbb{X}, \mu) \rightarrow \mathbb{R})$. The rest follows from Proposition 3.27. \square

Restatement of Proposition 3.29. *The following implications hold for real-valued functions (and all the computations are uniform).*

- (1) *If $f \in L^1_{\text{comp}}$ and A is effectively measurable, then $\int_A f d\mu$ is computable.*
- (2) *If \mathbb{X} is effectively compact (see [36])—as is $[0, 1]^d$, \mathbb{T}^d , and $2^{\mathbb{N}}$ —and $g: \mathbb{X} \rightarrow \mathbb{R}$ is computable, then g is L^1 -computable (since it has computable bounds).*
- (3) *If $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ is effectively measurable and $g \in L^1_{\text{comp}}(\mathbb{Y}, \mu_* f)$ (resp. $L^2_{\text{comp}}(\mathbb{Y}, \mu_* f)$), then $g \circ f \in L^1_{\text{comp}}(\mathbb{X}, \mu)$ (resp. $L^2_{\text{comp}}(\mathbb{X}, \mu)$).*

Proof. (1): By Proposition 3.28 (4), $\mathbf{1}_A \in L^1_{\text{comp}}$. Then use 3.20 (7).

(2): By Proposition 3.24, g is effectively measurable. Since \mathbb{X} is effectively compact, $\max_{x \in \mathbb{X}} g(x)$ and $\min_{x \in \mathbb{X}} g(x)$ are computable from g [36]. Now apply Proposition 3.20 (2).

(3): By Proposition 3.27 $g \circ f \in L^0_{\text{comp}}$, and moreover $\|g \circ f\|_{L^1(\mathbb{X}, \mu)} = \|g\|_{L^1(\mathbb{Y}, \mu_* f)}$ (similarly for L^2). Apply Proposition 3.20 (3). \square

Restatement of Proposition 3.30. *Given a measurable map $f: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$, the following are equivalent.*

- (1) *f is effectively measurable.*
- (2) *The push-forward measure $(\mathbb{Y}, \mu_* f)$ is computable and one (or all) of the following “pull-back” maps are computable:*
 - (a) *(L^1 functions) $g \in L^1(\mathbb{Y}, \mu_* f) \mapsto g \circ f \in L^1(\mathbb{X}, \mu)$.*
 - (b) *(L^2 functions) $g \in L^2(\mathbb{Y}, \mu_* f) \mapsto g \circ f \in L^2(\mathbb{X}, \mu)$.*
 - (c) *(Measurable sets) $B \subseteq (\mathbb{Y}, \mu_* f) \mapsto f^{-1}(B) \subseteq (\mathbb{X}, \mu)$.*

Proof. (1) \Rightarrow (2) follows from Propositions 3.25, 3.27, 3.28, and 3.29.

For the other direction, assume (2)(a) or (2)(b). Then $B \mapsto \mathbf{1}_B \mapsto \mathbf{1}_B \circ f \mapsto f^{-1}(B)$ is a chain of computable operators (using Proposition 3.28 (4)), and therefore (2)(c) holds.

For (2)(c) \Rightarrow (1), fix $\varepsilon > 0$. Since $(\mathbb{Y}, \mu_* f)$ is computable, effectively choose finitely many balls B_0, \dots, B_{k-1} of radius at most $\varepsilon/2$ from $\text{Basis}(\mathbb{Y}, \mu_* f)$ such that $\mu_* f(B_0 \cup \dots \cup B_{k-1}) \geq 1 - \varepsilon/2$. Let C_0, \dots, C_{2^k-1} be the cells formed by combining the elements of B_0, \dots, B_{k-1} . Let C_0 denote the cell $B_0^c \cap \dots \cap B_{k-1}^c$ which is the only cell without a diameter bounded by $\varepsilon/2$. For $i \geq 1$, effectively choose a point y_i inside the cell C_i (by choosing the center of the lowest indexed ball B_j for which $C_i \subseteq B_j$). Let $A_i = f^{-1}(C_i)$. By assumption, these are effectively measurable. Define $\varphi: (\mathbb{X}, \mu) \rightarrow \mathbb{Y}$ as the effectively measurable function which has value y_i on A_i for $i \geq 1$ and 0 otherwise. Notice on A_i ($1 \leq i \leq 2^k - 1$), that φ and f both take values in C_i .

$$\begin{aligned} d_{\text{meas}}(f, \varphi) &= \int \max\{d_{\mathbb{Y}}(f, \varphi), 1\} d\mu \\ &\leq 1 \cdot \mu(A_0) + \sum_{i=1}^{2^k-1} \int_{A_i} d_{\mathbb{Y}}(f, \varphi) d\mu \\ &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \cdot \sum_{i=1}^{2^k-1} \mu(A_i) \leq \varepsilon. \end{aligned}$$

Hence f is effectively measurable. \square

REFERENCES

- [1] Jeremy Avigad, Edward T. Dean, and Jason Rute. Algorithmic randomness, reverse mathematics, and the dominated convergence theorem. *Ann. Pure Appl. Logic*, 163(12):1854–1864, 2012.
- [2] Jeremy Avigad, Philipp Gerhardy, and Henry Towsner. Local stability of ergodic averages. *Trans. Amer. Math. Soc.*, 362(1):261–288, 2010.
- [3] Laurent Bienvenu, Adam R. Day, Mathieu Hoyrup, Ilya Mezhirov, and Alexander Shen. A constructive version of Birkhoff’s ergodic theorem for Martin-Löf random points. *Inform. and Comput.*, 210:21–30, 2012.
- [4] Errett Bishop. *Foundations of constructive analysis*. McGraw-Hill Book Co., New York, 1967.
- [5] Errett Bishop and Douglas Bridges. *Constructive analysis*. 279. Springer-Verlag, 1985.

- [6] Volker Bosserhoff. Notions of probabilistic computability on represented spaces. *J. Universal Computer Science*, 14(6):956–995, 2008.
- [7] Vasco Brattka, Peter Hertling, and Klaus Weihrauch. A tutorial on computable analysis. In *New computational paradigms*, pages 425–491. Springer, New York, 2008.
- [8] Vasco Brattka, Joseph S. Miller, and André Nies. Randomness and differentiability. *Submitted*.
- [9] Claude Dellacherie and Paul-André Meyer. *Probabilities and potential B: Theory of martingales*, volume 72 of *North-Holland Mathematics Studies*. North-Holland Publishing Co., Amsterdam, 1982. Translated from the French by J. P. Wilson.
- [10] O. Demut. The differentiability of constructive functions of weakly bounded variation on pseudo numbers. *Comment. Math. Univ. Carolinae*, 16(3):583–599, 1975.
- [11] Rodney G. Downey and Evan J. Griffiths. Schnorr randomness. *J. Symbolic Logic*, 69(2):533–554, 2004.
- [12] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic randomness and complexity*. Theory and Applications of Computability. Springer, New York, 2010.
- [13] R. M. Dudley. *Real analysis and probability*, volume 74 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2002. Revised reprint of the 1989 original.
- [14] Rick Durrett. *Probability: theory and examples*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, fourth edition, 2010.
- [15] Abbas Edalat. A computable approach to measure and integration theory. *Inform. and Comput.*, 207(5):642–659, 2009.
- [16] Johanna N. Y. Franklin, Noam Greenberg, Joseph S. Miller, and Keng Meng Ng. Martin-Löf random points satisfy Birkhoff’s ergodic theorem for effectively closed sets. *Proc. Amer. Math. Soc.*, 140(10):3623–3628, 2012.
- [17] Johanna N.Y. Franklin and Henry Towsner. Randomness and non-ergodic systems. *Submitted*.
- [18] Cameron E. Freer, Bjørn Kjos-Hanssen, André Nies, and Frank Stephan. Algorithmic aspects of Lipschitz functions. *Submitted*.
- [19] Cameron E. Freer and Daniel M. Roy. Computable de Finetti measures. *Annals of Pure and Applied Logic*, 2011.
- [20] Peter Gács. Uniform test of algorithmic randomness over a general space. *Theoret. Comput. Sci.*, 341(1-3):91–137, 2005.
- [21] Peter Gács, Mathieu Hoyrup, and Cristóbal Rojas. Randomness on computable probability spaces—a dynamical point of view. *Theory Comput. Syst.*, 48(3):465–485, 2011.
- [22] Stefano Galatolo, Mathieu Hoyrup, and Cristóbal Rojas. Computing the speed of convergence of ergodic averages and pseudorandom points in computable dynamical systems. In Xizhong Zheng and Ning Zhong, editors, *Proceedings Seventh International Conference on Computability and Complexity in Analysis*, Zhenjiang, China, 21–25th June 2010, volume 24 of *Electronic Proceedings in Theoretical Computer Science*, pages 7–18. Open Publishing Association, 2010.
- [23] Mathieu Hoyrup. Computability of the ergodic decomposition. *Annals of Pure and Applied Logic*, 164(5):542–549, 2013. Computability in Europe 2011.
- [24] Mathieu Hoyrup and Cristóbal Rojas. An application of Martin-Löf randomness to effective probability theory. In *Mathematical theory and computational practice*, volume 5635 of *Lecture Notes in Comput. Sci.*, pages 260–269. Springer, Berlin, 2009.
- [25] Mathieu Hoyrup and Cristóbal Rojas. Applications of effective probability theory to Martin-Löf randomness. In *Automata, languages and programming. Part I*, volume 5555 of *Lecture Notes in Comput. Sci.*, pages 549–561. Springer, Berlin, 2009.
- [26] Mathieu Hoyrup and Cristóbal Rojas. Computability of probability measures and Martin-Löf randomness over metric spaces. *Inform. and Comput.*, 207(7):830–847, 2009.
- [27] Mathieu Hoyrup and Cristóbal Rojas. Absolute continuity of measures and preservation of randomness. *Preprint*.
- [28] Mathieu Hoyrup, Cristóbal Rojas, and Klaus Weihrauch. Computability of the Radon-Nikodym derivative. In *Models of computation in context*, volume 6735 of *Lecture Notes in Comput. Sci.*, pages 132–141. Springer, Heidelberg, 2011.
- [29] Tahereh Jafarikhah, Massoud Pourmahdian, and Klaus Weihrauch. Computable Jordan decomposition of linear continuous functionals on $C[0; 1]$. In Mathieu Hoyrup, Ker-I Ko, Robert Rettinger, and Ning Zhong, editors, *CCA 2013 Tenth International Conference on Computability and Complexity in Analysis*, volume 367-7/2013, pages 48–59. Informatik Berichte FernUniversität in Hagen, 2013.
- [30] Yitzhak Katznelson. *An introduction to harmonic analysis*. John Wiley & Sons Inc., New York, 1968.
- [31] Ker-I Ko and Harvey Friedman. Computational complexity of real functions. *Theoret. Comput. Sci.*, 20(3):323–352, 1982.
- [32] Hong Lu and Klaus Weihrauch. Computable Riesz representation for the dual of $C[0; 1]$. *MLQ Math. Log. Q.*, 53(4-5):415–430, 2007.
- [33] Wolfgang Merkle, Nenad Mihailović, and Theodore A. Slaman. Some results on effective randomness. *Theory Comput. Syst.*, 39(5):707–721, 2006.
- [34] Kenshi Miyabe. L^1 -computability, layerwise computability and Solovay reducibility. *Computability*. To appear.
- [35] Michał Morayne and Sławomir Solecki. Martingale proof of the existence of Lebesgue points. *Real Anal. Exchange*, 15(1):401–406, 1989/90.
- [36] Takakazu Mori, Yoshiki Tsujii, and Mariko Yasugi. Computability structures on metric spaces. In *Combinatorics, complexity, & logic (Auckland, 1996)*, Springer Ser. Discrete Math. Theor. Comput. Sci., pages 351–362. Springer, Singapore, 1997.
- [37] Satyadev Nandakumar. An effective ergodic theorem and some applications. In *STOC’08*, pages 39–44. ACM, New York, 2008.
- [38] André Nies. *Computability and randomness*, volume 51 of *Oxford Logic Guides*. Oxford University Press, Oxford, 2009.
- [39] Noopur Pathak. A computational aspect of the Lebesgue differentiation theorem. *J. Log. Anal.*, 1(9):15, 2009.
- [40] Noopur Pathak, Cristóbal Rojas, and Stephen G. Simpson. Schnorr randomness and the Lebesgue differentiation theorem. *Proceedings of the American Mathematical Society*, To appear.

- [41] Marian B. Pour-El and J. Ian Richards. *Computability in analysis and physics*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1989.
- [42] Jason Rute. Computable randomness and betting for computable probability spaces. *Submitted*. arXiv:1203.5535.
- [43] N. A. Šanin. *Constructive real numbers and constructive function spaces*. Translated from the Russian by E. Mendelson. Translations of Mathematical Monographs, Vol. 21. American Mathematical Society, Providence, R.I., 1968.
- [44] Claus-Peter Schnorr. *Zufälligkeit und Wahrscheinlichkeit. Eine algorithmische Begründung der Wahrscheinlichkeitstheorie*. Lecture Notes in Mathematics, Vol. 218. Springer-Verlag, Berlin, 1971.
- [45] Matthias Schröder. Admissible representations for probability measures. *MLQ Math. Log. Q.*, 53(4-5):431–445, 2007.
- [46] Robert I. Soare. *Recursively enumerable sets and degrees*. Perspectives in Mathematical Logic. Springer-Verlag, Berlin, 1987. A study of computable functions and computably generated sets.
- [47] Hayato Takahashi. Bayesian approach to a definition of random sequences with respect to parametric models. In M.J. Dinneen, editor, *Proc. of IEEE ISOC ITW2005 on Coding and Complexity*, pages 217–220, 2005.
- [48] Terence Tao. *An introduction to measure theory*, volume 126 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [49] Michael E. Taylor. *Measure theory and integration*, volume 76 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2006.
- [50] V. V. V'yugin. Ergodic theorems for individual random sequences. *Theoret. Comput. Sci.*, 207(2):343–361, 1998.
- [51] Peter Walters. *An introduction to ergodic theory*, volume 79 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1982.
- [52] Klaus Weihrauch. *Computable analysis: An introduction*. Texts in Theoretical Computer Science. An EATCS Series. Springer-Verlag, Berlin, 2000.
- [53] Klaus Weihrauch. On computable metric spaces Tietze-Urysohn extension is computable. In *Computability and complexity in analysis (Swansea, 2000)*, volume 2064 of *Lecture Notes in Comput. Sci.*, pages 357–368. Springer, Berlin, 2001.
- [54] David Williams. *Probability with martingales*. Cambridge Mathematical Textbooks. Cambridge University Press, Cambridge, 1991.
- [55] Yongcheng Wu and Decheng Ding. Computability of measurable sets via effective topologies. *Arch. Math. Logic*, 45(3):365–379, 2006.
- [56] Yongcheng Wu and Klaus Weihrauch. A computable version of the Daniell-Stone theorem on integration and linear functionals. *Theoret. Comput. Sci.*, 359(1-3):28–42, 2006.
- [57] Xiaokang Yu. *Measure theory in weak subsystems of second-order arithmetic*. ProQuest LLC, Ann Arbor, MI, 1987. Thesis (Ph.D.)—The Pennsylvania State University.
- [58] Xiaokang Yu. Lebesgue convergence theorems and reverse mathematics. *Math. Logic Quart.*, 40(1):1–13, 1994.
- [59] Xiaokang Yu and Stephen G. Simpson. Measure theory and weak König's lemma. *Arch. Math. Logic*, 30(3):171–180, 1990.